

ON THE NUMBER OF REPRESENTATIONS OF INTEGERS BY CERTAIN QUADRATIC FORMS

SHAUN COOPER

(Received 1 October 2007)

Abstract

Generating functions are used to derive formulas for the number of representations of a positive integer by each of the quadratic forms $x_1^2 + x_2^2 + x_3^2 + 2x_4^2$, $x_1^2 + 2x_2^2 + 2x_3^2 + 2x_4^2$, $x_1^2 + x_2^2 + 2x_3^2 + 4x_4^2$ and $x_1^2 + 2x_2^2 + 4x_3^2 + 4x_4^2$. The formulas show that the number of representations by each form is always positive. Some of the analogous results involving sums of triangular numbers are also given.

2000 *Mathematics subject classification*: primary 11E25; secondary 05A19, 11D85, 33E05.

Keywords and phrases: sum of squares, sum of triangular numbers, quadratic form, theta function, elliptic function, Lambert series, Lagrange's four squares theorem.

1. Introduction

Lagrange's four-squares theorem says that every positive integer may be expressed as a sum of four squares. Jacobi went further and used elliptic functions to prove that the number of solutions of

$$x_1^2 + x_2^2 + x_3^2 + x_4^2 = n$$

in integers is

$$8 \sum_{\substack{d|n \\ d \not\equiv 0 \pmod{4}}} d.$$

Since $d = 1$ is a divisor of any positive integer n , the sum in Jacobi's theorem is positive and Lagrange's theorem follows immediately.

Fine [6, pp. 74–76] proved an analogue of Jacobi's theorem for the quadratic form $x_1^2 + x_2^2 + x_3^2 + 2x_4^2$, and showed that the number of representations of any positive integer by this form is always positive. The purpose of this paper is to present a short and simple proof of Fine's result, and to obtain analogous results for the quadratic forms $x_1^2 + 2x_2^2 + 2x_3^2 + 2x_4^2$, $x_1^2 + x_2^2 + 2x_3^2 + 4x_4^2$ and $x_1^2 + 2x_2^2 + 4x_3^2 + 4x_4^2$. The proof depends on simple properties of theta functions and a well-known formula which expresses a difference of two Weierstrass elliptic functions as an infinite product.

2. Notation and lemmas

Let q be a complex number that satisfies $|q| < 1$. Ramanujan's theta functions $\varphi(q)$ and $\psi(q)$ are defined by

$$\varphi(q) = \sum_{n=-\infty}^{\infty} q^{n^2},$$

$$\psi(q) = \sum_{n=0}^{\infty} q^{n(n+1)/2}.$$

Some of their basic properties are summarized in the following three lemmas.

LEMMA 2.1.

$$\varphi(q) + \varphi(-q) = 2\varphi(q^4), \quad (2.1)$$

$$\varphi(q) - \varphi(-q) = 4q\psi(q^8), \quad (2.2)$$

$$\varphi(q)^2 + \varphi(-q)^2 = 2\varphi(q^2)^2, \quad (2.3)$$

$$\varphi(q)^2 - \varphi(-q)^2 = 8q\psi(q^4)^2. \quad (2.4)$$

PROOF. These can be proved using series manipulations. A complete proof is given in [3, pp. 40–41]. \square

LEMMA 2.2.

$$\varphi(-q) = \prod_{j=1}^{\infty} \frac{(1 - q^j)^2}{(1 - q^{2j})}, \quad (2.5)$$

$$\psi(q) = \prod_{j=1}^{\infty} \frac{(1 - q^{2j})^2}{(1 - q^j)}. \quad (2.6)$$

PROOF. These are consequences of the Jacobi triple product identity. See [3, pp. 36–37] for proofs. \square

LEMMA 2.3.

$$\varphi(-q^2)^2 = \varphi(q)\varphi(-q), \quad (2.7)$$

$$\psi(q)^2 = \varphi(q)\psi(q^2). \quad (2.8)$$

PROOF. These are consequences of Lemma 2.2. See [3, p. 40] for proofs. \square

The final lemma in this section, and the main tool for this work, is a classical result that expresses a quotient of Weierstrass sigma functions as a difference of two Weierstrass elliptic functions.

LEMMA 2.4. *Let x, z and q be any complex numbers satisfying $|q| < |x|, |z| < |q|^{-1}$, and $x, z \neq 1$. Then*

$$\begin{aligned} & \frac{(1-xz)(x-z)}{(1-x)^2(1-z)^2} \\ & \times \prod_{j=1}^{\infty} \frac{(1-xzq^j)(1-q^jx/z)(1-q^jz/x)(1-q^j/(xz))(1-q^j)^4}{(1-xq^j)^2(1-zq^j)^2(1-q^j/x)^2(1-q^j/z)^2} \\ & = \frac{x}{(1-x)^2} - \frac{z}{(1-z)^2} + \sum_{j=1}^{\infty} \frac{jq^j}{1-q^j} (x^j + x^{-j} - z^j - z^{-j}). \end{aligned}$$

PROOF. There are many proofs of this result in the literature. For a simple one, see [5]. For an equivalent statement in terms of the Weierstrass functions, see [7, p. 451, Exercise 1]. For generalizations, see [4]. □

We shall use the following special values of Jacobi’s symbol:

$$\left(\frac{2}{n}\right) = \begin{cases} 1 & \text{if } n \equiv 1 \text{ or } 7 \pmod{8}; \\ -1 & \text{if } n \equiv 3 \text{ or } 5 \pmod{8}; \\ 0 & \text{otherwise;} \end{cases}$$

where n is a positive integer. The Jacobi symbol is multiplicative, that is,

$$\left(\frac{2}{mn}\right) = \left(\frac{2}{m}\right)\left(\frac{2}{n}\right)$$

for all positive integers m and n .

3. Theta function identities

The results in this section all hinge on the following result.

THEOREM 3.1.

$$\begin{aligned} \varphi(-q)\varphi(-q^2)\varphi(-q^4)^2 &= 1 - 2 \sum_{j=1}^{\infty} \binom{2}{j} \frac{jq^j}{1-q^j}, \\ q\psi(q)^2\psi(q^2)\psi(q^4) &= \sum_{j=1}^{\infty} \frac{j(q^j - q^{3j} - q^{5j} + q^{7j})}{1-q^{8j}}. \end{aligned}$$

PROOF. Let $\omega = \exp(i\pi/4)$ and take $x = \omega^3, z = \omega$, in Lemma 2.4 to get

$$\begin{aligned} & \frac{2(\omega^3 - \omega)}{(1-\omega)^2(1-\omega^3)^2} \prod_{j=1}^{\infty} \frac{(1+q^j)^2(1+q^{2j})(1-q^j)^4}{(1-\omega q^j)^2(1-\omega^3 q^j)^2(1-\omega^5 q^j)^2(1-\omega^7 q^j)^2} \\ & = \frac{\omega^3}{(1-\omega^3)^2} - \frac{\omega}{(1-\omega)^2} + \sum_{j=1}^{\infty} \frac{jq^j}{1-q^j} (\omega^{3j} + \omega^{-3j} - \omega^j - \omega^{-j}). \quad (3.1) \end{aligned}$$

By straightforward calculations,

$$\frac{2(\omega^3 - \omega)}{(1 - \omega)^2(1 - \omega^3)^2} = \frac{\omega^3}{(1 - \omega^3)^2} - \frac{\omega}{(1 - \omega)^2} = \sqrt{2}.$$

Next, using the algebraic rearrangements

$$1 + x = \frac{1 - x^2}{1 - x}$$

and

$$(1 - \omega x)(1 - \omega^3 x)(1 - \omega^5 x)(1 - \omega^7 x) = \frac{1 - x^8}{1 - x^4},$$

the left-hand side of (3.1) simplifies to

$$\sqrt{2} \prod_{j=1}^{\infty} \frac{(1 - q^j)^2(1 - q^{2j})(1 - q^{4j})^3}{(1 - q^{8j})^2},$$

which, by (2.5), equals

$$\sqrt{2}\varphi(-q)\varphi(-q^2)\varphi(-q^4)^2. \tag{3.2}$$

Furthermore, by checking the residue classes modulo 8, we find that

$$\omega^{3j} + \omega^{-3j} - \omega^j - \omega^{-j} = -2\sqrt{2}\left(\frac{2}{j}\right).$$

Therefore, the right-hand side of (3.1) simplifies to

$$\sqrt{2} - 2\sqrt{2} \sum_{j=1}^{\infty} \left(\frac{2}{j}\right) \frac{jq^j}{1 - q^j}. \tag{3.3}$$

Combining (3.2) and (3.3), we complete the proof of the first result.

Now let us prove the second result. In Lemma 2.4, replace (q, x, z) with (q^8, q, q^3) to get

$$\begin{aligned} & q \prod_{j=1}^{\infty} \frac{(1 - q^{8j-4})^2(1 - q^{8j-6})(1 - q^{8j-2})(1 - q^{8j})^4}{(1 - q^{8j-7})^2(1 - q^{8j-5})^2(1 - q^{8j-3})^2(1 - q^{8j-1})^2} \\ &= \frac{q}{(1 - q)^2} - \frac{q^3}{1 - q^3} + \sum_{j=1}^{\infty} \frac{jq^{8j}}{1 - q^{8j}}(q^j + q^{-j} - q^{3j} - q^{-3j}). \end{aligned} \tag{3.4}$$

The left-hand side of (3.4) simplifies to

$$q \prod_{j=1}^{\infty} \frac{(1 - q^{2j})^3(1 - q^{4j})(1 - q^{8j})^2}{(1 - q^j)^2}$$

which, by (2.6), equals

$$q\psi(q)^2\psi(q^2)\psi(q^4). \tag{3.5}$$

The right-hand side of (3.4) is

$$\begin{aligned} & \sum_{j=1}^{\infty} j(q^j - q^{3j}) + \sum_{j=1}^{\infty} \sum_{k=1}^{\infty} j\{q^{(8k+1)j} + q^{(8k-1)j} - q^{(8k+3)j} - q^{(8k-3)j}\} \\ &= \sum_{j=1}^{\infty} \sum_{k=1}^{\infty} j\{q^{(8k-7)j} + q^{(8k-1)j} - q^{(8k-5)j} - q^{(8k-3)j}\} \\ &= \sum_{j=1}^{\infty} \frac{j(q^j - q^{3j} - q^{5j} + q^{7j})}{1 - q^{8j}}. \end{aligned} \tag{3.6}$$

Combining (3.5) and (3.6), we complete the proof of the second result. □

The next theorem gives Lambert series expansions for four different products of theta functions.

THEOREM 3.2. *Let*

$$f_1(q) = 1 - 2 \sum_{j=1}^{\infty} \binom{2}{j} \frac{jq^j}{1 - q^j} \quad \text{and} \quad f_2(q) = \sum_{j=1}^{\infty} \frac{j(q^j - q^{3j} - q^{5j} + q^{7j})}{1 - q^{8j}}.$$

Then

$$\begin{aligned} \varphi(q)^3\varphi(q^2) &= f_1(q) + 8f_2(q), \\ \varphi(q)\varphi(q^2)^3 &= f_1(q) + 4f_2(q), \\ \varphi(q)^2\varphi(q^2)\varphi(q^4) &= f_1(q^2) + 4f_2(q), \\ \varphi(q)\varphi(q^2)\varphi(q^4)^2 &= f_1(q^2) + 2f_2(q). \end{aligned}$$

PROOF. By (2.4) followed by two applications of each of (2.7) and (2.8),

$$\begin{aligned} \varphi(q)^3\varphi(q^2) &= \varphi(q)\varphi(q^2)\{\varphi(-q)^2 + 8q\psi(q^4)^2\} \\ &= \varphi(-q)\varphi(-q^2)^2\varphi(q^2) + 8q\varphi(q)\psi(q^2)^2\psi(q^4) \\ &= \varphi(-q)\varphi(-q^2)\varphi(-q^4)^2 + 8q\psi(q)^2\psi(q^2)\psi(q^4). \end{aligned} \tag{3.7}$$

Similarly, by (2.3) and (2.4), followed by two applications of each of (2.7) and (2.8),

$$\begin{aligned} \varphi(q)\varphi(q^2)^3 &= \varphi(q)\varphi(q^2)\{\varphi(-q)^2 + 4q\psi(q^4)^2\} \\ &= \varphi(-q)\varphi(-q^2)^2\varphi(q^2) + 4q\varphi(q)\psi(q^2)^2\psi(q^4) \\ &= \varphi(-q)\varphi(-q^2)\varphi(-q^4)^2 + 4q\psi(q)^2\psi(q^2)\psi(q^4). \end{aligned} \tag{3.8}$$

By (2.2) followed by three applications of each of (2.7) and (2.8),

$$\begin{aligned}\varphi(q)^2\varphi(q^2)\varphi(q^4) &= \varphi(q)\varphi(q^2)\varphi(q^4)\{\varphi(-q) + 4q\psi(q^8)\} \\ &= \varphi(-q^2)^2\varphi(q^2)\varphi(q^4) + 4q\varphi(q)\varphi(q^2)\psi(q^4)^2 \\ &= \varphi(-q^2)\varphi(-q^4)^2\varphi(q^4) + 4q\varphi(q)\psi(q^2)^2\psi(q^4) \\ &= \varphi(-q^2)\varphi(-q^4)\varphi(-q^8)^2 + 4q\psi(q)^2\psi(q^2)\psi(q^4).\end{aligned}\quad (3.9)$$

Similarly, by (2.1) and (2.2), followed by three applications of each of (2.7) and (2.8), we have (omitting some details which are similar to the above)

$$\begin{aligned}\varphi(q)\varphi(q^2)\varphi(q^4)^2 &= \varphi(q)\varphi(q^2)\varphi(q^4)\{\varphi(-q) + 2q\psi(q^8)\} \\ &= \varphi(-q^2)\varphi(-q^4)\varphi(-q^8)^2 + 2q\psi(q)^2\psi(q^2)\psi(q^4).\end{aligned}\quad (3.10)$$

Substituting the results of Theorem 3.1 into the right-hand sides of (3.7)–(3.10), we complete the proof. \square

Results involving the function $\psi(q)$ (and sometimes $\varphi(q)$ as well) can be obtained by dissecting one of the results in Theorem 3.1. These are given in the following result.

THEOREM 3.3.

$$\begin{aligned}q\psi(q^8)\varphi(q^8)^3 &= \sum_{n=0}^{\infty} \left(\sum_{d|8n+1} \left(\frac{2}{d} \right) d \right) q^{8n+1}, \\ q^3\psi(q^8)^3\varphi(q^8) &= -\frac{1}{2} \sum_{n=0}^{\infty} \left(\sum_{d|8n+3} \left(\frac{2}{d} \right) d \right) q^{8n+3}, \\ q^5\psi(q^8)^3\psi(q^{16}) &= -\frac{1}{4} \sum_{n=0}^{\infty} \left(\sum_{d|8n+5} \left(\frac{2}{d} \right) d \right) q^{8n+5}, \\ q^7\psi(q^8)\psi(q^{16})^3 &= \frac{1}{8} \sum_{n=0}^{\infty} \left(\sum_{d|8n+7} \left(\frac{2}{d} \right) d \right) q^{8n+7}.\end{aligned}$$

PROOF. If $f(q) = \sum_{n=0}^{\infty} c(n)q^n$, let us define $[q; a, b]f(q)$ by

$$[q; a, b]f(q) = \sum_{n=0}^{\infty} c(an + b)q^{an+b}.$$

By Lemma 2.1,

$$\begin{aligned}\varphi(-q)\varphi(-q^2)\varphi(-q^4)^2 \\ = \{\varphi(q^4) - 2q\psi(q^8)\}\{\varphi(q^8) - 2q^2\psi(q^{16})\}\{\varphi(q^8)^2 - 4q^4\psi(q^{16})^2\}.\end{aligned}$$

If we expand and extract terms of the form q^{8n+1} , q^{8n+3} , q^{8n+5} and q^{8n+7} , we obtain

$$[q; 8, 1]\varphi(-q)\varphi(-q^2)\varphi(-q^4)^2 = -2q\psi(q^8)\varphi(q^8)^3, \tag{3.11}$$

$$\begin{aligned} [q; 8, 3]\varphi(-q)\varphi(-q^2)\varphi(-q^4)^2 &= 4q^3\psi(q^8)\psi(q^{16})\varphi(q^8)^2 \\ &= 4q^3\psi(q^8)^3\varphi(q^8) \quad \text{by (2.8),} \end{aligned} \tag{3.12}$$

$$\begin{aligned} [q; 8, 5]\varphi(-q)\varphi(-q^2)\varphi(-q^4)^2 &= 8q^5\psi(q^8)\varphi(q^8)\psi(q^{16})^2 \\ &= 8q^5\psi(q^8)^3\psi(q^{16}) \quad \text{by (2.8)} \end{aligned} \tag{3.13}$$

and

$$[q; 8, 7]\varphi(-q)\varphi(-q^2)\varphi(-q^4)^2 = -16q^7\psi(q^8)\psi(q^{16})^3, \tag{3.14}$$

respectively. Next, by the first part of Theorem 3.1,

$$\varphi(-q)\varphi(-q^2)\varphi(-q^4)^2 = 1 - 2 \sum_{n=1}^{\infty} \left(\sum_{d|n} \left(\frac{2}{d} \right) d \right) q^n. \tag{3.15}$$

Therefore, for $1 \leq k \leq 8$,

$$[q; 8, k]\varphi(-q)\varphi(-q^2)\varphi(-q^4)^2 = -2 \sum_{n=0}^{\infty} \left(\sum_{d|8n+k} \left(\frac{2}{d} \right) d \right) q^{8n+k}. \tag{3.16}$$

Using (3.16) in (3.11)–(3.14) we complete the proof. □

4. Combinatorial interpretations

In this section, we give combinatorial interpretations of the results in Theorems 3.1–3.3. We begin with a lemma.

LEMMA 4.1. *Let n be a positive integer and write $n = 2^\alpha m$, where m is odd and α is a nonnegative integer. Then:*

- (1) *the coefficient of q^n in $\varphi(-q)\varphi(-q^2)\varphi(-q^4)^2$ is $-2 \sum_{d|m} (2/d)d$;*
- (2) *the coefficient of q^n in $q\psi(q)^2\psi(q^2)\psi(q^4)$ is $2^\alpha (2/m) \sum_{d|m} (2/d)d$.*

PROOF. The first result is immediate from (3.15), together with the fact that $(2/d) = 0$ if d is even. To prove the second result, observe that from Theorem 3.1,

$$q\psi(q)^2\psi(q^2)\psi(q^4) = \sum_{j=1}^{\infty} \frac{j(q^j - q^{3j} - q^{5j} + q^{7j})}{1 - q^{8j}} = \sum_{j=1}^{\infty} \sum_{k=1}^{\infty} \left(\frac{2}{k} \right) j q^{jk}.$$

The coefficient of q^n is therefore

$$\begin{aligned} \sum_{jk=n} \binom{2}{k} j &= \sum_{\substack{jk=n \\ k \text{ odd}}} \binom{2}{k} j \\ &= \sum_{ik=m} \binom{2}{k} 2^\alpha i \\ &= 2^\alpha \binom{2}{m} \sum_{ik=m} \binom{2}{i} i \\ &= 2^\alpha \binom{2}{m} \sum_{d|m} \binom{2}{d} d, \end{aligned}$$

where the multiplicative property of the Jacobi symbol was used in the penultimate step. □

Let a, b, c and d be positive integers. Let $r_{(a,b,c,d)}(n)$ denote the number of solutions of $ax_1^2 + bx_2^2 + cx_3^2 + dx_4^2 = n$ in integers (positive, negative and/or zero). The generating function is

$$\sum_{n=0}^{\infty} r_{(a,b,c,d)}(n)q^n = \varphi(q^a)\varphi(q^b)\varphi(q^c)\varphi(q^d).$$

THEOREM 4.2. *If n is a positive integer, write $n = 2^\alpha m$, where m is odd and α is a nonnegative integer. Then*

$$\begin{aligned} r_{(2,1,1,1)}(n) &= \left\{ 2^{\alpha+3} \binom{2}{m} - 2 \right\} \sum_{d|m} \binom{2}{d} d, \\ r_{(2,2,2,1)}(n) &= \left\{ 2^{\alpha+2} \binom{2}{m} - 2 \right\} \sum_{d|m} \binom{2}{d} d, \\ r_{(4,2,1,1)}(n) &= \left\{ 2^{\alpha+2} \binom{2}{m} - 1 - (-1)^n \right\} \sum_{d|m} \binom{2}{d} d, \\ r_{(4,4,2,1)}(n) &= \left\{ 2^{\alpha+1} \binom{2}{m} - 1 - (-1)^n \right\} \sum_{d|m} \binom{2}{d} d. \end{aligned}$$

PROOF. Use Lemma 4.1 to extract the coefficient of q^n from each of the results in Theorem 3.2. □

THEOREM 4.3. *Let o_j and e_j denote an odd integer and an even integer, respectively. For a positive integer n , let $A(n), B(n), C(n), D(n)$ and $E(n)$ denote the number of*

solutions of

$$\begin{aligned} o_1^2 + 2e_2^2 + 2e_3^2 + 2e_4^2 &= n, \\ o_1^2 + o_2^2 + o_3^2 + 2e_4^2 &= n, \\ o_1^2 + o_2^2 + o_3^2 + 2o_4^2 &= n, \\ o_1^2 + 2o_2^2 + 2o_3^2 + 2o_4^2 &= n \end{aligned}$$

and

$$o_1^2 + o_2^2 + 2o_3^2 + 4o_4^2 = n,$$

respectively. Then

$$\sum_{d|n} \left(\frac{2}{d}\right) d = \begin{cases} \frac{1}{2}A(n) & \text{if } n \equiv 1 \pmod{8}, \\ -\frac{1}{4}B(n) & \text{if } n \equiv 3 \pmod{8}, \\ -\frac{1}{4}C(n) & \text{if } n \equiv 5 \pmod{8}, \\ \frac{1}{2}D(n) & \text{if } n \equiv 7 \pmod{8}. \end{cases} \tag{4.1}$$

Since $A(n)$, $B(n)$, $C(n)$ and $D(n)$ are zero unless $n \equiv 1, 3, 5$ or 7 , respectively, (4.1) completely determines the values of $A(n)$, $B(n)$, $C(n)$ and $D(n)$.

If we write $n = 2^\alpha m$, where m is odd and α is a nonnegative integer, then

$$E(n) = 2^{\alpha+1} \left(\frac{2}{m}\right) \sum_{d|n} \left(\frac{2}{d}\right) d \quad \text{if } n \equiv 0 \pmod{8}, \tag{4.2}$$

and $E(n) = 0$ otherwise.

PROOF. Observe that

$$\begin{aligned} \sum_{n=0}^{\infty} A(n)q^n &= \sum_{x_1, x_2, x_3, x_4=-\infty}^{\infty} q^{(2x_1+1)^2+2(2x_2)^2+2(2x_3)^2+2(2x_4)^2} \\ &= 2q\psi(q^8)\varphi(q^8)^3. \end{aligned} \tag{4.3}$$

Similarly, we find that

$$\sum_{n=0}^{\infty} B(n)q^n = 8q^3\psi(q^8)^3\varphi(q^8), \tag{4.4}$$

$$\sum_{n=0}^{\infty} C(n)q^n = 16q^5\psi(q^8)^3\psi(q^{16}), \tag{4.5}$$

$$\sum_{n=0}^{\infty} D(n)q^n = 16q^7\psi(q^8)\psi(q^{16})^3, \tag{4.6}$$

$$\sum_{n=0}^{\infty} E(n)q^n = 16q^8\psi(q^8)^2\psi(q^{16})\psi(q^{32}). \tag{4.7}$$

The identity (4.1) follows from these by comparing coefficients using Theorem 3.3, while (4.2) follows from Lemma 4.1. \square

The results for $C(n)$, $D(n)$ and $E(n)$ in Theorem 4.3 may be re-expressed in terms of sums of triangular numbers. Let a, b, c and d be positive integers, and let $t_{(a,b,c,d)}(n)$ denote the number of solutions in nonnegative integers of

$$\frac{ax_1(x_1 + 1)}{2} + \frac{bx_2(x_2 + 1)}{2} + \frac{cx_3(x_3 + 1)}{2} + \frac{dx_4(x_4 + 1)}{2} = n.$$

The generating function for $t_{(a,b,c,d)}(n)$ is

$$\sum_{n=0}^{\infty} t_{(a,b,c,d)}(n)q^n = \psi(q^a)\psi(q^b)\psi(q^c)\psi(q^d).$$

THEOREM 4.4. *Let n be a positive integer and write $n = 2^\alpha m$, where m is odd and α is a nonnegative integer. Then*

$$\begin{aligned} t_{(2,1,1,1)}(n) &= -\frac{1}{4} \sum_{d|8n+5} \left(\frac{2}{d}\right)d, \\ t_{(2,2,2,1)}(n) &= \frac{1}{8} \sum_{d|8n+7} \left(\frac{2}{d}\right)d, \\ t_{(4,2,1,1)}(n-1) &= 2^\alpha \left(\frac{2}{m}\right) \sum_{d|n} \left(\frac{2}{d}\right)d. \end{aligned}$$

PROOF. From the generating function for $t_{(2,1,1,1)}(n)$ and (4.5),

$$\sum_{n=0}^{\infty} t_{(2,1,1,1)}(n)q^{8n+5} = q^5\psi(q^8)^3\psi(q^{16}) = \frac{1}{16} \sum_{n=0}^{\infty} C(n)q^n.$$

Equating coefficients and applying Theorem 4.3, we find that

$$t_{(2,1,1,1)}(n) = \frac{1}{16}C(8n + 5) = -\frac{1}{4} \sum_{d|8n+5} \left(\frac{2}{d}\right)d.$$

The other two results may be proved similarly. \square

Analogues of Lagrange’s theorem can be deduced from Theorems 4.2 and 4.3.

THEOREM 4.5. *Let n be a nonnegative integer. Then $r_{(2,1,1,1)}(n)$, $r_{(2,2,2,1)}(n)$, $r_{(4,2,1,1)}(n)$ and $r_{(4,4,2,1)}(n)$ are all positive, as are $A(8n + 1)$, $B(8n + 3)$, $C(8n + 5)$, $D(8n + 7)$ and $E(8n + 8)$.*

PROOF. Let m be an odd integer, and let $m = \prod_p p^{\alpha_p}$ be its prime factorization. By geometric series,

$$\left(\frac{2}{m}\right) \sum_{d|m} \left(\frac{2}{d}\right) d = \prod_p \frac{p^{\alpha_p+1} - (2/p)^{\alpha_p+1}}{p - (2/p)},$$

which is clearly positive. The truth of the theorem follows from this, together with the results in Theorems 4.2 and 4.3. The trivial observation

$$r_{(2,1,1,1)}(0) = r_{(2,2,2,1)}(0) = r_{(4,2,1,1)}(0) = r_{(4,4,2,1)}(0) = 1$$

takes care of the case $n = 0$. □

REMARK 4.6. It is interesting to compare the results for $r_\lambda(n)$ with the corresponding results for $t_\lambda(n)$ in Theorem 4.4. We find that

$$r_{(2,1,1,1)}(8n + 5) = -10 \sum_{d|8n+5} \left(\frac{2}{d}\right) d = 40t_{(2,1,1,1)}(n)$$

and

$$r_{(2,2,2,1)}(8n + 7) = 2 \sum_{d|8n+7} \left(\frac{2}{d}\right) d = 16t_{(2,2,2,1)}(n).$$

These are instances of a theorem in [1], which says that $r_{(\lambda_1, \dots, \lambda_k)}(8n + \lambda_1 + \dots + \lambda_k)$ is a constant (independent of n) multiple of $t_{(\lambda_1, \dots, \lambda_k)}(n)$, provided $\lambda_1 + \dots + \lambda_k < 8$. Furthermore,

$$\begin{aligned} r_{(4,2,1,1)}(8n) - r_{(4,2,1,1)}(2n) &= (2^{\alpha+5} - 2^{\alpha+3}) \left(\frac{2}{m}\right) \sum_{d|m} \left(\frac{2}{d}\right) d \\ &= 24 \times 2^\alpha \left(\frac{2}{m}\right) \sum_{d|m} \left(\frac{2}{d}\right) d \\ &= 24t_{(4,2,1,1)}(n - 1), \end{aligned}$$

and this is an instance of a theorem in [2].

References

- [1] C. Adiga, S. Cooper and J. H. Han, ‘A general relation between sums of squares and sums of triangular numbers’, *Int. J. Number Theory* **1** (2005), 175–182.
- [2] N. D. Baruah, S. Cooper and M. Hirschhorn, ‘Sums of squares and sums of triangular numbers induced by partitions of 8’, *Int. J. Number Theory* to appear.
- [3] B. C. Berndt, *Ramanujan’s Notebooks, Part III* (Springer-Verlag, New York, 1991).
- [4] S. H. Chan, ‘Generalized Lambert series identities’, *Proc. London Math. Soc.* (3) **91** (2005), 598–622.
- [5] J. M. Dobbie, ‘A simple proof of some partition formulae of Ramanujan’s’, *Quart. J. Math. Oxford Ser.* (2) **6** (1955), 193–196.

- [6] N. J. Fine, *Basic Hypergeometric Series and Applications* (American Mathematical Society, Providence, RI, 1988).
- [7] E. T. Whittaker and G. N. Watson, *A Course of Modern Analysis*, 4th edn (Cambridge University Press, Cambridge, 1927).

SHAUN COOPER, Institute of Information and Mathematical Sciences,
Massey University, Private Bag 102904, North Shore Mail Centre,
Auckland, New Zealand
e-mail: s.cooper@massey.ac.nz