

INDEX FOR VOLUME 113

	PAGE
Armitage, D. H. & Nelson, C. S. A harmonic quadrature formula characterizing open strips	147
Armitage, D. H., Gardiner, S. J. & Netuka, I. Separation of points by classes of harmonic functions.	561
Auinger, K. The word problem for the bifree combinatorial strict regular semigroup	519
Baker, J. W. & Lau, A. T. Compact left ideal groups in semigroup compactification of locally compact groups	507
Baxter, M. Symmetry characterizations of certain distributions. 3. Discounted additive functionals and large deviations for a general finite-state Markov chain	381
Boston, N. & Ullom, S. V. Representations related to CM elliptic curves	71
Brown, B. M., Kirby, V. G., Evans, W. D. & Plum, M. Safe numerical bounds for the Titchmarsh–Weyl $m(\lambda)$ -function	583
Cariñena, J., Martínez, E. & Sarlet, W. Geometric characterization of separable second-order differential equations	205
Castillo, J. M. F. & Sánchez, F. Upper l_p -estimates in vector sequence spaces, with some applications	329
Dales, H. G. & Millington, A. Translation-invariant linear operators	161
Díaz, J. C. & Miñarro, M. A. On Fréchet Montel spaces and their projective tensor product	335
Donkin, S. Invariant functions on matrices	23
Drury, S. W. & Guo, K. Some remarks on the restriction of the Fourier transform to surfaces	153
Edmunds, D. E., Opic, B. & Pick, L. Poincaré and Friedrichs inequalities in abstract Sobolev spaces	355
Evans, M. J. T -systems of certain finite simple groups.	9
Evans, W. D., Plum, M., Brown, B. M. & Kirby, V. G. Safe numerical bounds for the Titchmarsh–Weyl $m(\lambda)$ -function	583
Farr, G. E. A generalization of the Whitney rank generating function	267
Gardiner, S. J., Netuka, I. & Armitage, D. H. Separation of points by classes of harmonic functions.	561
Gibson, C. G. & Hobbs, C. A. Simple singularities of space curves	297
Green, D. J. On the cohomology of the sporadic simple group J_4	253
Guo, K. & Drury, S. W. Some remarks on the restriction of the Fourier transform to surfaces	153
Hennings, M. A. Strongly spin-preserving solutions of the Yang–Baxter equation and their link variants	401
Higgins, P. M. Combinatorial results for semigroups of order-preserving mappings	281
Hinkkanen, A. Julia sets of rational are functions uniformly perfect	543
Hobbs, C. A. & Gibson, C. G. Simple singularities of space curves	297
Hobson, D. G. & Rogers, L. C. G. Recurrence and transience of reflecting Brownian motion in the quadrant	387
Hofmann, K. H. & Morris, S. A. Generators on the arc component of compact connected groups	479
Holland, D. & Wilson, S. M. J. Localization and class groups of module categories with exactness defects	233
Johnson, F. E. A. A rational invariant for certain infinite discrete groups	473

Index

Kamiya, S. On H -balls and canonical regions of loxodromic elements in complex hyperbolic space	573
Kesten, H. Convergence in distribution of lightly trimmed and untrimmed sums are equivalent	615
Kiesel, R. Power series methods and almost sure convergence	195
Kirby, V. G., Evans, W. D., Plum, M. & Brown, B. M. Safe numerical bounds for the Titchmarsh–Weyl $m(\lambda)$ -function	583
Lau, A. T. & Baker, D. W. Compact left ideal groups in semigroup compactification of locally compact groups	507
Maeda, H. A criterion for a smooth surface to be Del Pezzo	1
Martinez, E., Cariñena, J. & Sarlet, W. Geometric characterization of separable second-order differential equations	205
Majid, S. Transmutation theory and rank for quantum braided groups	45
Mbekhta, M. Semi-Fredholm perturbations and commutators	173
Millington, A. & Dales, H. G. Translation-invariant linear operators	161
Miñarro, M. A. & Díaz, J. C. On Fréchet Montel spaces and their projective tensor product	335
Morris, S. A. & Hofmann, K. H. Generators on the arc component of compact connected groups	479
Nelson, C. S. & Armitage, D. H. A harmonic quadrature formula characterizing open strips	147
Netuka, I., Armitage, D. H. & Gardiner, S. J. Separation of points by classes of harmonic functions	561
Opic, B., Pick, L. & Edmunds, D. E. Poincaré and Friedrichs inequalities in abstract Sobolev spaces	355
Oxley, J. & Whittle, G. On sign-representable matroids	499
Pan, Y. L^2 estimates for convolution operators with oscillating kernels	179
Pick, L., Edmunds, D. E. & Opic, B. Poincaré and Friedrichs inequalities in abstract Sobolev spaces	355
Plum, M., Brown, B. M., Kirby, V. G. & Evans, W. D. Safe numerical bounds for the Titchmarsh–Weyl $m(\lambda)$ -function	583
Rickert, J. H. Simultaneous rational approximations and related diophantine equations	461
Robinson, P. L. Quasifree representations of Clifford algebras	487
Roach, G. F. & Zhang, B. Spectral representation and scattering theory for the wave equation with two unbounded media	423
Rogers, L. C. G. & Hobson, D. G. Recurrence and transience of reflecting Brownian motion in the quadrant	387
Sánchez, F. & Castillo, J. M. F. Upper l_p -estimates in vector sequence spaces, with some applications	329
Sander, J. W. On primes not dividing binomial coefficients	225
Sarlet, W., Martinez, E. & Cariñena, J. Geometric characterization of separable second-order differential equations	205
Schwärzler, W. & Welsh, D. J. A. Knots, matroids and the Ising model	107
Slack, M. Infinite loop spaces with trivial Dyer–Lashof operations	311
Spanoudakis, N. K. Operators in finite distributive subspace lattices, I	141
Sugie, J. & Yoneyama, T. On the Liénard system which has no periodic solutions	413
Swinnerton-Dyer, Sir Peter. The Brauer group of cubic surfaces	449
Szücs, A. Double point surfaces of smooth immersions $M^n \rightarrow \mathbb{R}^{2n-2}$	601
Taniyama, K. Cobordism of theta curves in S^3	97
Teragaito, M. Roll-spun knots	91
Ullom, S. V. & Boston, N. Representations related to CM elliptic curves	71

Index

Urbanski, M. & Zamboni, L. On free actions of Λ -trees	535
Velani, S. L. Diophantine approximation and Hausdorff dimension in Fuchsian groups	343
Welsh, D. J. A. & Schwärzler, W. Knots, matroids and Ising model	107
Whittle, G. & Oxley, J. On sign-representable matroids	499
Wilson, R. A. Matrix for Fischer's group Fi_{24}	5
Wilson, S. M. J. & Holland, D. Localization and class groups of module categories with exactness defects	233
Yoneyama, T. & Sugie, J. On the Liénard system which has no periodic solutions	413
Zamboni, L. & Urbanski, M. On free actions of Λ -trees	535
Zhang, B. & Roach, G. F. Spectral representation and scattering theory for the wave equation with two unbounded media	423
Zimmermann, B. On a hyperbolic 3-manifold with some special properties.	87

THE PREPARATION OF MANUSCRIPTS

The attention of authors is particularly directed to the following requests.

1. Papers should be typed, double-spaced, on one side of white paper (of which A4, 210 by 297 mm, is a suitable size). The pages must be numbered. Margins of 30 mm should be left at the side, top and bottom of each page. Two clear copies should be sent.

A cover page should give the title, the author's name and institution, with the address at which mail is to be sent.

The title, while brief, must be informative (e.g. *A new proof of the prime-number theorem*, whereas *Some applications of a theorem of G. H. Hardy* would be useless).

The first paragraph or two should form a summary of the main theme of the paper, providing an abstract intelligible to mathematicians.

For a typescript to be accepted for publication, it must accord with the standard requirements of publishers, and be presented in a form in which the author's intentions regarding symbols etc. are clear to a printer (who is not a mathematician).

The following notes are intended to help the author in preparing the typescript. New authors may well enlist the help of senior colleagues, both as to the substance of their work and the details of setting it out correctly and attractively.

2. Notation

Notation should be chosen carefully so that mathematical operations are expressed with all possible neatness, to lighten the task of the compositor and to reduce the chance of error.

For instance n_k (n sub k) is common usage, but avoid if possible using c sub n sub k . Fractions are generally best expressed by a solidus. Complicated exponentials like

$$\exp\{z^2 \sin \theta / (1 + y^2)\}$$

should be shown in this and no other way.

In the manuscript, italics, small capitals and capitals are specified by single, double and triple underlinings. Bold faced type is shown by wavy underlining; wavy will be printed wavy.

It helps if displayed equations or statements which will be quoted later are numbered in order on the right of their line. They can then be referred to by, for example, 'from (7)'.
The author must enable the printer (if necessary by pencilled notes in the margin) to distinguish between similar symbols such as o , O , o , O ; x , X , \times ; ϕ , Φ , \emptyset ; l , 1 ; ε , \in ; κ , k .

Greek letters can be denoted by Gk in the margin.

If an author wishes to mark the end of the proof of a theorem, the sign \blacksquare may be used.

Footnotes should be avoided.

3. Diagrams

It is extremely helpful if diagrams are drawn in Indian ink on white card, faintly blue or green-lined graph paper, or tracing cloth or paper. *Symbols, legends and captions should be given on a transparent overlay.* Each text figure must be numbered as Figure 1, Figure 2, ... and its intended position clearly indicated in the manuscript:

Figure 1 here

The author's name in pencil must be on all separate sheets of diagrams.

A figure is expensive to reproduce and should be included only when the subject matter demands it, or when it greatly clarifies the exposition.

The Society recognizes that some authors do not have the facilities for producing drawings of a sufficiently high standard to be reproduced directly and it is therefore willing to have such diagrams re-drawn, provided that they are clear.

4. Tables

Tables should be numbered (above the table) and set out on separate sheets. Indicate the position of each in the text as for figures:

Table 3 here

5. References

References should be collected at the end of the paper numbered in alphabetical order of the authors' names. Titles of journals should be abbreviated as in *Mathematical Reviews*. The following examples show the preferred style for references to a paper in a journal, a paper in a proceedings volume, a book and an unpublished dissertation:

- [1] J. F. ADAMS. On the non-existence of elements of Hopf invariant one. *Ann. of Math.* (2) **72** (1960), 20–104.
- [2] M. P. FOURMAN and D. S. SCOTT. Sheaves and logic. In *Applications of Sheaves*, Lecture Notes in Math. vol. 753 (Springer-Verlag, 1979), pp. 302–401.
- [3] P. T. JOHNSTONE. *Stone Spaces*. Cambridge Studies in Advanced Math. no. 3 (Cambridge University Press, 1982).
- [4] F. W. LAWVERE. Functorial semantics of algebraic theories. Ph.D. thesis. Columbia University (1963).

*Mathematical Proceedings of
the Cambridge Philosophical Society*

MPCPCO 113 (Pt 3) 449-638 (1993) 0305-0041 May 1993


CONTENTS

	PAGE
SWINNERTON-DYER, SIR PETER. The Brauer group of cubic surfaces	449
RICKERT, J. H. Simultaneous rational approximations and related diophantine equations	461
JOHNSON, F. E. A. A rational invariant for certain infinite discrete groups	473
HOFMANN, K. H. & MORRIS, S. A. Generators on the arc component of compact connected groups	479
ROBINSON, P. L. Quasifree representations of Clifford algebras	487
OXLEY, J. & WHITTLE, G. On sign-representable matroids	499
BAKER, J. W. & LAU, A. T. Compact left ideal groups in semigroup compactification of locally compact groups.	507
AUINGER, K. The word problem for the bifree combinatorial strict regular semigroup	519
URBANSKI, M. & ZAMBONI, L. On free actions of Λ -trees	535
HINKKANEN, A. Julia sets of rational functions uniformly perfect	543
ARMITAGE, D. H., GARDINER, S. J. & NETUKA, I. Separation of points by classes of harmonic functions	561
KAMIYA, S. On H -balls and canonical regions of loxodromic elements in complex hyperbolic space	573
BROWN, B. M., KIRBY, V. G., EVANS, W. D. & PLUM, M. Safe numerical bounds for the Titchmarsh-Weyl $m(\lambda)$ -function	583
SZÜCS, A. Double point surfaces of smooth immersions $M^n \rightarrow \mathbb{R}^{2n-2}$	601
KESTEN, H. Convergence in distribution of lightly trimmed and untrimmed sums are equivalent.	615

© The Cambridge Philosophical Society 1993

Printed in Great Britain by the University Press, Cambridge

CAMBRIDGE
UNIVERSITY PRESS



0305-0041(199305)113:3;1-H