

P47.06

APD-induced plasticity and ERK1/2 translocation to the nucleus

P.J. Rogue¹*, S. Freyermuth²*, ¹U338 de INSERM, Centre de Neurochimie, Strasbourg; ²Faculté de Médecine, Université Louis Pasteur, Strasbourg, France

Objectives: Dopamine D₂ receptors are typical G_{i/o}-coupled receptors that represent key targets for antipsychotic drugs (APD). The therapeutic effects of APDs are delayed, suggesting long-term adaptations mediated by changes in gene expression possibly involving activation of the mitogen-activated protein kinase ERK1/2 cascade.

Methods: Using Western blot analysis, we investigated the regulation of ERK1/2 in both total and nuclear fractions from HEK293 or COS-7 cells transiently expressing D_{2L} or D_{2s}, following stimulation by quinpirole.

Summary of the results obtained: Western blot analysis with antibodies to the basal and phosphorylated forms of ERK1/2 in whole cell extracts from D_{2L}-COS7 cells, showed that QUIN stimulates a marked increase in the levels of p42 and p44 kDa phosphorylation whereas the total ERK1/2 is unaffected. QUIN-induced phosphorylation of ERK1/2 is prevented by prior treatment with APDS. The same overall pattern is observed with D_{2L}-HEK293 cells, or using D_{2s}. Pharmacological analysis of ERK1/2 phosphorylation following D_{2L} activation revealed that the signal transduction pathways involved seem to lead to activation of Ras and the Raf1/MEK/ERK cascade. After addition of QUIN to D_{2L}-COS7 cells, downstream transcription factor Elk-1 is phosphorylated following a similar time course to that of ERK1/2 phosphorylation. Regulation of the nuclear import of activated ERK1/2 by D₂ receptors was studied in digitonin permeabilised cells through Western blot analysis after subcellular fractionation. Upon activation by QUIN, a strong nuclear staining appeared. WGA pre-treatment inhibited QUIN-induced phospho-ERK1/2 staining at the level of the nucleus but not in the total extract. Similar results were obtained following incubation with a dominant negative importing. Furthermore, a marked increase in p-Elk-1 immunolabelling was observed in the nuclear fraction immediately following QUIN, and this effect was not completely abolished by WGA.

Conclusions: Overall, these results provide evidence that ERK1/2 activation and nuclear translocation dopamine D₂ receptors, suggesting these events play a role in APD-induced gene expression via Elk-1.

P48. Quality assurance**P48.01**

Patient satisfaction as a marker of psychiatric treatment quality – the TÜBB2000

G. Längle*, M. Schmidt-Hackenberg, H. Waschulewski. Tübingen University, Department of Psychiatry and Psychotherapy, Germany

In the industrialized countries, increasing account is being taken of the “consumer standpoint”, also in the context of psychiatric hospitalisation, as a significant element of quality management.

For the German-speaking region, an instrument applicable to both clinical routine and scientific investigations has been developed in recent years: the “Tübingen Treatment Satisfaction Form – TÜBB2000”.

Based on treatment areas of relevance to satisfaction from the patient's standpoint, this questionnaire was developed in a number of phases. 25 items were extracted by means of item analysis, and a significantly higher variance was achieved in the answers by using a

change-oriented question style. The validity of the instrument was verified inter alia by comparison with established instruments such as the CSQ 8. The accuracy of the registration detailing, however, proved to be greater with the TÜBB2000. Factor analysis was used to identify three factors reflecting key aspects of satisfaction: atmosphere (8 items), quality of treatment (8 items), and autonomy (7 items).

Within the framework of a cohort study of schizophrenics (N=169), regression analyses indicated that treatment satisfaction was of prognostic significance with respect to compliance in the post-hospitalisation course of treatment as well as to the re-hospitalisation of schizophrenic patients. Using the “TÜBB2000”, quality defects in routine clinical practice could be identified and appropriate measures planned for improvement and quality development.

P48.02

Evaluation of psychiatric day treatment

P. Garlipp, H. Haltenhof, B. Brüggemann, H. Vosschulte, K. Stritt, K.-P. Seidler. Hannover Medical School, Department of Social Psychiatry & Psychotherapy, Germany

In order to establish a continuous quality management in the psychiatric day clinic of the Hannover Medical School we developed a documentation system which combines the professionals' and the patients' perspectives. Besides ICD-10 and the GAF-Scale the therapists rated three items concerning awareness of illness, insight into illness and compliance. For self-report assessment the patients filled in the SCL-90-R and a weekly ward rating scale, which highlights among other things aspects of structure, atmosphere and interaction with the therapists. Prior to discharge the therapists and the patients rated changes of symptoms and general well-being and their satisfaction with the therapeutic outcome. Based on 84 treatment episodes over a one-year-period we present and discuss preliminary results focussing on four diagnostic groups (schizophrenia, depressive, personality and adjustment disorders).

P48.03

Data base of quality in psychiatric outpatient treatment. DIPSY

J. Aagaard*. Aarhus Psychiatric Hospital, Department of Psychiatric Demography, Denmark

Objective: During the last decade, the number of patients offered treatment in the CMH-centres or other outpatient clinics has rapidly increased. A large variation in target group, organisation and treatment modalities has been found. No closer knowledge of the quality of this treatment is available. A nation-wide database to document the quality of the treatment started in the spring 2001.

Methods: An interdisciplinary working group (DIPSY) selected quality indicators from clinical experience and evidence-based literature. Data is derived from the Psychiatric Case Register (DPC-Dipsy) and from supplementary clinical variables (WEB-Dipsy) obtained at yearly follow-up or at the end of an outpatient treatment period. Each treatment unit will be able to compare its results with average statistics from own county the entire country, and with the recommended standards of quality.

Results: The indicators and standards are as follows:

1. Symptoms/function: Gaf (80% og seve psychotic yield a Gaf>40).
2. Treatment: Bipolar affective psychosis, Maintenance therapy (90%). Schizophrenia, novel neuroleptic (50%). Psychotherapy (40% of patients with borderline personality disorders).
3. Social: Homelessness (5%).

4. Patients satisfaction with the information offered about disease and treatment (80%).
5. Continuity: Duration of outpatient treatment without admission to hospital (30% of severe psychotic >2 years).
6. Service: Waiting time (50% within 14 days after referral)

Conclusion: The database in progress may in time documents necessary aspects of the quality of the outpatient treatment in Denmark.

P48.04

Patterns of use of acute psychiatric units in Andalusia (Spain)

J.Miguel Pena-Andreu, J.L. Martínez González, M.I. Rodríguez Idígoras, G. Jiménez Lérica, M. Ruiz Ruiz, M.M. Jiménez Sartertero. *Unidad de Calidad, Delegacion Provincial de Salud, Malaga, Spain*

Objective: To study, for the first time, how patients are using the 17 General Hospital Psychiatric Hospitalization Units (GHPHUA) in Andalusia: The biggest area of Spain (population 7.236.459). (see the Poster: "Analysis of subgroups of psychiatric admissions in Andalusia (Spain)).

Methods: Using the CMBD (Andalusian Case Register) we have made an observational retrospective study of the 16873 users of the GHPHUA during 5 years.

Results:

1. The use of GHPHUA has been: 3060 patients (1995); 3013 (1996); 3557 (1997); 3590 (1998); and 3653 (1999).
2. The use by heavy users patients – with more than three admissions along the year- has been: 698 patients (1995); 683 (1996); 924 (1997), 915 (1998); and 1009 (1999).
3. There is an ascending evolution of this phenomenon in the analyzed temporary series: 22,81% (1995); 22,67% (1996); 25,98% (1997); 25,49% (1998) and 27,62% (1999).

Conclusions:

- i. The use of the GHPHUA is increasing progressively.
- ii. The fourth part of the patients who use them are heavy users.
- iii. The number of "revolving door" patients is being increasing every year.

P48.05

Analysis of subgroups of psychiatric admissions in Andalusia (Spain)

J.Miguel Pena-Andreu, J.L. Martínez González, M.I. Rodríguez Idígoras, M.J. Fernández Terrón, M.D. Martínez Cabrera. *Unidad de Calidad, Delegacion Provincial de Salud, Malaga, Spain*

Objective: To analyze admissions of single users and "revolving door" patients in the 17 General Hospital Psychiatric Hospitalization Units (GHPHUA) of Andalusia.

Methods: See the Poster: "Patterns of use of Acute Psychiatric Units in Andalusia (Spain)": Evaluated 45995 GHPHUA admissions during 5 years.

Results:

1. The number of admissions in GHPHUA has been: 7679 admissions (1995); 7640 (1996); 9853 (1997), 10062 (1998) and 10761 (1999).
2. Heavy users patients – with more than three admissions along the year- have had: 3557 admissions (1995); 3570 (1996); 5233 (1997), 5349(1998) and 6094 (1999).
3. There is an ascending evolution of admissions for "revolving door" patients: 46,32% (1995); 46,73% (1996); 53,11% (1997); 53,16% (1998) and 56,63% (1999); and descendent for not heavy users: 14,29%, 20,05% and 19,34% (1995); 14,19%,

19,69% and 19,4% (1996); 12,43%, 16,83% and 17,63% (1997); 12,09%, 17,49% and 17,26% (1998); 10,97%, 16,78% and 15,61% (1999), respectively for patients with 1, 2, or 3 admissions.

Conclusions:

- i. "Revolving door" patients consumes 50% of the admissions.
- ii. Such consumption is also progressively ascending.
- iii. Consumption is progressively descendent for not heavy users.

P48.06

A model to evaluate the reliability of the GAF-scale

P. Söderberg¹*, S. Tungstrom¹, B.-A. Armelius². ¹*Psychiatry R&D Department, Säter;* ²*Department of Psychology, Umeå University, Sweden*

The aim of this study is to develop a model of controlling the reliability of the GAF-scale, between sites in routine clinical practice.

Method: A database of 8 269 GAF-ratings from different patients in 34 clinics has been analysed. The material came from a nation-wide project, were every patient during two weeks was rated with the GAF-scale. To analyse the reliability between sites, a regression model was made, considering the database distribution of patients diagnoses, treatment frequency and patient data divided up in psychiatric clinics.

Result: The result shows overall that the GAF-scale can be reliable, in a sense of a global measurement in psychiatry when it used in clinical settings. Sites that have used the GAF- scale as a routine practice indicate good variability and their GAF-ratings fit the regression model according to the predicted values, when it was controlled for diagnoses, patients ages and treatment periods. This study also pointed out a reliability problem between sites, that "local cultures" have been developed in different clinics and highlights the need of quality assurance programme for routine assessments.

P48.07

Comparable unit descriptions for mental health and social services

M.-J. Haack¹*, D. Oberg², B. Sallmen³. ¹*University of Maastricht, The Netherlands* ²*Karolinska Institutet Stockholm, Stockholm;* ³*Lund University, Sweden*

The description of treatment settings and interventions is considered increasingly important in researching and planning mental health care. European Service Mapping Schedule (ESMS) was developed to describe services, but does not describe the interventions that are available within the services.

Monitoring Area and Phase System (MAPS) provides a method for describing treatment interventions, based on standardized item categories. The MAPS-Unit form describes units in the same dimensions as clients and is cross culturally applicable, theoretically based and valid in different fields of care. These dimensions are AREA and PHASE.

The method was originally developed within drug and alcohol treatment, but aimed to be applicable for other services because of frequent use of health and social services of the clients. The poster will introduce the categorization method and give examples of comparable unit descriptions within psychiatry, youth care, prison and drug treatment.