

Azodrin

does its job inside out-outside in

... and that means double trouble for cotton pests, because Shell's powerful new organo-phosphorus insecticide Azodrin works *two* ways.

By contact, immediately destroying insects on the plant surface. And systemically, by penetrating foliage rapidly where it kills sucking and chewing pests as they try to eat.

This sophisticated dual action makes Azodrin by far the most wide-ranging insecticide yet devised giving *continuing* control over *more* pests *more* effectively.


Azodrin has already been employed with outstanding success in the United States and

Central America, and is now proving world-wide to be one of the cotton grower's most powerful allies.


Suddenly pests that took a bite out of profits are losing their appetite.

For further information or supplies of Azodrin contact your Shell company or Shell chemicals distributor. Shell works today for your more profitable tomorrow.

Shell Chemicals



What's so special about the 240 Atomic Absorption Spectrophotometer?



HIGH PERFORMANCE: Signal integration. Scale expansion. Zero suppression. Stability

VERSATILITY: Provides both flame emission and absorption facilities.

FLEXIBILITY: Provides for a full range of lamp sources including electrodeless microwave lamps.

SIMPLICITY: Easy and speedy operation.

There are many other reasons that make the EEL 240 so special, our brochure and price list show most of the answers.



EVANS ELECTROSELENIUM LIMITED
HALSTEAD · ESSEX · ENGLAND

Telephone : Halstead 2461

Telex : 98204

For further advice Quote L.E.D. A

(ii)

ANIMAL PRODUCTION
Journal of the British Society of Animal Production

CONTENTS OF VOLUME 12 PART 2 MAY 1970 INCLUDE THE FOLLOWING

- DONALD and RUSSELL. The relationship between live weight of ewe at mating and weight of newborn lamb.
- DONALD, READ and RUSSELL. Influence of litter size and breed of sire on carcass weight and quality of lambs.
- STRANG. Litter productivity in Large White pigs. 1. The relative importance of some sources of variation.
- STRANG and KING. Litter productivity in Large White pigs. 2. Heritability and repeatability estimates.
- LAWRENCE. Effects on the bacon pig of adding protein and vitamin-mineral supplements to barley without mixing and at different time intervals.
- BROADBENT, MCINTOSH and SPENCE. The evaluation of a device for feeding group-housed animals individually.
- CURRAN, WIMBLE and HOLMES. Prediction of the voluntary intake of food by dairy cows. 1. Stall-fed cows in late pregnancy and early lactation.
- CURRAN and HOLMES. Prediction of the voluntary intake of food by dairy cows. 2. Lactating grazing cows.
- LYONS, CAFFREY and O'DONNELL. The effect of energy, protein and vitamin supplementation on the performance and voluntary intake of barley straw by cattle.
- KAY, MACDEARMID and MACLEOD. Intensive beef production. 10. Replacement of cereals with chopped straw.
- WOOD. The relationship between the month of calving and milk production.
- ROWLAND. The rumenitis and liver abscess complex and vitamin A status in beef cattle.
- OSMAN, EL-SAYED, AKKADA, ABOU and AGABAWI. Influence of rumen ciliate protozoa on conversion of food and growth rate in early-weaned Zebu calves.
- ANDREWS and ØRSKOV. A note on the effect of bulk density and digestibility on the voluntary intake of concentrate diets by sheep of two ages.
- DEIF, ABOU, AKKADA and EL-SHAZLY. A note on the utilization of urea nitrogen by sheep.
- TERNOUTH and BEATTIE. A note on the voluntary food consumption and the sodium-potassium ratio of sheep after shearing.
- Proceedings of the 51st meeting of the British Society of Animal Production, Harrogate, 23-25 March 1970.

OLIVER & BOYD LTD

Tweeddale Court, 14 High Street, Edinburgh EH1 1YL, Scotland

THE FOOD AND AGRICULTURE
ORGANIZATION OF THE
UNITED NATIONS

has an opening for a
**PESTICIDE FORMULATION
CHEMIST**

Applicants must have the appropriate University degree and at least 7 years' experience in pesticides formulation.

A good knowledge of English is essential and French or Spanish are desirable.

We offer liberal tax-free emoluments based on depth and breadth of training and experience.

Send brief résumé to the Chief, Recruitment Section, Personnel Division, FAO, Via delle Terme di Caracalla, 00100 Rome, Italy, quoting reference PL/GAPF/PI.

Application forms and detailed information will be sent to qualified applicants.

THE FOOD AND AGRICULTURE
ORGANIZATION OF THE
UNITED NATIONS

has openings in its various
programs for

General Agronomist
Irrigation Agronomist
Tropical Agronomist

Applicants must have the appropriate University degree and at least 7 years' experience in either of the above fields of specialization, preferably with working knowledge in dry or humid tropics.

Knowledge of English, French or Spanish is essential.

We offer liberal tax-free emoluments based on depth and breadth of training and experience.

Send brief résumé to the Chief, Recruitment Section, Personnel Division, FAO, Via delle Terme di Caracalla, 00100 Rome, Italy, specifying field of interest, quoting reference PL/GA/PI.

Application forms and detailed information will be sent to qualified applicants.

CONTENTS

Vol. 74 Part 2 April 1970

	PAGE
PIKE, I. H. The effect on nitrogen utilization in the sow of variation in dietary protein concentration and pattern of feeding in pregnancy	209
BROSTER, W. H., SUTTON, J. D., SMITH, T., BROSTER, VALERIE J. and BALCH, C. C. The effect of supplements of sucrose and of glucose monohydrate on the milk production and live weight of dairy cows. (With 4 text-figures)	217
REID, D. The effects of a wide range of nitrogen application rates on the yields from a perennial ryegrass sward with and without white clover. (With 8 text-figures)	227
THOMAS, N. W. and JUDGE, M. D. Alteration of porcine skeletal muscle myoglobin by the environment	241
ALLEN, T. E., BENNETT, J. W., DONEGAN, SUSAN M. and HUTCHINSON, J. C. D. Moisture, its accumulation and site of evaporation in the coats of sweating cattle. (With 3 text-figures)	247
MORRIS, J. G. and PAYNE, E. Ammonia and urea toxicoses in sheep and their relation to dietary nitrogen intake. (With 1 plate and 6 text-figures)	259
MEADLEY, J. T. and MILBOURN, G. M. The growth of vining peas. II. The effect of density of planting. (With 5 text-figures)	273
LEA, C. H., SWOBODA, P. A. T. and GATHERUM, D. P. A chemical study of soft fat in cross-bred pigs. (With 2 text-figures)	279
GORDON, J. G. and MCALLISTER, I. K. The circadian rhythm of rumination. (With 2 text-figures)	291
ROBERTSON, I. S., PAVER, H. and WILSON, J. C. Effect of castration and dietary protein level on growth and carcass composition in beef cattle. (With 3 text-figures)	299
WAINMAN, F. W., BLAXTER, K. L. and PULLAR, J. D. The nutritive value for ruminants of a complete processed diet based on barley straw	311
MORAN, J. B. Brahman cattle in a temperate environment. I. Live-weight gains and carcass characteristics. (With 2 plates and 2 text-figures)	315
MORAN, J. B. Brahman cattle in a temperate environment. II. Adaptability and grazing behaviour. (With 1 text-figure)	323
MONYO, J. H. and WHITTINGTON, W. J. Genetic analysis of root growth in wheat. (With 4 text-figures)	329
CONNIFFE, D., BROWNE, D. and WALSHE, M. J. Experimental design for grazing trials. (With 2 text-figures)	339
PERRY, D. A. The relation of seed vigour to field establishment of garden pea cultivars	343
OCARIZ, J. L. H., LITTLEJOHN, A. and ROBERTSON, I. S. A comparison of entire and ovariectomized beef heifers treated with ethylestrenol	349
SEARLE, T. W. Body composition in lambs and young sheep and its prediction <i>in vivo</i> from tritiated water space and body weight. (With 3 text-figures)	357
OJEHOMON, O. O. A comparison of the vegetative growth, development and seed yield of three varieties of cowpea, <i>Vigna unguiculata</i> (L.) Walp. (With 8 text-figures)	363
OJEHOMON, O. O. Effect of continuous removal of open flowers on the seed yield of two varieties of cowpea, <i>Vigna unguiculata</i> (L.) Walp. (With 3 text-figures)	375
BACHE, B. W. and ROGERS, N. E. Soil phosphate values in relation to phosphate supply to plants from some Nigerian soils. (With 2 text-figures)	383
DAVIES, J. M. L. and GARETH JONES, D. The response of seven barley varieties to inoculation in the field with different isolates of <i>Cercospora herpotrichoides</i> Fron. (With 1 text-figure)	391
BROCKMAN, J. S., SHAW, P. G. and WOLTON, K. M. The effect of phosphate and potash fertilizers on cut and grazed grassland. (With 6 text-figures)	397

SUBSCRIPTIONS. Two volumes of three parts are published annually. The subscription price is £6 net (U.S.A. \$19.50) per volume (post free); single parts are available at £2 12s. net (U.S.A. \$8.00) plus postage. Orders or enquiries may be sent to any bookseller or subscription agent, or to Cambridge University Press, P.O. Box 92, London, N.W. 1. (U.S.A. and Canada, Cambridge University Press American Branch, 32 East 57th Street, New York, N.Y. 10022, U.S.A.)