

Assessment of intake inadequacy and food sources of zinc of people in China

Guansheng Ma^{1,2,*}, Yanping Li^{1,2}, Ying Jin¹, Songming Du¹, Frans J Kok² and Xiaoguang Yang¹

¹National Institute for Nutrition and Food Safety, Chinese Center for Disease Control and Prevention, 29 Nan Wei Road, Beijing 100050, People's Republic of China: ²Division of Human Nutrition, Wageningen University, Wageningen, The Netherlands

Submitted 28 March 2006: Accepted 24 October 2006: First published online 19 February 2007

Abstract

Objectives: To assess the intake inadequacy and food sources of zinc of people in China.

Design and subjects: Diets of 68962 subjects aged 2–101 years (urban 21 103, rural 47 859) in the 2002 China National Nutrition and Health Survey were analysed. Dietary intake was assessed using 24-hour recall for three consecutive days. Zinc intake inadequacy was calculated based on values suggested by the World Health Organization.

Results: The median zinc intake ranged from 4.9 mg day⁻¹ (urban girls, 2–3 years) to 11.9 mg day⁻¹ (rural males, 19+ years). The zinc density of urban residents (2–3 to 19+ years) was 5.0–5.3 mg day⁻¹ (1000 kcal)⁻¹, significantly higher than that of their rural counterparts (4.7–4.8 mg day⁻¹ (1000 kcal)⁻¹). Differences in food sources of zinc from cereal grains (27.4–45.1 vs. 51.6–63.2%) and animal foods (28.4–54.8 vs. 16.8–30.6%) were found between urban and rural residents. Zinc from vegetables and fruits (8.2–13.8 vs. 9.7–12.4%) and legumes (1.3–3.3 vs. 2.5–3.4%) was comparable between urban and rural residents. The proportion of zinc intake inadequacy ranged between 2.8% (urban females, 19+ years) and 29.4% (rural lactating women). Rural residents had higher proportions of zinc intake inadequacy than their urban counterparts. Significantly higher proportions of zinc inadequacy were found in the category of phytate/zinc molar ratio > 15 for both rural and urban residents.

Conclusions: About 20% of rural children are at risk of inadequate zinc intake, with phytate as a potential important inhibitor. Moreover, lactating women are also considered a vulnerable group.

Keywords

Zinc
Phytate
Dietary intake
China National Nutrition and Health Survey

Zinc is an essential mineral that performs biochemical functions important for maintaining human health^{1–6}. Zinc deficiency may be widespread in developing countries but the true magnitude of mild and moderate zinc deficiency is unknown, in part because of the lack of a reliable and specific index of zinc status^{7,8}. Use of the Recommended Dietary Allowance⁹ and Estimated Average Requirement^{10,11} is suggested for estimating the prevalence of nutrient inadequacy in population groups.

Zinc deficiency may arise from low dietary intake, low bioavailability and/or interaction with other nutrients, and losses through disease processes^{12–14}. Inhibitors of zinc absorption are believed to be the more likely causative factor¹⁵. Phytate, present in whole grains, cereals and legumes, is a strong inhibitor of zinc absorption. Phytate is high in plant-based diets, which is a major inhibitory factor for zinc absorption. The phytate/zinc molar ratio is used to

predict the inhibitory effect of phytate on the bioavailability of zinc^{8,16,17}. The World Health Organization (WHO) suggests that the assessment of dietary zinc status should take inhibitory factors into account, and phytate data must be available⁸. However, the inhibitory effect of phytate on the bioavailability of zinc has not been examined in China owing to a lack of information on the phytate content of foods in the *China Food Composition Table*¹⁸. In 2005, the phytate content of 60 food samples commonly consumed in China was analysed¹⁹ using an anion-exchange method, which made it possible to assess the dietary zinc intake of people in China taking the inhibitory effect of phytate into account. Thus the aims of the present study were to: (1) assess the prevalence of zinc intake inadequacy in relation to the effect of phytate in Chinese diets; (2) examine if zinc intake inadequacy differs by age, sex and region in China; and (3) describe the food sources of zinc in Chinese diets.

*Corresponding author: Email mags@chinacdc.net.cn

Subjects and methods

Sampling

The 2002 China National Nutrition and Health Survey (CNNHS) is a nationally representative cross-sectional survey that covered 31 provinces, autonomous regions and the municipalities directly affiliated with the Central Government (Hong Kong, Macao and Taiwan were not included). A multi-stage cluster sampling method was used for subject selection. In stage 1, all 2860 counties/districts/cities of China were divided into six categories (big cities, medium and small cities, rural 1, 2, 3 and 4) based on type and level of economic development (from high to low). Twenty-two counties/districts/cities were randomly selected from each category, thus a total of 132 counties/districts/cities were randomly selected at this stage. In stage 2, three townships/sub-districts were randomly selected from each of the selected counties/districts/cities, resulting in 396 townships/sub-districts being randomly selected at this stage. In stage 3, two villages/neighbourhood committees were randomly selected from the selected townships/sub-districts, giving a total of 792 villages/neighbourhood committees. In stage 4, 90 households were randomly selected from each selected village/neighbourhood, which resulted in a final total of 71 971 households that were randomly selected to represent the national data.

Dietary intake assessment

The dietary survey was conducted among all members of 30 households that were randomly selected from the pre-selected 90 households. All family members above 2 years old from the selected households were invited for the dietary intake assessment. A total of 23 470 from 71 971 households participated in the dietary intake assessment²⁰.

The protocol of the survey was approved by the Ethical Committee of the National Institute for Nutrition and Food Safety, Chinese Center for Disease Control and Prevention. A signed consent form was obtained from each subject or his/her parent or guardian.

Information on food intake was collected using the 24-hour dietary recall method for three consecutive days (two weekdays and one weekend day) by trained interviewers. The parent or guardian was interviewed for children aged 2–16 years.

Zinc intake was calculated using the data of dietary recall in conjunction with the *China Food Composition Table*¹⁸. The composition of infant formula is available in the tables and was included in the calculation. Zinc content of foods in the *China Food Composition Table* was determined by atomic absorption spectrophotometry¹⁸. Duplicate food samples were analysed. Standard reference materials were obtained from the China National Center of Standard Material for quality control. The relative standard deviation was within 10%.

Sixty food samples, including rice, wheat flour, corn and soybean products commonly consumed in China, were selected based on the frequency of foods consumed in the 2002 CNNHS¹⁹. The phytate content of these food samples was determined using an anion-exchange method²¹. Phytate intake was calculated using the phytate content in conjunction with the dietary recall data. The average daily dietary intakes of zinc and phytate were calculated using the mean value of the three days' intakes.

The contributions of individual foods to zinc were calculated by summing the amount of zinc consumed from each food by all subjects in each age group and dividing by the total intake from all foods for all subjects in the respective age group.

Phytate/zinc molar ratio was calculated as the phytate intake per day in mmol divided by the zinc intake per day in mmol. The proportions of subjects with different phytate/zinc molar ratios (<5, 5–15, >15) were calculated. The percentage of people with zinc intake below the WHO normative requirement⁸ at different phytate/zinc molar ratios was calculated. The average individual normative requirements for zinc from the diet are developed according to zinc availability, the three (high, moderate and low) bioavailability levels corresponding to 50%, 30% and 15% absorption⁸. The normative requirements of the first trimester for pregnant women, and of first three months for lactating women, were used to assess the inadequacy of zinc intake for these groups.

Statistical analysis

Values of dietary zinc intake are expressed as median and interquartile range. Zinc intake is expressed as mg day⁻¹ and zinc density as mg day⁻¹(1000 kcal)⁻¹. Normal probability plots and Kolmogorov–Smirnov tests were used to determine whether variables followed a normal distribution. To analyse the associations of sex, age, region and their interactions with dietary zinc intake, a general linear model factorial analysis was applied with Tukey's *post hoc* comparisons. The results are presented by sex, age and region for the significant interactions found between sex and age, and between age and region. Differences in zinc intake of subjects by sex and region were compared using Wilcoxon's signed rank sum test. The proportion of subjects with different phytate/zinc molar ratios was calculated, while the proportion of zinc intakes less than the WHO suggested normative requirement⁸ was calculated by region and phytate/zinc molar ratio. Differences in the above-mentioned proportions between urban and rural areas were compared using the χ^2 test. All statistical analyses were conducted with the SAS Statistical Package (8.2e for Windows; SAS Institute Inc.). Statistical significance was set at 0.05.

Results

Characteristics of the study population

A total of 68 962 subjects were included in the present analysis. Of these, 21 103 (males 10 027, females 11 076) were from urban areas, while 47 859 (males 23 524, females 24 335) were from rural areas. There were 310 pregnant women and 470 lactating women. The means of body mass index and dietary energy intake are presented by age group and region in Table 1.

Dietary zinc intake of people in China

The median zinc intakes by age, sex and region are presented in Table 2. There were significant effects of age ($F(6, 68\,956) = 372.8; P < 0.001$) and sex ($F(1, 68\,961) = 113.8; P < 0.001$), and first-level interactions between region and age ($F(6, 68\,956) = 22.5; P < 0.001$) and between age and sex ($F(6, 68\,956) = 15.5; P < 0.001$), for dietary zinc intakes. Median zinc intake ranged from 4.9 mg day^{-1} (urban girls, 2–3 years) to 11.9 mg day^{-1} (rural males, 19+ years). The zinc intake of urban boys aged 2–3 years was higher than that of their rural counterparts (6.2 vs. 5.1 mg day^{-1}). In contrast, the zinc intake of rural adults (19+ years) was significantly higher than that of their urban counterparts (males 11.9 vs. 10.5 mg day^{-1} ; females 10.2 vs. 8.9 mg day^{-1}). The younger the age group, the less the zinc intake. Rural lactating women consumed more zinc than their urban counterparts (11.0 vs. 9.8 mg day^{-1}) whereas urban pregnant women consumed comparable amounts of zinc to their rural counterparts (10.7 vs. 10.9 mg day^{-1}).

There were significant effects of age ($F(6, 68\,956) = 4.9; P < 0.001$) and region ($F(1, 68\,961) = 278.8; P < 0.001$), first-level interactions between region and age ($F(6, 68\,956) = 4.2; P < 0.001$) and between age and sex ($F(6, 68\,956) = 3.3; P < 0.001$), as well as second-level

interactions between sex, region and age ($F(4, 68\,958) = 3.9; P < 0.001$) for dietary zinc density. The median zinc density of all age groups was between 4.7 and $5.7\text{ mg day}^{-1}(1000\text{ kcal})^{-1}$ (Table 2). The zinc density of urban residents (2–3 to 19+ years) was 5.0 – $5.3\text{ mg day}^{-1}(1000\text{ kcal})^{-1}$, significantly higher than that of their rural counterparts (4.7 – $4.8\text{ mg day}^{-1}(1000\text{ kcal})^{-1}$). No significant difference was found between different age groups. Variation in median zinc intake was found between males and females within the same region and age group. Zinc density of the diets of urban boys (2–3 years) was higher than that of their female counterparts; in contrast, the rural boys (2–3 years) and rural adults consumed diets of lower zinc density than their female counterparts. Urban pregnant women had higher dietary zinc density than their rural counterparts (5.7 vs. $4.7\text{ mg day}^{-1}(1000\text{ kcal})^{-1}$). No difference in zinc density was found between urban and rural lactating women.

Food sources of zinc

Cereal grains including rice, wheat flour and its products contributed 27.4–63.2% to dietary zinc intake, while 16.8–54.8% zinc came from animal foods. Vegetables and fruits provided 8.2–13.8% of dietary zinc, while legumes contributed only 1.3–3.4% of dietary zinc. Large discrepancies were found in the food sources of zinc from staple and animal foods between urban and rural residents within the same age group. Staple foods provided 27.4–45.1% zinc for urban residents, 51.6–63.2% for rural residents. Animal foods contributed 28.4–54.8% of dietary zinc for urban residents, much higher than that for their respective rural counterparts (16.8–30.6%). Vegetables, fruits and legumes provided comparable zinc for both urban and rural residents. Detailed information on the food sources of dietary zinc intake stratified by age, sex and region is shown in Table 3.

Table 1 Characteristics of the study population

Group	Urban					Rural				
	n	BMI (kg m^{-2})		Energy (kcal day^{-1})		n	BMI (kg m^{-2})		Energy (kcal day^{-1})	
		Mean	SD	Mean	SD		Mean	SD	Mean	SD
Males										
2–3 years	154	16.2	1.8	1386.8	1006.7	623	16.2	1.7	1212.8	613.4
4–8 years	455	15.9	2.3	1528.3	694.6	2049	15.2	1.5	1589.7	585.9
9–13 years	641	17.4	2.9	1858.0	711.9	2641	16.5	2.4	2007.1	652.4
14–18 years	554	19.7	3.1	2196.0	808.5	1461	18.8	2.5	2414.2	700.2
19+ years	8223	24.0	3.5	2129.9	790.0	16 750	22.4	3.2	2600.2	750.1
Females										
2–3 years	108	15.7	1.4	1439.0	2510.9	466	15.6	1.5	1160.7	537.7
4–8 years	436	15.3	2.0	1392.8	557.0	1758	14.8	1.4	1512.6	618.4
9–13 years	598	17.0	2.9	1700.1	645.2	2351	16.4	2.4	1830.0	571.6
14–18 years	484	20.3	3.0	1816.0	656.3	1259	19.8	2.5	2076.9	694.1
19+ years	9340	23.9	3.8	1808.0	749.2	17 831	22.9	3.5	2238.6	678.1
Pregnant	55	22.5	3.2	1938.7	603.3	255	24.2	3.5	2380.7	714.4
Lactating	55	22.3	2.0	1979.1	635.6	415	23.2	4.4	2421.6	742.7

BMI – body mass index; SD – standard deviation.

Table 2 Median (P50) and interquartile range (P25, P75) of zinc intake and zinc density by age, sex and region

Group	Zinc intake (mg day ⁻¹)						Zinc density (mg day ⁻¹ (1000 kcal) ⁻¹)					
	Urban			Rural			Urban			Rural		
	P25	P50	P75	P25	P50	P75	P25	P50	P75	P25	P50	P75
Males												
2–3 years	4.7	6.2 ^c	7.9	3.9	5.1 ^{*,e}	6.9	4.5	5.3	6.4	4.1	4.7 [*]	5.4
4–8 years	5.2	7.1 ^c	9.3	5.5	7.1 ^d	9.0	4.3	5.0	6.0	4.2	4.7 [*]	5.3
9–13 years	6.9	9.1 ^b	12.1	7.3	9.0 ^c	11.2	4.4	5.0	6.3	4.3	4.7 [*]	5.2
14–18 years	8.4	10.7 ^a	13.8	8.9	11.0 ^b	13.6	4.3	5.1	6.0	4.3	4.7 [*]	5.2
19+ years	8.1	10.5 ^a	13.6	9.7	11.9 ^{*,a}	14.4	4.4	5.1	6.1	4.3	4.7 [*]	5.2
Females												
2–3 years	3.7	4.9 ^{†cd}	7.2	3.7	5.1 ^a	7.0	4.2	5.1 [†]	6.1	4.2	4.8 [†]	5.5
4–8 years	5.1	6.7 ^c	8.9	5.2	6.8 ^{†b}	8.6	4.4	5.1	6.1	4.2	4.7	5.3
9–13 years	6.3	8.2 ^{†ad}	10.9	6.7	8.3 ^{†c}	10.3	4.4	5.1	6.0	4.3	4.8 [*]	5.3
14–18 years	7.0	8.9 ^{†ab}	11.8	7.6	9.3 ^{†d}	11.8	4.4	5.1	6.3	4.3	4.8 [*]	5.2
19+ years	6.9	8.9 ^{†b}	11.6	8.3	10.2 ^{†+e}	12.5	4.4	5.2	6.2	4.3	4.8 [†]	5.3
Pregnant	8.1	10.7 ^{ab}	14.1	8.9	10.9 ^{ef}	13.6	4.5	5.7	6.6	4.2	4.7 [*]	5.3
Lactating	7.1	9.8 ^{abcd}	11.9	9.0	11.0 ^{*,fg}	13.3	4.2	4.9	5.9	4.3	4.8	5.3

A general linear model was performed with Tukey *post hoc* analysis to compare the effects of age (by sex and by region). Values not sharing the same superscript letter (a–g) denote a significant difference between age groups of the same sex, $P < 0.05$.

* Significant difference between urban and rural residents within the same age and sex group, Wilcoxon's signed rank sum test: $P < 0.05$.

† Sex difference within the same age and region, Wilcoxon's signed rank sum test: $P < 0.05$.

Proportions of subjects with different phytate/zinc molar ratios

Variations in the proportion of subjects with the same phytate/zinc molar ratio were found between the different age and/or sex subgroups (Table 4). Some 28.6–46.8% of urban residents and 3.9–8.0% of rural residents had phytate/zinc molar ratio < 5 , while 48.7–65.7% of residents had phytate/zinc molar ratio between 5 and 15, and 4.5–18.2% of urban residents and 27.8–36.4% of rural residents had phytate/zinc molar ratio > 15 . The proportion of rural residents with phytate/zinc molar ratio > 15 was significantly higher than that of their urban counterparts. A significant effect of age on the proportion of subjects with different phytate/zinc molar ratios was found in all subgroups. Significant sex difference was only found in rural children aged 9–13 years (Table 4).

Prevalence of zinc intake inadequacy

Table 5 presents the prevalence of zinc intake inadequacy by age, sex and region. The proportion of zinc intake inadequacy ranged between 2.8% (urban females, 19+ years) and 29.4% (rural lactating women). The distribution was much higher when phytate/zinc molar ratio > 15 was considered: 10.3% for rural females, 19+ years and 100% for urban lactating women. The proportion of zinc intake inadequacy among urban residents was 2.8–16.4%, significantly lower than that of their rural counterparts (3.6–29.4%). Prevalence of zinc intake inadequacy was significantly higher among male adults (19+ years) than their female counterparts. The zinc intake of 5.5% of pregnant women and 29.4% of lactating women was inadequate.

Discussion

Information on zinc intake in developing countries is limited because of the paucity of data on the zinc content of local staple foods²². China developed food composition tables in the 1960s and keeps them updated using reliable and valid determination methods¹⁸. The composition of fortified foods is also available in the *China Food Composition Table*. The strength of the present study is the use of three days of 24-hour dietary recall data of a national representative sample population in conjunction with the *China Food Composition Table*. One limitation of this study is that the zinc from dietary supplements was not taken into account. As the use of dietary supplements in China²³ is low (urban 10.3%, rural 2.9%) compared with that (39.5%) in the USA⁹, we assumed that zinc intake was underestimated to a small extent. This limitation suggests that the use of dietary supplements and their contribution to total nutrient intake in the Chinese population, especially for urban persons, should be monitored in future surveys.

The present study found that the zinc intake of people in China is lower than that of the US population^{9,24,25}, but comparable to that in the UK^{26,27}, Germany²⁸ and New Zealand²⁹. People in China consume more zinc than their counterparts in Chile³⁰, India³¹ and Korea³². However, the adequacy of zinc intake depends not only on the amount, but also its bioavailability. The inhibitory factors that may interact with zinc should be examined when assessing the dietary intake of zinc^{8,33}. People consuming a diet that provides marginal zinc intake may not absorb an adequate amount of zinc if they are also consuming foods high in phytate. The average phytate intake of people in China (1186 mg day⁻¹)³⁴ is higher than that in Western

Table 3 Food sources of dietary zinc by age, sex and region (%)

Group	Urban						Rural					
	Vegetables & fruits			Legumes			Nuts, cake, sauce, oil			Other		
	Cereal grains	Animal foods	Vegetables & fruits	Legumes	Nuts, cake, sauce, oil	Other	Cereal grains	Animal foods	Vegetables & fruits	Legumes	Nuts, cake, sauce, oil	Other
Males												
2-3 years	28.9	51.8	8.4	2.4	2.4	6.0	51.6	30.6	9.7	3.2	1.6	3.2
4-8 years	40.0	38.8	10.0	1.3	3.8	6.3	60.0	20.0	12.0	2.7	2.7	2.7
9-13 years	41.6	34.7	10.9	3.0	4.0	5.9	61.2	18.4	11.2	3.1	3.1	3.1
14-18 years	42.2	35.3	10.3	2.6	4.3	5.2	63.2	17.9	11.1	2.6	2.6	2.6
19+ years	42.2	34.5	11.2	2.6	4.3	5.2	61.9	18.3	11.1	3.2	3.2	2.4
Females												
2-3 years	27.4	54.8	8.2	1.4	2.7	5.5	53.4	27.6	10.3	3.4	1.7	3.4
4-8 years	41.3	34.7	10.7	2.7	4.0	6.7	58.9	20.5	12.3	2.7	2.7	2.7
9-13 years	41.8	34.1	12.1	3.3	3.3	5.5	61.8	16.9	12.4	3.4	3.4	2.2
14-18 years	37.6	38.6	10.9	2.0	4.0	6.9	62.4	16.8	11.9	3.0	3.0	3.0
19+ years	40.0	34.0	13.0	3.0	4.0	6.0	62.0	17.6	12.0	2.8	2.8	2.8
Pregnant	43.1	28.4	13.8	2.6	4.3	7.8	58.6	18.1	12.1	3.4	3.4	4.3
Lactating	45.1	30.4	11.8	2.9	4.9	4.9	60.2	20.3	11.9	2.5	2.5	2.5

Table 4 Proportion of subjects with different phytate/zinc molar ratio (%)

Group	Urban			Rural		
	Phytate/zinc molar ratio			Phytate/zinc molar ratio*		
	<5	5-15	>15	<5	5-15	>15
Males						
2-3 years	46.8	48.7	4.5	8.0	64.2	27.8*
4-8 years	28.6	62.9	8.6	5.0	65.2	29.9*
9-13 years	32.6	58.8	8.6	5.6	61.8	32.5*
14-18 years	32.1	59.7	8.1	4.8	61.7	33.5*
19+ years	32.4	60.6	7.0	5.2	61.7	33.1*
Age effect	$\chi^2 = 21.6, P < 0.01$			$\chi^2 = 25.3, P < 0.01$		
Females						
2-3 years	35.2	59.3	5.6	7.7	64.4	27.9*
4-8 years	30.3	59.6	10.1	5.7	65.7	28.6*
9-13 years	33.1	57.5	9.4	3.9	63.1	33.0*†
14-18 years	34.1	59.9	6.0	5.8	59.5	34.7*
19+ years	32.7	60.2	7.1	4.9	61.1	34.0*
Pregnant	30.9	50.9	18.2	6.3	61.6	32.2*
Lactating	29.1	63.6	7.3	5.1	58.6	36.4*
Age effect	$\chi^2 = 22.0, P < 0.05$			$\chi^2 = 45.4, P < 0.01$		

* Significant difference between urban and rural residents, χ^2 test: $P < 0.05$.

† Sex difference within the same age and region, χ^2 test: $P < 0.05$.

countries^{35,36} and less than that of Korean³⁷ and Indian people³⁸. The proportion of people with inadequate zinc intake was considerably high in the phytate/zinc molar ratio > 15 category, indicating that zinc in the diets of these populations would be affected by their high phytate intake. The vulnerable groups of zinc deficiency in China are children, adolescents and lactating women, especially those in rural areas. As zinc plays an important role in children's growth³⁹ and during pregnancy^{40,41}, zinc deficiency will impact severely on human health⁴², with adverse effects that include growth retardation, abnormal

Table 5 Proportion of zinc intakes less than the normative requirement of the World Health Organization (%)

Group	Urban				Rural			
	Phytate/zinc molar ratio				Phytate/zinc molar ratio			
	<5	5-15	>15	Total	<5	5-15	>15	Total
Males								
2-3 years	0.0	8.8	33.3	6.3	4.5	9.1	69.5	25.8*
4-8 years	0.9	9.0	69.7	11.9	0.0	1.1*	46.1*	14.7
9-13 years	0.0	7.3	55.3	8.9	0.0	0.9*	49.3	16.8*
14-18 years	2.8	6.3	67.6	10.2	1.9	1.6*	47.4*	17.2*
19+ years	0.2	2.8	43.6	4.8	0.4	0.3*	19.4*	6.6*
Females								
2-3 years	2.9	5.9	80.0	8.8	3.2	6.7	62.5	21.7*
4-8 years	1.0	3.4†	55.0	8.6	0.0	2.6†	46.5	15.1*
9-13 years	0.0	6.0	51.0	8.3	0.0	1.9*†	42.5†	15.6*
14-18 years	0.7	5.1	50.0	5.9†	1.6	2.5*	49.7	18.8*
19+ years	0.1	1.3†	27.8†	2.8†	0.1	0.2*†	10.3*†	3.6*†
Pregnant	0.0	0.0	40.0	7.3	0.0	0.0	17.1	5.5
Lactating	0.0	14.3	100.0	16.4	0.0	2.9*	76.2	29.4*

* Significant difference between urban and rural residents, χ^2 test: $P < 0.05$.

† Sex difference within the same age and region, χ^2 test: $P < 0.05$.

immune function, impaired cognitive function, increased maternal morbidity, prolonged gestation and increased risk to the foetus.

Information on the inadequacy of dietary zinc intakes should be interpreted together with data derived from other assessment methods. The national prevalence of stunting among children <5 years old can be used as an indirect indicator of a population's risk of zinc deficiency^{43,44}. The national prevalence of stunting in China is 3.3–8.6% for urban children <5 years and 17.3–20.9% for their rural counterparts⁴⁵. It can be concluded that the level of risk for zinc deficiency is low for the urban population and moderate for the rural population in China. Further population assessment of zinc status by serum zinc concentrations in a representative sample of the population should be made in order to identify segments of the population at high risk. Appropriate intervention strategies including supplementation, food fortification, dietary diversification and biofortification should be developed to target different populations.

In conclusion, urban people in China are at low risk of zinc deficiency, whereas the rural population is at a moderate level of risk for zinc deficiency. Lactating women and rural children are vulnerable populations in China. Effective strategies correcting these problems should be developed and implemented in targeted risk groups.

Acknowledgements

Sources of funding: The 2002 China National Nutrition and Health Survey was funded by the Ministry of Health and Ministry of Science and Technology, China (2001DEA30035, 2003DIA6N008).

Conflict of interest declaration: There is no interest conflict.

Authorship responsibilities: G.M. was responsible for study design, project management, data analysis, manuscript writing and interpretation of results. Y.L. was responsible for data cleaning and analysis, manuscript writing and interpretation of results. Y.J. was responsible for determining the phytate content of foods, manuscript writing and interpretation of results. S.D. was responsible for manuscript writing and interpretation of results. F.J.K. was responsible for manuscript writing and interpretation of results. X.Y. was responsible for study design, project management and interpretation of results.

Acknowledgements: We thank all team members and all participants from the 31 provinces. We are very appreciative of the support of UNICEF, WHO, Unilever China and the Danone Nutrition Institute China.

References

- 1 Abdel-Mageed AB, Oehme FW. A review of the biochemical roles, toxicity and interactions of zinc, copper and iron: I. zinc. *Veterinary and Human Toxicology* 1990; **32**: 34–9.
- 2 Black RE, ed. Zinc for Child Health: Proceedings of a Symposium [special issue]. *American Journal of Clinical Nutrition* 1998; **68**: 409S–516S.
- 3 Brown KH, Wuehler SE, eds. *Zinc and Human Health: Results of Recent Trials and Implications for Program Interventions and Research*. Ottawa: The Micronutrient Initiative, 2000.
- 4 Hambidge M, Cousins RJ, Costello RB, eds. Zinc and Health: Current Status and Future Directions [special issue]. *Journal of Nutrition* 2000; **130**: 1341S–519S.
- 5 Institute of Medicine, Food and Nutrition Board. *Dietary Reference Intakes for Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine, Iron, Manganese, Molybdenum, Nickel, Silicon, Vanadium, and Zinc*. Washington, DC: National Academy Press, 2001.
- 6 Mertz W, Morris ER, Smith JC Jr, Udomkesmalee E, Fields M, Levander OA, *et al.* Trace elements in the elderly: metabolism, requirements, and recommendations for intakes. In: Munro HN, Danford DE, eds. *Nutrition, Aging, and the Elderly*. New York: Plenum Press, 1989; 205–17.
- 7 Hambidge KM. Assessing the trace element status of man. *Proceedings of the Nutrition Society* 1988; **47**: 37–44.
- 8 World Health Organization (WHO). *Trace Elements in Human Nutrition and Health*. Geneva: WHO, 1996.
- 9 Briefel RR, Bialostosky K, Kennedy-Stephenson J, McDowell AM, Ervin RB, Wright JD. Zinc intake of the US population: findings from the Third National Health and Nutrition Examination Survey, 1988–1994. *Journal of Nutrition* 2000; **130**: 1367S–73S.
- 10 Institute of Medicine, Food and Nutrition Board. *Dietary Reference Intakes: Applications in Dietary Assessment*. Washington, DC: National Academy Press, 2000.
- 11 Barr SI, Murphy SP, Poos MI. Interpreting and using the dietary reference intakes in dietary assessment of individuals and groups. *Journal of the American Dietetic Association* 2002; **102**: 780–8.
- 12 Cousins RJ. Zinc. In: Ziegler EE, Filer LJ, eds. *Present Knowledge in Nutrition*. Washington, DC: ILSI Press, 1996; 293–306.
- 13 Wada L, King JC. Trace element nutrition during pregnancy. *Clinical Obstetrics and Gynecology* 1994; **37**: 574–86.
- 14 Walsh CT, Sandstead HH, Prasad AS, Newberne PM, Fraker PJ. Zinc: health effects and research priorities for the 1990's. *Environmental Health Perspectives* 1994; **102**(Suppl. 2): 5–46.
- 15 Lönnerdal B. Dietary factors influencing zinc absorption. *Journal of Nutrition* 2000; **130**: 1378S–83S.
- 16 Turnlund JR, King JC, Keyes WR, Gong B, Michel MC. A stable isotope study of zinc absorption in young men: effects on phytate and α -cellulose. *American Journal of Clinical Nutrition* 1984; **40**: 1071–7.
- 17 Rossander L, Sandberg AS, Sandström B. The influence of dietary fibre on mineral absorption and utilization. In: Schweizer TF, Edwards CA, eds. *Dietary Fibre – A Component of Food*. London: Springer-Verlag, 1992; 197–216.
- 18 Yang Y, Wang G, Pan X. *China Food Composition Table 2002*. Beijing: Beijing Medical University Publishing House, 2002; 343pp.
- 19 Ma G, Jin Y, Piao J, Kok F, Bonnema G, Jacobsen E. Phytate, calcium, iron, and zinc contents and their molar ratios in foods commonly consumed in China. *Journal of Agricultural and Food Chemistry* 2005; **53**: 10285–90.
- 20 Wang L. *Report of China National Nutrition and Health Survey 2002 (1): Summary Report*. Beijing: People's Medical Publishing House, 2005; 18–45.
- 21 Harland BF, Oberleas D. Anion-exchange method for determination of phytate in foods: collaborative study. *Journal of the Association of Official Analytical Chemists* 1986; **69**: 667–70.

- 22 Gibson RS. Zinc: the missing link in combating micronutrient malnutrition in developing countries. *Proceedings of the Nutrition Society* 2006; **65**: 51–60.
- 23 Ma G, Cui Z, Li Y, Hu X, Wang J, Yang X. The survey about the use of dietary supplements by Chinese Adults. *Acta Nutrimenta Sinica* 2006; **28**: 8–10.
- 24 Ervin RB, Kennedy-Stephenson JK. Mineral intakes of elderly adult supplement and non-supplement users in the Third National Health and Nutrition Examination Survey. *Journal of Nutrition* 2002; **132**: 3422–7.
- 25 Arsenault EJ, Brown HK. Zinc intake of US preschool children exceeds new dietary reference intakes. *American Journal of Clinical Nutrition* 2003; **78**: 1011–7.
- 26 Bailey AL, Maisey S, Southon S, Wright AJA, Finglas PM, Fulcher RA. Relationships between micronutrient intake and biochemical indicators of nutrient adequacy in a 'free-living' elderly UK population. *British Journal of Nutrition* 1997; **77**: 225–42.
- 27 Thane CW, Bates CJ, Prentice A. Zinc and vitamin A intake and status in a national sample of British young people aged 4–18 y. *European Journal of Clinical Nutrition* 2004; **58**: 363–75.
- 28 Schulze MB, Linseisen J, Krobe A, Boening H. Macro-nutrient, vitamin, and mineral intakes in the EPIC–Germany cohorts. *Annals of Nutrition & Metabolism* 2001; **45**: 181–9.
- 29 Jong de N, Gibson SR, Thomson DC, Ferguson LE, McKenzie EJ, Green JT, *et al.* Selenium and zinc status are suboptimal in a sample of older New Zealand women in a community-based study. *Journal of Nutrition* 2001; **131**: 2677–84.
- 30 Olivares M, Pizarro F, de Pablo S, Araya M, Uauy R. Iron, zinc, and copper: contents in common Chilean foods and daily intakes in Santiago, Chile. *Nutrition* 2004; **20**: 205–12.
- 31 Ram BS, Mohammad AN, Shanti SR, Sarita B, Zhang G, Zhu S. Current zinc intakes and risk of diabetes and coronary artery disease and factors associated with insulin resistance in rural and urban populations of North India. *Journal of the American College of Nutrition* 1998; **17**: 564–70.
- 32 Joung H, Nam G, Yoon S, Lee J, Shim JE, Paik HY. Bioavailable zinc intake of Korean adults in relation to the phytate content of Korean foods. *Journal of Food Composition and Analysis* 2004; **17**: 713–24.
- 33 Oberleas D, Harland BF. Phytate contents of food: effect on dietary zinc bioavailability. *Journal of the American Dietetic Association* 1981; **79**: 433–6.
- 34 Ma G, Li Y, Jin Y, Zhai F, Kok FJ, Yang X. Phytate intake and molar ratios of phytate to zinc, iron and calcium in the diets of people in China. *European Journal of Clinical Nutrition* 2006 Aug 23 [Epub ahead of print].
- 35 Plaami S, Kumpulainen J. Inositol phosphate content of some cereal-based foods. *Journal of Food Composition and Analysis* 1995; **8**: 324–35.
- 36 Carnovale E, Lombardi-Boccia G, Lugaro E. Phytate and zinc content of Italian diets. *Human Nutrition: Applied Nutrition* 1987; **41A**: 180–6.
- 37 Kwun IS, Kwon CS. Dietary molar ratios of phytate:zinc and millimolar ratios of phytate × calcium: zinc in South Koreans. *Biological Trace Element Research* 2000; **75**: 29–41.
- 38 Khokhar S, Pushpanjali, Fenwick GR. Phytate content of Indian foods and intakes by vegetarian Indians of Hisar Region, Haryana State. *Journal of Agricultural and Food Chemistry* 1994; **42**: 2440–4.
- 39 Brown KH, Peerson JM, Rivera J, Allen LH. Effect of supplemental zinc on the growth and serum zinc concentrations of prepubertal children: a meta-analysis of randomized controlled trials. *American Journal of Clinical Nutrition* 2002; **75**: 1062–71.
- 40 Caulfield LE, Zavaleta N, Shankar AH, Merialdi M. Potential contribution of maternal zinc supplementation during pregnancy to maternal and child survival. *American Journal of Clinical Nutrition* 1998; **68**: 499S–508S.
- 41 Shah D, Sachdev HPS. Zinc deficiency in pregnancy and fetal outcome. *Nutrition Reviews* 2006; **64**: 15–30.
- 42 Prasad AS. *Biochemistry of Zinc*. New York: Plenum, 1993; 193–218.
- 43 Hotz C, Lowe NM, Araya M, Brown KH. Assessment of the trace element status of individuals and populations: the example of zinc and copper. *Journal of Nutrition* 2003; **133**: 1563S–8S.
- 44 Hotz C, Brown KM. Assessment of the risk of zinc deficiency in populations and options for its control. *Food and Nutrition Bulletin* 2004; **25**: S99–199.
- 45 Yang X, Wang Z, He Y, Yu W, Hu Y, Zhai F. Trends and prevalence of malnutrition among Chinese children under five years. *Acta Nutrimenta Sinica* 2005; **27**: 185–8.