JOURNAL of the

AUSTRALIAN MATHEMATICAL SOCIETY

Series	A —	- Pure Mat	hema	tics	and	Statistics
		Volume 4	6 P	art	1	
		FEBRUA	ARY	1989)	

CARLO PUCCI	A maximum principle related to level surfaces of solutions of parabolic equations			
DON BRUNKER and DENIS HIGGS	Constructions of Σ -groups, relatively free Σ -groups	8		
N. S. HEKSTER	Varieties of groups and isologisms	22		
Józef Banaś	Integrable solutions of Hammerstein and Urysohn integral equations			
STEVEN N. EVANS	The maximum distribution of a Gaussian stochas- tic process indexed by a local field			
D. C. KENT and G. D. RICHARDSON	Ordered probabilistic metric spaces	88		
RICHARD GLENDINNING	The growth of the expected number of real zeros of a random polynomial	100		
ERICH W. ELLERS and HUBERTA LAUSCH	Length theorems for the general linear group of a module over a local ring	122		
J. L. MENNICKE and B. H. NEUMANN	More on finite groups with few defining relations	132		
J. C. FERRANDO and M. LÓPEZ-PELLICER	Quasi-suprabarrelled spaces	137		
ANTHONY G. PAKES	An infinite alleles version of the Markov branching process	146		

Set by the American Mathematical Society

https://doi.org/10.1017/S1446788700039329 Publishee online by Cambridge University Press, Michigan, U.S.A.

THE JOURNAL

The Journal of the Australian Mathematical Society began publication in 1959, and from 1975 has appeared in two series, Series A (Pure Mathematics and Statistics) and Series B (Applied Mathematics). Information about Series B may be found on the inside back cover.

Series A, Pure Mathematics and Statistics

Submission of research papers in pure mathematics and mathematical statistics is invited. Authors intending to submit papers for publication should prepare the typescript following the instructions below, and should send two complete copies to the Editor, or to an appropriate Associate Editor (see back cover of this issue).

Two volumes of three parts each are planned for 1989. These are Volume 46 and Volume 47. The annual subscription price to non-members is A\$181.00 (US\$146.00) for 1989. Each part in 1989 will have 20% more pages than in 1988.

PREPARATION OF TYPESCRIPTS

1. The author should keep a complete copy of the submitted article; the Society will not accept responsibility for any loss.

2. Each typescript should include (i) an abstract of not more than 150 words, preferably containing no formulae, and certainly containing no complicated formulae and no references, (ii) a 1980 Mathematics subject classification (Amer. Math. Soc.) (1985 Revision), (iii) if the title is long, a shortened form of it, no more than forty characters in length, including spaces.

3. The article should be typed or photocopied on high quality A4 or quarto bond paper, on one side only, with at least double spacing, and with a generous margin (at least 3 cm) all around.

4. The conventions of *A* manual for authors of mathematical papers published by the American Mathematical Society should be observed. Two alternative styles for references and quotations are described below. Authors are requested to use one and only one style consistently throughout their typescript.

STYLE 1. A typical reference in the list of references would be:

T. M. Cherry (1965), "Infinite linear systems with homogeneous kernel of degree -1", J. Austral. Math. Soc. 5, 129-168.

A corresponding reference in the text would be: Cherry (1965), p. 155.

STYLE 2. A typical reference in the list of references would be:

[3] L. Grüschen, "Pseudo-quasi-ergodic theorems", J. Austral. Math. Soc. 4 (1960), 2-3.

A corresponding reference in the text would be: Grüschen [3]; or ([3], [4], [9]); or Grüschen ([2, 3]); or Grüschen [3, Theorem 1].

In either style, the list of references should be in alphabetical order of surnames of first authors.

5. Explain clearly what symbols are to be set in special typefaces (such as cursive, fraktur, script, bold) and what is required with any unusual symbols (which should wherever possible be chosen from symbols available to the printer). This is best done by supplying a separate page entitled "Notes to the Compositor" in which conventions and requirements are fully set out, and by drawing attention to particular symbols at their first appearance in the typescript. The Notes to the Compositor should include a list of all symbols and foreign letters used in the article. Be careful to distinguish between similar symbols, such as v, ν , V, k, κ , K, 1, 1, 1, e, ϕ (phi), \emptyset (empty set), ε (epsilon), \in (membership), and so on. Normally 0 will be set as zero; if o or O is required, then show this. Distinguish between inequalities <, > and angular brackets \langle , \rangle .

Journal of the Australian Mathematical Society, Series A — Pure Mathematics and Statistics, is published bimonthly by the Australian Mathematical Society, c/-Department of Mathematics, University of Queensland, St. Lucia, Qld. 4067, Australia. Second class postage is paid at Ann Arbor, Michigan 48106. Postmaster: Send address change notices to Dr B. D. Jones, Australian Mathematical Society, c/-Department of Mathematics, University of Queensland, St. Lucia, Qld. 4067, Australia.

JOURNAL

of the

AUSTRALIAN MATHEMATICAL SOCIETY

Series A — Pure Mathematics and Statistics

Volume 46 1989