

The Association for marine professionals

The Marine Biological Association



"The future now is in the hands of those who understand the real importance of the blue part of the planet"

Dr Sylvia Earle Hon.MBA

Join a global community promoting marine science excellence.

Professional Membership of the MBA provides:

- o use of the post-nominal Mem.MBA
- o *The Marine Biologist* magazine
- o access to an Experts Database to connect with other professionals
- o opportunities to influence marine policy
- o a Certificate of Membership
- o an exclusive MBA pin badge
- o access to research awards and grants

And much more!

Fellow FMBA £120*, Professional Mem.MBA £45*, Associate £40*, Student £25
*£5 Direct Debit discount

Providing an authoritative and independent voice for the marine biological community.

T: +44 (0) 1752 426493 | E: membership@mba.ac.uk | www.mba.ac.uk

@thembauk



PUBLISHED BY THE PRESS SYNDICATE OF THE UNIVERSITY OF CAMBRIDGE
The Pitt Building, Trumpington Street, Cambridge, CB2 1RP, United Kingdom

This journal issue has been printed on FSC-certified paper and cover board. FSC is an independent, non-governmental, not-for-profit organization established to promote the responsible management of the world's forests. Please see www.fsc.org for information.

Printed by Bell and Bain Limited, Glasgow

CAMBRIDGE UNIVERSITY PRESS

Journals Fulfillment Department, UPH, Shaftesbury Road, Cambridge CB2 8BS, United Kingdom
1 Liberty Plaza, Floor 20, New York, NY 10006, USA
477 Williamstown Road, Port Melbourne, VIC 3207, Australia
C/ Orense, 4, planta 13, 28020, Madrid, Spain
Dock House, The Waterfront, Cape Town 8001, South Africa

CONTENTS

REVIEW

- Luna, A., Rocha, F. and Perales-Raya, C. A review of cephalopods (Phylum: Mollusca) of the Canary Current Large Marine Ecosystem (Central-East Atlantic, African coast) 1

ORIGINAL ARTICLES

- Highfield, A., Ward, A., Pipe, R. and Schroeder, D.C. Molecular and phylogenetic analysis reveals new diversity of *Dunaliella salina* from hypersaline environments 27
- Shiomoto, A. and Kamuro, Y. Seasonal variation of water-column light utilization efficiency for primary production in Saroma-ko Lagoon 39
- Harino, H. and Yamato, S. Distribution of antifouling biocides in a coastal area of Tanabe Bay, Japan 49
- Nogueira, M.M., Neves, E. and Johnsson, R. Effects of habitat structure on the mollusc assemblage in *Mussismilia* corals: evaluation of the influence of different coral growth morphology 61
- Dizon, E.G.S., Da-Anoy, J.P., Roth, M.S. and Conaco, C. Fluorescent protein expression in temperature tolerant and susceptible reef-building corals 71
- Oya, Y. and Kajihara, H. Description and phylogenetic relationships of a new genus of Planoceridae (Polycladida, Acotylea) from Shimoda, Japan 81
- Sepahvand, V., Keikhosravi, A. and Kihara, T.C. Population differentiation in the Persian Gulf and Gulf of Oman: insights from the ghost shrimp's associated copepods, *Clausidium iranensis* and *Clausidium persiaensis* 89
- Espinosa-Leal, L., Medellín-Mora, J., Corredor-Acosta, A. and Escibano, R. The community structure of hyperiid amphipods associated with two seamount regions in the South-east Pacific 97
- Prakash, S., Muthu, A. and Kumar, A. Population structure and reproductive performance in the sea anemone associated shrimp *Ancyllocaris brevicarpalis* (Caridea: Palaemonidae) 109
- Salvati, D. da S., Perroca, J.F., Simões, S.M., Castilho, A.L. and Costa, R.C.da Variation in size distribution of juvenile pink shrimps *Farfantepenaeus brasiliensis* and *F. paulensis* in the estuarine-adjacent ocean area of Cananéia, south-eastern coast of Brazil 117
- Fernández, L., García-Soler, C. and Alborés, I. Reproductive strategies under different environmental conditions: total output vs investment per egg in the slipper lobster *Scyllarus arctus* 131
- McLeay, L., Linnane, A., McGarvey, R., Bryars, S. and Hawthorne, P. Response of a southern rock lobster (*Jasus edwardsii*) population to three years of Marine Protected Area implementation within South Australia 141
- Van Volkom, K.S., Harris, L.G. and Dijkstra, J.A. The influence of invasive ascidian diets on the growth of the sea star *Henricia sanguinolenta* 151
- Khalidi, A., Chater, I., Elleboode, R., Mahé, K. and Chakroun-Marzouk, N. Age, growth and mortality of the striped seabream *Lithognathus mormyrus* (Linnaeus, 1758) in the Gulf of Tunis (Central Mediterranean Sea) 159
- Diez, G., Arregi, L., Basterretxea, M., Cuende, E. and Oyarzabal, I. Preliminary observations on abundance and distribution of fish fauna in a canyon of the Bay of Biscay (ICES Division 8c) 169
- Andrade-Tubino, M.F., Salgado, F.L.K., Uehara, W., Utsunomia, R. and Araújo, F.G. *Opsanus beta* (Goode & Bean, 1880) (Acanthopterygii: Batrachoididae), a non-indigenous toadfish in Sepetiba Bay, south-eastern Brazil 179
- Roussos, I. and Megalofonou, P. Ontogenetic shifts and sexual dimorphism in the brain organization of the small-spotted catshark *Scyliorhinus canicula* 189
- Tverin, M., Granroth, J., Abrahamsson, A., Tang, P., Pihlström, H., Lundström, K. and Käkelä, R. Adipose tissue fatty acids suggest spatial and temporal dietary differences in great cormorants of the Baltic Sea area 199
- Harino, H. and Yamato, S. Distribution of antifouling biocides in a coastal area of Tanabe Bay, Japan — Erratum 215
-