

Implementation, Policy and Community Engagement Brief Report

Cite this article: Ridgeway JL, Albertie ML, Williams S, Pantoja E, Stephenson N, Pacheco-Spann L, Ball CT, Radecki Breitkopf C, Bakkum-Gamez JN, Chase L, DeStephano CC, and Sherman ME. Development of novel home-based tampon sampling for endometrial cancer: findings from community-based focus groups with Black and White women. *Journal of Clinical and Translational Science* 5: e125, 1–5. doi: [10.1017/cts.2021.787](https://doi.org/10.1017/cts.2021.787)

Received: 8 December 2020

Revised: 19 April 2021

Accepted: 28 April 2021

Keywords:

Endometrial cancer; health disparities; home testing; formative evaluation; focus groups

Address for correspondence:

J. L. Ridgeway, PhD, Kern Center for the Science of Health Care Delivery, Mayo Clinic, Rochester, MN 55905, USA.

Email: ridgeway.jennifer@mayo.edu

†Deceased.


*These authors are co-senior authors.

© Mayo Clinic, 2021. Published by Cambridge University Press on behalf of The Association for Clinical and Translational Science. This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is unaltered and is properly cited. The written permission of Cambridge University Press must be obtained for commercial re-use or in order to create a derivative work.



Clinical Research
FORUM
Analysis. Advocacy. Action.

Development of novel home-based tampon sampling for endometrial cancer: findings from community-based focus groups with Black and White women

Jennifer L. Ridgeway¹ , Monica L. Albertie², Shantel Williams², Elizabeth Pantoja², Noreen Stephenson², Laura Pacheco-Spann³, Colleen T. Ball³, Carmen Radecki Breitkopf^{4,†}, Jamie N. Bakkum-Gamez⁵, Lori Chase⁶, Christopher C. DeStephano^{7,*} and Mark E. Sherman^{3,*}

¹Robert D. and Patricia E. Kern Center for the Science of Health Care Delivery, Mayo Clinic, Rochester, MN, USA;

²Center for Health Equity and Community Engagement Research, Mayo Clinic, Jacksonville, FL, USA;

³Department of Health Sciences Research, Mayo Clinic, Jacksonville, FL, USA; ⁴Department of Health Sciences Research, Mayo Clinic, Rochester, MN, USA; ⁵Department of Obstetrics and Gynecology, Mayo Clinic, Rochester, MN, USA; ⁶Research Integrator Program, Mayo Clinic, Jacksonville, FL, USA and ⁷Department of Medical and Surgical Gynecology, Mayo Clinic, Jacksonville, FL, USA

Abstract

Novel endometrial cancer (EC) early-detection approaches may reduce racial disparities in mortality. We conducted six community-based focus groups with White and Black women ($N = 57$ participants) in February–March 2020 to explore acceptability of a home-based tampon sampling approach for EC. Participants also completed a survey. Data were analyzed using qualitative content analysis. Awareness of EC and risk factors was low. Acceptability regarding home sampling was high, but participants expressed concerns about instruction complexity and potential risks. Black women reported lower comfort with tampons. Increasing EC awareness, self-efficacy, and familiarization with tampons would advance prospects for at-home sample collection for EC testing.

Introduction

Endometrial cancer (EC) is the most common gynecologic cancer in the USA [1]. Among women over 50 with intact uteri, the incidence of EC among non-Hispanic Black women has surpassed that among non-Hispanic White women [1,2] and their mortality rate is nearly double that of non-Hispanic White women [3]. The prevalence of obesity and diabetes, among the strongest risk factors for EC, is higher for Black women [4,5]. Delays in care seeking, disparities in access, and potential differences in tumor biology are also implicated as causes of racial disparities in mortality [6,7].

Symptom recognition is central to early diagnosis. Natural menopause occurs on average at age 51 years in the USA, but frequent occurrence of episodic bleeding after menopause likely contributes to uncertainties about the definition of post-menopausal bleeding (PMB) associated with EC risk [8], especially given that 90% of women with PMB do not have EC, with potentially higher frequencies among Black women [9]. Patients and caregivers may dismiss PMB as a normal variant and delay care seeking [10,11]. Cost, discomfort, and medical expertise required for diagnostic evaluation may also pose barriers to work-up of PMB [12]. Transvaginal sonography, endometrial biopsy, and hysteroscopy produce discomfort and risk of complications, especially among women with severe obesity. Less invasive self-sampling has shown promise for improving cervical cancer screening [13,14]. However, acceptability of self-sampling in the context of triaging women with PMB for diagnostic work-up is unknown. Acceptability of tampon collection is also unknown; HPV sampling for cervical cancer is done using other types of self-sampling devices for fluid collection. The success of in-clinic “proof-of-principle” studies using vaginal tampon collection, combined with sensitive and specific molecular testing for EC, suggests that home-based self-collection may have utility in diagnostic work-up of PMB [15] and it may increase access and early detection [11]. However, most new testing approaches for early cancer detection fail when deployed in real-world settings, in part because researchers did not engage potential users to understand feasibility and acceptability [16]. We conducted community-based focus groups to explore perceptions of White and Black women, as well as gather formative feedback on a sampling kit designed for use in a pilot study.

Methods and Materials

Setting and Participants

Participants were recruited near Jacksonville, Florida, via flyers in the local academic medical center and through distribution by a local community research advisory group. Eligibility was age 40 years or older and self-report Black or White race. Participants were offered \$50 remuneration. This study was reviewed by a community research advisory board and approved by the affiliated Institutional Review Board (IRB# 19-001140).

Data Collection

Members of the study team with experience in qualitative research and community engagement designed and conducted the focus groups. Black and White women were included in separate focus groups in community locations. Women provided oral consent and completed a brief survey about tampon use and attitudes prior to the discussion (Appendix A). Results on questions related to participant characteristics and attitudes are presented here, to provide context for the focus group findings. The semi-structured moderator guide had three parts: awareness of EC and EC risk factors; experience and impressions of tampon use; and feedback on the proposed tampon collection kit. Participants were provided testing kits and instruction sheets for the third part of the discussion. Focus groups were audio recorded and transcribed verbatim for analysis. Moderators answered questions but did not present formal education on EC, as the intent of the groups was to understand the experiences and impressions of women without educational intervention.

Data Analysis

We tabulated the frequency and percentage of participants by categorical and ordered survey responses. The sample median and interquartile range were used to describe continuous survey responses. Comparisons between Black and White participants were made using Fisher's exact test (categorical responses) and Wilcoxon rank sum test (continuous and ordered responses). We used methods of qualitative content analysis, completed by study team members with relevant and complementary social science and medical research backgrounds. Transcripts were reviewed independently, and then researchers met to discuss impressions and develop a coding framework of *a priori* topics from the interview guide (e.g., knowledge of EC) and inductively derived topics. Transcripts were independently coded by two researchers, and coding differences were discussed and reconciled. Transcripts were entered into qualitative software (NVivo 12.6; QSR International) to facilitate queries for analysis, including exploration of differences between groups.

Results

Six focus groups were held in February and March 2020, three with Black women ($N = 31$) and three with White women ($N = 26$). Participant characteristics are presented in Table 1. Compared to White participants, Black participants were younger and reported a lower frequency of ever having used a tampon. More White participants reported being post-menopausal. No participants in either group reported Hispanic ethnicity.

Participant survey responses on tampon use are summarized in Table 2. Nearly a quarter of all Black participants reported that they had no opinion on tampon attitude questions. Among White

participants, 96% agreed or strongly agreed with the statement "Tampons are generally safe to use," whereas only 60% of Black participants gave these responses. Black participants were also more likely to say that tampons were uncomfortable (26% vs 15% agreed or strongly agreed), and significantly fewer said that they were confident that they could insert a tampon correctly, although a majority in both groups agreed that they could do so.

Qualitative Findings

Qualitative findings are organized in two themes, which are displayed in Table 3 with supporting quotations. Themes represent data from all six focus groups. Potential differences between participants in the Black and White focus groups are highlighted below when the study team agreed that differences were notable.

Awareness of EC and Perceived Risk

Low awareness of EC and symptoms

Awareness of EC was low in all focus groups, with most women saying they had never heard of EC, although two women reported knowing someone with EC. Participants expected that symptoms would include bloating, heavy bleeding or discharge, and pain with intercourse. In terms of menstruation, women spoke about abnormal cycles most frequently, especially more frequent bleeding than expected for a pre-menopausal woman.

Expected risks related to health history and lifestyle

Reported knowledge of EC risk factors was also low. Suspected risk factors included personal or family history of cancer, heavy bleeding at young age, and history of pregnancy or hormone therapy. Participants in all groups discussed endometriosis when asked about EC familiarly. Several Black participants stated that either they or a contact had the condition, and the similarity of the terms led participants to assume they were somehow related. Women also mentioned behavioral risk factors including smoking and drug use, obesity, having multiple sexual partners, having a "stressed" lifestyle, and not attending regular health care visits. Race and ethnicity were mentioned in two groups as potentially being related to EC, including risk related to delays in seeking care among Black women.

Perspectives on EC Testing Using Tampons

Tampon test acceptability was high but tampon use and perceived risks were considerations

The idea of using a tampon for testing had high acceptability (e.g., minimally invasive and convenient), but personal tampon experience was a primary consideration. Older age was reported as potentially associated with lower acceptability because older or post-menopausal women might experience dryness-related discomfort. Noted risks of tampon use included those related to forgetting about tampon placement. In terms of a tampon-based testing procedure, perceived risks included any chemicals that might be used on the tampon. Some women also noted concern related to DNA-based research, which may carry risks related to confidentiality. Discussion of risk perception was especially prevalent among Black participants, who reported that they talked with other women, including family members, about tampons. These conversations were occasionally a source of myths, including association of tampons and sexual activity.

Table 1. Participant characteristics

Characteristic	Black Participants (N = 31)	White Participants (N = 26)	P value
Median age (IQR), years	54 (50, 62) ^a	60 (56, 68)	0.027 ^b
Ethnicity, N (%)			1.00 ^c
Hispanic, Latino or of Spanish origin	0 (0%)	0 (0%)	
Highest level of schooling completed, N (%)			0.33 ^b
8–11 years	1 (3%)	0 (0%)	
12 years or high school	5 (16%)	2 (8%)	
Post high school training other than college	0 (0%)	1 (4%)	
Some college	8 (26%)	4 (15%)	
College graduate	10 (32%)	11 (42%)	
Post-graduate	8 (26%)	8 (31%)	
Occupational status, N (%)			0.35 ^c
Employed	19 (61%)	12 (46%)	
Unemployed	0 (0%)	0 (0%)	
Homemaker	0 (0%)	1 (4%)	
Student	0 (0%)	0 (0%)	
Retired or disabled	12 (39%)	13 (50%)	
Marital status, N (%)			
Married or partnered	9 (29%)	18 (69%)	
Separated or divorced	13 (42%)	6 (23%)	
Widowed	2 (6%)	2 (8%)	
Single, never married	7 (23%)	0 (0%)	
Confidence filling out medical forms alone, N (%)	^a		0.13 ^b
Extremely	19 (63%)	21 (81%)	
Quite a bit	7 (23%)	4 (15%)	
Somewhat	2 (7%)	1 (4%)	
A little bit	2 (7%)	0 (0%)	
Not at all	0 (0%)	0 (0%)	
Health care coverage of any kind, N (%)			1.00 ^c
Yes	30 (97%)	26 (100%)	
No	1 (3%)	0 (0%)	
Ever used a tampon (yes), N (%)	19 (61%)	25 (96%)	0.003 ^c
Menopausal status, N (%)	^a		0.042 ^b
Pre-menopausal	7 (23%)	2 (8%)	
Peri-menopausal	5 (17%)	2 (8%)	
Post-menopausal	18 (60%)	22 (85%)	

^aInformation was not available for 1 participant.^bWilcoxon rank sum test.^cFisher exact test.**Table 2.** Views on tampon use

Characteristic	Black Participants (N = 31)	White Participants (N = 26)	P value ^b
Tampons are generally safe to use, N (%)	^a		<0.001
Strongly agree	2 (7%)	12 (46%)	
Agree	16 (53%)	13 (50%)	
No opinion	7 (23%)	1 (4%)	
Disagree	1 (3%)	0 (0%)	
Strongly disagree	4 (13%)	0 (0%)	
Tampons are uncomfortable to wear/use, N (%)			0.004
Strongly agree	4 (13%)	0 (0%)	
Agree	4 (13%)	4 (15%)	
No opinion	8 (26%)	1 (4%)	
Disagree	13 (42%)	12 (46%)	
Strongly disagree	2 (6%)	9 (35%)	
It is easy to forget that you are wearing a tampon, N (%)			0.61
Strongly agree	0 (0%)	2 (8%)	
Agree	14 (45%)	12 (46%)	
No opinion	9 (29%)	3 (12%)	
Disagree	5 (16%)	8 (31%)	
Strongly disagree	3 (10%)	1 (4%)	
Tampon removal can be painful, N (%)	^a		0.22
Strongly agree	2 (7%)	1 (4%)	
Agree	8 (27%)	7 (27%)	
No opinion	8 (27%)	1 (4%)	
Disagree	10 (33%)	15 (58%)	
Strongly disagree	2 (7%)	2 (8%)	
I am confident I can insert a tampon correctly, N (%)			<0.001
Strongly agree	7 (23%)	18 (69%)	
Agree	16 (52%)	7 (27%)	
No opinion	5 (16%)	1 (4%)	
Disagree	2 (6%)	0 (0%)	
Strongly disagree	1 (3%)	0 (0%)	

^aInformation was not available for 1 participant.^bWilcoxon rank sum test.

Self-efficacy was high but instructions need to be simple and procedures clear

Especially among women with experience using tampons, reported self-efficacy (i.e., confidence in ability to complete home testing) was high. Participant review of the pilot kit highlighted opportunities to make the instructions clearer, simpler, and easier to follow, but some women thought the instructions were overwhelming

Table 3. Qualitative findings by theme

Theme	Subtheme	Supporting quotations
Awareness of EC and perceived risk	<ul style="list-style-type: none"> • Low awareness of EC and symptoms • Expected risks related to health history and lifestyle 	<ul style="list-style-type: none"> • “However, as evidenced by the fact that none of us mentioned – when you asked about what cancers are you concerned about – none of us mentioned endometrial cancer. It is not a cancer that comes to mind . . . it’s not discussed.” • “I knew nothing about it until a year ago. One of our best friends was diagnosed and she lived five months, and so I learned quick it’s not a good cancer to have.” • “Women who are still having their period, having bleeding in between when they thought their periods should be (might signal an EC symptom).” • “And my thing was I’ve met people that had endometriosis. And like the first things they go for- the doctors check for is cancer.” • “(EC risk factor would be) a history of heavy bleeding maybe when she was younger and having periods and painful.”
Perspectives on EC testing using tampons	<ul style="list-style-type: none"> • Tampon test acceptability was high, but tampon use and perceived risks were considerations • Self-efficacy was high, but instructions need to be simple and procedures clear 	<ul style="list-style-type: none"> • “I think a lot of things you hear from your parents, your aunts, but also your peers because some of the peers would hear it from their parents, so they would say that ‘loose’ (girls use tampons).” • “I think the kit is great because it’s very- it’s noninvasive. It’s something you prepare yourself at home. So for people who have anxiety about invasive procedures, testing, this is very simple.” • “Age because the further out you get from using tampons and some being comfortable with the process and maybe being comfortable with your anatomy and whatever’s going on down there because it’s not always, you know- whatever [laughs]. So I think, as you get older, maybe that’s easier for older people, ‘Oh I don’t know if I want to do that.’” • “I would just worry about the integrity of the sample with like temperature. I mean it can be like 100 degrees outside in Florida, and like is it in a mail truck somewhere baking.” • “What’s on that thing that I’m inserting in me, making sure there is no harmful- later on, we hear that solution A is harmful to your health; and so don’t tell me that later on if it’s something that’s just as safe as using a real tampon.”

EC, endometrial cancer.

enough that they expressed a preference for in-clinic tampon collection instead, especially in response to instructions dictating a clean environment, a time range of tampon placement (30 min to 2 h), and the need to store the sample at a specified temperature. Participants expressed concern that mistakes would impact the usefulness of the sample or the accuracy of the results.

Discussion

Risk perception is an important motivator of behaviors [17]. Knowledge of EC risk factors was minimal in this study, including the link with obesity, which is the strongest and most prevalent risk [4]. Although women linked abnormal uterine bleeding (AUB) to EC, concerns focused primarily on frequent or heavier bleeding among pre-menopausal women. High prevalence of irregular bleed secondary to leiomyomata among Black women in particular may lead some patients to “normalize PMB,” which, along with stigma discussing bleeding and inadequate efforts to elicit histories of PMB, may delay care seeking [7,10]. Furthermore, despite low awareness of EC, this study found relatively high familiarity with endometriosis. Presumed connection between these conditions could serve as a barrier to early detection if women associate lack of an endometriosis diagnosis as indicating lower EC risk. These data strongly suggest the need for education to increase awareness of PMB specifically and EC risk factors more broadly.

Currently, work-up for EC diagnosis involves clinical procedures that may be costly, inconvenient, and, in some cases, risky. The science of home-based testing, combined with sensitive and specific molecular testing for EC, makes home-based testing to triage potential cases with pre-menopausal AUB and PMB a compelling strategy to increase access to care if women find it acceptable.

Participants in this study said home-testing might be convenient, but acceptability was associated with personal tampon experience and perceived risk. Tampon use has been found to be lower among Black women than among White women, but the research is limited and has been focused primarily on younger women [18,19]. Further research is needed on perceptions of tampons and tampon use among a broader population, including women closer to post-menopausal ages, especially considering our finding that women were concerned about vaginal dryness common with older age. Future research could also explore whether women view one-time, short-duration tampon use for sample collection differently than regular, repeated use.

Participants also raised concerns about the complexity of testing instructions and procedures. Improved formatting and visuals may alleviate these concerns, as might videos that demonstrate testing procedures or smartphone apps that monitor tampon insertion time. However, it is notable that the participants—most with high educational attainment—reported that in-clinic tampon testing might be preferable if instructions or procedures were too complex. Instructions may also need to detail benefits and demands of all options and safeguards in place for home sampling accuracy. Our pilot study is currently deploying a test kit, refined based on the findings reported here, to further assess feasibility, including sample stability. Further, this study team has pilot-tested markers for benign endometrial DNA, which can provide information about the quality of samples that tested negative for EC-specific markers.

The strengths of this study include community health advisory board review of study procedures and data collection instruments. Analysis by the multidisciplinary study team enriched interpretation and bolsters credibility of the findings. This study also has

limitations. The number of focus groups and participants was suitable for qualitative research on personal experiences. Survey responses provide context for the focus group findings and generate hypotheses for further exploration, but the small number of participants is not suitable for statistical generalizability. Likewise, the use of focus groups with both Black and White women was intended to ensure representation of Black voices and to explore potential differences in attitudes, experiences, and information needs, but findings should not be generalized broadly. These differences should be studied further with larger-scale quantitative methods appropriate for hypothesis testing. Women in this study also had high levels of education and were insured, and none of the participants reported Hispanic ethnicity. Future research is needed to understand the views of women with different levels of education or lack of insurance coverage, as well as to further explore issues of race, ethnicity, and culture. We did not administer a post-focus group survey, so it is not known whether attitudes changed because of the group discussions. Finally, it is important to acknowledge that EC disparities are multi-level and include racial disparities related to factors like clinician bias [20]. Future research should include exploration of health-system barriers that compound EC disparities. This study team is currently establishing a local women's health advisory board to advance this work.

Conclusion

Education about EC and risk factors, especially PMB, is needed to motivate action related to early detection. Using the tampon as a self-collection testing device may be acceptable, but further effort is needed to address the concerns of Black women.

Supplementary Material. To view supplementary material for this article, please visit <https://doi.org/10.1017/cts.2021.787>

Acknowledgements. The study team would like to thank the focus group participants and the members of the community advisory board for their contributions to this study.

This research was funded by a grant from Mayo Clinic Office of Women's Health (M.E.S).

Disclosures. The authors have no conflicts of interest to declare.

References

- Doll KM, Winn AN. Assessing endometrial cancer risk among US women: long-term trends using hysterectomy-adjusted analysis. *American Journal of Obstetrics & Gynecology* 2019; **221**: 318.e311–318.e319.
- Jamison PM, Noone AM, Ries LAG, Lee NC, Edwards BK. Trends in endometrial cancer incidence by race and histology with a correction for the prevalence of hysterectomy, SEER 1992 to 2008. *Cancer Epidemiology Biomarkers & Prevention* 2013; **22**: 233–241.
- DeSantis CE, Miller KD, Goding Sauer A, Jemal A, Siegel RL. Cancer statistics for African Americans, 2019. *CA: A Cancer Journal for Clinicians* 2019; **69**: 211–233.
- Hales CM, Carroll MD, Fryar CD, Ogden CL. *Prevalence of Obesity and Severe Obesity among Adults: United States, 2017–2018*. NCHS Data Brief, no 360. Hyattsville, MD: National Center for Health Statistics, 2020.
- Centers for Disease Control and Prevention. National Diabetes Statistics Report, 2020. Centers for Disease Control and Prevention, U.S. Dept of Health and Human Services. 2020 [cited Mar 29, 2021]. (<https://www.cdc.gov/diabetes/pdfs/data/statistics/national-diabetes-statistics-report.pdf>)
- Collins Y, Holcomb K, Chapman-Davis E, Khabele D, Farley JH. Gynecologic cancer disparities: A report from the Health Disparities Taskforce of the Society of Gynecologic Oncology. *Gynecologic Oncology* 2014; **133**: 353–361.
- Doll KM, Khor S, Odem-Davis K, et al. Role of bleeding recognition and evaluation in Black-white disparities in endometrial cancer. *American Journal of Obstetrics & Gynecology* 2018; **219**: 593.e591–593.e514.
- Astrup K, Olivarius Nde F. Frequency of spontaneously occurring postmenopausal bleeding in the general population. *Acta Obstetrica et Gynecologica Scandinavica* 2004; **83**: 203–207.
- Ghoubara A, Sundar S, Ewies AAA. Black women with postmenopausal bleeding have lower prevalence of endometrial cancer than other ethnic groups. *Climacteric* 2019; **22**: 632–636.
- Clarke MA, Long BJ, Del Mar Morillo A, Arbyn M, Bakkum-Gamez JN, Wentzensen N. Association of Endometrial cancer risk with postmenopausal bleeding in women: a systematic review and meta-analysis. *JAMA Internal Medicine* 2018; **178**: 1210–1222.
- Doll KM, Hempstead B, Alson J, Sage L, Lavallee D. Assessment of pre-diagnostic experiences of Black women with endometrial cancer in the United States. *JAMA Network Open* 2020; **3**: e204954.
- DeStephano CC, Bakkum-Gamez JN, Kaunitz AM, Ridgeway JL, Sherman ME. Intercepting endometrial cancer: Opportunities to expand access using new technology. *Cancer Prevention Research* 2020; **13**: 563–568.
- Sanner K, Wikström I, Strand A, Lindell M, Wilander E. Self-sampling of the vaginal fluid at home combined with high-risk HPV testing. *British Journal of Cancer* 2009; **101**: 871–874.
- Winer RL, Lin J, Tiro JA, et al. Effect of mailed human papillomavirus test kits vs usual care reminders on cervical cancer screening uptake, precancer detection, and treatment: a randomized clinical trial. *JAMA Network Open* 2019; **2**: e1914729.
- Bakkum-Gamez JN, Wentzensen N, Maurer MJ, et al. Detection of endometrial cancer via molecular analysis of DNA collected with vaginal tampons. *Gynecologic Oncology* 2015; **137**: 14–22.
- Walter FM, Thompson MJ, Wellwood I, et al. Evaluating diagnostic strategies for early detection of cancer: the CanTest framework. *BMC Cancer* 2019; **19**: 586.
- Maddux JE, Rogers RW. Protection motivation and self-efficacy: a revised theory of fear appeals and attitude change. *Journal of Experimental Social Psychology* 1983; **19**: 469–479.
- Finkelstein JW, von Eye A. Sanitary product use by white, black, and Mexican American women. *Public Health Reports* 1990; **105**: 491–496.
- Romo LF, Berenson AB. Tampon use in adolescence: differences among European American, African American and Latina women in practices, concerns, and barriers. *Journal of Pediatric and Adolescent Gynecology* 2012; **25**: 328–333.
- Doll KM. Investigating Black-White disparities in gynecologic oncology: theories, conceptual models, and applications. *Gynecologic Oncology* 2018; **149**: 78–83.