

¹⁴C SOURCES AND DISTRIBUTION IN THE VICINITY OF LA HAGUE NUCLEAR REPROCESSING PLANT: PART II—MARINE ENVIRONMENT

D Maro^{1,2} • M Fontugne^{3,4} • C Hatté³ • D Hebert¹ • M Rozet¹

ABSTRACT. Carbon dioxide partial pressure and radiocarbon activity were measured in air and seawater in the Bay of Seine and around the COGEMA-La Hague nuclear reprocessing plant (northwest France) during 3 cruises in 2000 and 2002. Results clearly show that the sea is a source of CO₂ and ¹⁴C to the atmosphere. High ¹⁴C concentrations in air and water related to the La Hague liquid waste are clearly recorded. For the restricted area of the Bay of Seine, CO₂ carbon and ¹⁴C fluxes were estimated, indicating that less than 3% of the liquid ¹⁴C release is introduced in the atmosphere.

INTRODUCTION

The COGEMA-La Hague nuclear reprocessing plant is located in the northwest of the Cotentin Peninsula, near Cherbourg (France). This nuclear plant releases radioelements in the atmosphere and in the English Channel. About 8.5 TBq.yr⁻¹ of radiocarbon is released as liquid waste through a pipe a few km off the shore, west of the reprocessing plant (COGEMA data 2000). Recent studies in the peninsula (Fontugne et al., this issue) show anomalously high ¹⁴C contents in vegetation near the coast which suggest a supplementary marine contribution through the degassing of the ¹⁴C excess supplied by the liquid release of the nuclear plant.

The aim of this study is to estimate the ¹⁴C fluxes between seawater and atmosphere in the northwest part of the Cotentin Peninsula and in the Bay of Seine.

MATERIALS AND METHODS

Methods

In order to estimate ¹⁴CO₂ fluxes across the surface seawater and the atmosphere interface, the partial pressure of carbon dioxide (pCO₂) was calculated using measurements of the total alkalinity and pH of water. According to Henry's law, the difference between pCO₂ in air and water indicates if seawater is a source of CO₂ to the atmosphere.

Carbon dioxide fluxes were calculated using Equation 1:

$$\Phi(\text{CO}_2) \text{ (mole.m}^{-2}\text{.s}^{-1}\text{)} = K \cdot S \cdot \Delta p\text{CO}_2 \quad (1),$$

where K (m.s⁻¹) is the CO₂ transfer coefficient between seawater and the atmosphere, S is solubility of CO₂ (mole.m⁻³.atm⁻¹), and $\Delta p\text{CO}_2$ (atm) is the difference between partial pressure in water and air. pCO₂ in air was considered as a constant mean value of 367 μatm according to values published by Copin-Montégut (1996), Boehme et al. (1998), Frankignoulle and Borges (2001), and Keir et al. (2001). K is a parameter depending on wind speed; its values are available in Liss and Merlivat (1986), Tans et al. (1990), or Wanninkhof and McGillis (1999). As these K values are slightly different in these 3 studies, we present the 3 CO₂ flux estimates.

¹Laboratoire de Radioécologie de Cherbourg-Octeville (IRSN/DEI/SECRE/LRC), F-50130 Cherbourg-Octeville, France.

²Email: denis.maro@irsn.fr.

³Laboratoire des Sciences du Climat et de l'Environnement, UMR 1572-CEA/CNRS, Domaine du CNRS, F-91198-Gif sur Yvette cedex, France.

⁴Corresponding author. Email: Michel.Fontugne@lsce.cnrs-gif.fr.

The exchange of CO₂ between the atmosphere and the surface seawater is an equilibrium process; the net CO₂ flux is the difference between gas going from water to air and gas going from air to water. Both these fluxes carry ¹⁴C at concentrations appropriate to the medium from where they originate, and the net ¹⁴C flux is, once again, the difference. The expression for the net ¹⁴C flux from the ocean surface to the atmosphere is given in Equation 2:

$$\Phi(^{14}\text{C}) = [^{14}\text{C}]_{\text{O}} \Phi_{\text{OA}}(\text{CO}_2) - [^{14}\text{C}]_{\text{A}} \Phi_{\text{AO}}(\text{CO}_2) \quad (2).$$

Regarding the constant factor due to the use of appropriate units, ¹⁴C fluxes were calculated following Equation 3:

$$\Phi(^{14}\text{C}) (\text{Bq.km}^{-2}.\text{d}^{-1}) = 10^9 ([^{14}\text{C}]_{\text{O}} \cdot \Phi_{\text{OA}} - [^{14}\text{C}]_{\text{A}} \Phi_{\text{AO}}) \quad (3),$$

where [¹⁴C]_O and [¹⁴C]_A are the ¹⁴C concentrations (Bq.Kg⁻¹C) in the surface ocean and air, respectively, and Φ_{OA} and Φ_{AO} are the CO₂ fluxes (mole.m⁻².s⁻¹) from ocean to air, and air to ocean, respectively.

Sampling and Analytical Procedures

Seawater and air samples were collected simultaneously during 3 cruises (TE-SEA cruise, 1–4 June 2000; TRANSAT 1 cruise, 24–28 February 2002; and TRANSAT 2 cruise, 27–31 August 2002) around the northern Cotentin Peninsula and the Bay of Seine (Figure 1). During these cruises, krypton-85 (⁸⁵Kr) was measured continuously at a frequency of 1 measurement.s⁻¹ (Maro et al. 2002). The detection of ⁸⁵Kr emitted from the chimney of the reprocessing plant allows air to be sampled for ¹⁴C measurements free of direct contamination coming from the plume of the chimney. CO₂ in a few liters of air was trapped by bubbling air in sodium hydroxide solution. Sodium hydroxide was prepared a few hours before sampling to avoid contamination by sodium carbonate, generally present within sodium hydroxide tablets. In the laboratory, barium hydroxide was added to a sodium hydroxide solution to get a precipitate of barium carbonate. This carbonate was then rinsed with degassed distilled water and dried at 50 °C before reaction with orthophosphoric acid under vacuum to evolve CO₂. Water used to prepare the solution was previously degassed. Precipitation, filtration of precipitate, and rinsing were performed under a controlled nitrogen atmosphere. The blank was determined following this procedure on the same solution used for sampling.

Water samples were collected in glass bottles and poisoned with mercury chloride. In the laboratory, total CO₂ (ΣCO₂) was extracted from seawater following the procedure described by Bard et al. (1988) and Leboucher et al. (1999).

The temperature and salinity of seawater were measured using SBE 19-03 Seabird equipment, and wind direction and speed were recorded. For the TRANSAT cruises, CO₂ partial pressure in surface seawater was calculated using the pH and total alkalinity measurements following the “Standard Operating Procedures SOP3 and SOP6” of the US Department of Energy (DOE 1994).

The ¹⁴C activity was measured at the Laboratoire de Sciences du Climat et de l’Environnement using the Gif accelerator mass spectrometry (AMS) facilities. Results are expressed in Bq.kg⁻¹ C (100 pMC = 226 Bq.kg⁻¹ C). The relative precision varies between 0.5 and 2%.




Figure 1 Location of sampling stations during the TE-SEA, TRANSAT 1, and TRANSAT 2 cruises

RESULTS AND DISCUSSION

Carbon Dioxide Flux

pCO₂ values (Figure 2 a, b) vary from 373 to 614 μatm for the TRANSAT 1 cruise data, and between 423 and 1408 μatm for TRANSAT 2 cruise data. pCO₂ measurements in water show similar high values compared to air during the winter and summer cruises and increase toward the Seine River estuary. This indicates CO₂ flux to the atmosphere since the mean atmospheric pCO₂ value is 367 μatm. These results are easily predictable because coastal, shelf, and estuarine areas exhibit high biological activity due to the nutrient input near the river mouths and organic matter recycling in the water column (Savoye et al. 2003). This heterotrophic activity, resulting in high degradation rates at the sediment surface and in the water column, produces high dissolved CO₂ concentrations (April 1999).

CO₂ fluxes to the atmosphere calculated from Equation 1 are reported in Figure 3 a, b. Carbon fluxes range from very few to 911 and 493 kg C km⁻².d⁻¹ for TRANSAT 1 and TRANSAT 2 cruises, respectively. These values are strongly dependent on the *K* transfer coefficient chosen (Liss and Merlivat [1986], Tan et al. [1990], or Wanninkhof and McGillis [1999]), the difference between estimations may vary by a factor of 2 or more, especially considering high wind speed.




Figure 2 Variation of CO₂ partial pressure in surface seawater versus distance from pipe outlet during TRANSAT 1 and 2 cruises




Figure 3 Variations of CO₂ fluxes versus distance from pipe outlet during TRANSAT 1 (above) and 2 (below) cruises

¹⁴C Flux

Preliminary experiments performed during the TE-SEA cruise are reported in Table 1 and Figure 4. The ¹⁴C activities of ΣCO₂ in seawater vary between 342.2 Bq.kg⁻¹ C at station 11 in the northwest part of the studied area to 580.8 Bq.kg⁻¹ C near the mouth of the pipe. The values (except station 11) range between 479.6 to 580.8 Bq.kg⁻¹ C, decreasing from south to north and recording clearly the plume and the dilution of waste. These values are about twice the reference values in 2000 (248 Bq.kg⁻¹ C) for modern surface waters measured in Brittany, off Brest. ¹⁴C values in air samples are more homogenous, varying between 257.6 to 292.2 Bq.kg⁻¹ C. However, these values are higher than the atmospheric reference (248 Bq.kg⁻¹ C).

Table 1 Location of sampling, distance from the pipe outlet, ¹⁴C activity in seawater and air during the TE-SEA cruise. Shaded rows are under plume influence (see Fontugne et al., these proceedings).

Station	Latitude (N)	Longitude (W)	Distance	¹⁴ C in water	¹⁴ C in air
TE-SEA	(°)	(°)	(km)	(Bq/kg C)	(Bq/kg C)
1	49.70	2.00	5.2	480.7	268.5
2	49.68	1.97	2.2	506.0	279.1
3	49.65	1.96	1.5	527.0	257.6
4	49.63	1.92	4.4	562.3	269.8
5	49.66	1.94	1.2	520.3	259.4
6	49.71	2.00	5.9	479.6	428.9
7	49.66	1.97	0.9	539.7	468.7
8	49.66	1.94	1.4	559.6	435.1
9	49.63	1.91	5.1	557.8	908.1
10	49.64	1.96	2.5	506.2	426.0
11	49.74	2.11	14.0	342.2	271.2
12	49.66	1.96	0.0	580.8	292.2




Figure 4 Variations of ¹⁴C activities (Bq.kg⁻¹ C) in surface seawater and air versus distance from the pipe outlet during the TE-SEA cruise

Table 2 Location of sampling, distance from the pipe outlet, ^{14}C activity in seawater and air, wind speed at 10 m, temperature, salinity, and CO_2 partial pressure in surface seawater during TRANSAT cruises.

Station	Latitude (N) (°)	Longitude (°)	Distance (km)	^{14}C water (Bq/kg C)	u10 (m/s)	Temperature (°C)	Salinity (pm)	p CO_2 (μatm)	^{14}C air (Bq/kg C)
TRANSAT 1									
1	49.73	1.88W	15.0	587.8	12.0	9.7	33.91	373.4	238.8
2	49.70	1.73W	22.4	—	11.0	9.8	33.86	378.6	—
3	49.67	1.63W	29.4	410.2	12.0	9.5	33.49	377.1	239.4
4	49.67	1.62W	29.6	—	11.0	9.5	33.61	386.3	—
5	49.67	1.62W	29.6	—	10.0	9.6	33.55	384.1	—
6	49.67	1.62W	29.6	—	10.0	9.6	33.42	386	—
8	49.67	1.62W	29.6	—	10.0	9.6	33.63	373.5	—
9	49.57	1.24W	58.3	378.4	8.0	8.7	33.01	387.5	237.1
10	49.51	1.03W	74.4	379.5	12.0	9.2	33.43	378.5	239.3
11	49.48	0.93W	82.5	—	9.0	8.8	33.05	402.8	—
12	49.46	0.82W	90.7	353.3	9.0	8.7	32.88	400.7	236.4
13	49.44	0.71W	98.5	—	9.0	8.6	32.41	416.8	—
14	49.43	0.60W	107.2	331.0	10.0	8.5	32.35	415.3	236.2
15	49.42	0.48W	115.5	—	10.0	8.5	32.31	419.9	—
16	49.40	0.35W	125.4	324.8	10.0	8.5	32.17	430.3	241.3
17	49.40	0.17W	137.9	—	10.0	8.5	32.37	421	—
18	49.39	0.09W	143.6	—	10.0	8.4	31.26	441.2	242.7
19	49.44	0.09W	142.8	—	10.0	8.4	31.65	448.6	—
20	49.47	0.04E	151.5	286.3	10.0	8.1	25.63	613.6	236.2
TRANSAT 2									
1	49.70	1.73W	22.40	342.5	7.3	17.9	34.78	465.7	226.4
5	49.42	2.25W	33.7	270.7	6.2	18.0	35.10	422.7	222.2
7	49.64	2.44W	34.4	—	3.4	16.7	35.10	424.0	—
9	49.69	1.96W	3.6	706.7	3.4	17.8	34.89	462.9	346.4
10	50.00	1.94W	37.7	281.6	4.1	17.1	35.06	438.0	240.9
13	50.00	1.01W	83.4	—	3.4	17.7	34.90	442.5	—
17	50.00	0.35E	175.4	343.6	1.8	18.6	34.56	452.0	262.6
18	49.47	0.04E	151.3	258.3	3.1	19.6	29.01	759.0	243.5
19	49.45	0.05E	152.7	—	5.9	19.5	22.40	1407.6	—
34	49.38	0.09W	144.4	—	5.5	19.4	31.66	464.3	—
36	49.46	0.66W	101.7	—	4.8	19.6	33.30	402.0	—
37	49.51	1.04W	74.0	351.5	4.6	18.7	34.35	441.0	262.2

During the TRANSAT cruises, ¹⁴C activities in the surface seawater vary between 258.3 to 706.7 Bq.kg⁻¹ C (Table 2). The highest value corresponds to a station within the plume of the liquid waste near the pipe outlet (Figures 5 a,b). All these values are higher than the reference value of 239 Bq.kg⁻¹ C (year 2002). During TRANSAT 1, winter cruise atmospheric values are near the reference value mainly due to the stormy meteorological conditions which induce a greater mixing of the atmosphere. During the TRANSAT 2 summer cruise, ¹⁴C activities in the air present a positive correlation with ¹⁴C in surface water, confirming transfer from the sea.




Figure 5 Variations of ¹⁴C activities (Bq.kg⁻¹ C) in surface seawater and air versus distance from pipe outlet during TRANSAT 1(above) and 2 (below) cruises

For each estimate of carbon dioxide flux, the ¹⁴C fluxes were estimated using Equation 3. Maximum fluxes are 6.4 10⁵ and 1.7 10⁵ Bq.km⁻².d⁻¹ for TRANSAT 1 and 2, respectively (Figures 6 a,b). During TRANSAT 1, the highest ¹⁴C flux is encountered near the city of Cherbourg, and the lowest between Cherbourg and the Seine River estuary. The TRANSAT 2 cruise presents a maximum value near Cherbourg and a minimum value near the Seine River estuary due to lower CO₂ fluxes during these cruises (low wind speed near the Seine estuary). Following estimates derived from the Tans et al. model, the mean flux in the Bay of Seine due to the COGEMA-La Hague liquid waste would

reach $2.3 \cdot 10^5$ and $3.9 \cdot 10^4$ $\text{Bq}\cdot\text{km}^{-2}\cdot\text{d}^{-1}$ for TRANSAT 1 and 2, respectively. A rough annual estimation would give 216 GBq for the 4400 km^{-2} of the Bay of Seine, representing less than 3% of the liquid release from COGEMA-La Hague nuclear reprocessing plant.




Figure 6 Variations of ^{14}C fluxes versus distance from pipe outlet during TRANSAT 1 (above) and 2 (below) cruises

CONCLUSIONS

Simultaneous measurements of CO_2 partial pressure and ^{14}C activity in air and seawater indicate that the English Channel and Bay of Seine are a source a carbon dioxide to the atmosphere in good agreement with previous studies. ^{14}C activities decrease from the west to the east according to the dilution of the waste plume of the plant. Estimations of CO_2 and ^{14}C flux show that a minor part of the ^{14}C liquid release by industrial activity is recycled to the atmosphere. These estimates are not very accurate due to the variation of CO_2 transfer coefficients proposed by different models. However, the ^{14}C releases in the Bay of Seine could provide a good opportunity to perform new experiments in order to get better estimates of CO_2 transfer coefficients between water and atmosphere.

ACKNOWLEDGEMENTS

Thanks are due to M Paterne for helpful discussions and M Arnold for AMS measurements and S Le Bar and M L Fitament from COGEMA for their help during the cruises. We are very grateful to Rodger Sparks for suggestions and corrections of the manuscript. We thank Mr Le Bourhis and Mr Henri captains of the crew of the R/V “Côtes de la Manche.” TE-SEA and TRANSAT cruises were supported by CNRS.LSCE contribution nr 1102.

REFERENCES

- Abril G. 1999. Dynamique du carbone dans les estuaires européens: processus de minéralisation et transfert continent-océan-atmosphère [PhD dissertation]. Université de Bordeaux I. p 173–209.
- Bard E, Arnold M, Ostlund HG, Maurice P, Monfray P, Duplessy JC. 1988. Penetration of bomb radiocarbon in the tropical Indian Ocean measured by means of accelerator mass spectrometry. *Earth and Planetary Sciences Letters* 87:379–89.
- Boehme SE, Sabine CL, Reimers CE. 1998. CO₂ fluxes from a coastal transect: a time-series approach. *Marine Chemistry* 63:49–67.
- COGEMA. 2000. Surveillance trimestrielle de l'environnement de la Hague. Rapport Hag. 055000120013. 84 p.
- Copin-Montégut G. 1996. *Chimie de l'eau de mer*. Paris: Institut Océanographique. 319 p.
- Department of Energy (DOE). Dickson AG, Goyet C, editors. 1994. *Handbook of Methods for the Analysis of the Various Parameters of the Carbon Dioxide System in Seawater*. Version 2. ORNL/CDIAC-74.
- Fontugne M, Maro D, Baron Y, Hatté C, Hébert D, Douville E. Sources and distribution of radiocarbon in the vicinity of La Hague nuclear reprocessing plant: part I—terrestrial environment. *Radiocarbon*, these proceedings.
- Frankignoulle M, Borges AV. 2001. European continental shelf as a significant sink for atmospheric carbon dioxide. *Global Biogeochemical Cycles* 15(3):569–76.
- Keir RS, Rehder G, Frankignoulle M. 2001. Partial pressure and air-sea flux of CO₂ in the Northeast Atlantic during September 1995. *Deep Sea Research Part II: Topical Studies in Oceanography* 48(14–15):3179–89.
- Leboucher V, Orr J, Jean-Baptiste P, Arnold M, Monfray P, Tisnerat-Laborde N, Poisson A, Duplessy JC. 1999. Oceanic radiocarbon between Antarctica and South Africa along WOCE section 16 at 30°E. *Radiocarbon* 41(1):51–73.
- Liss PS, Merlivat L. 1986. Air-sea exchange rates: introduction and synthesis. In: Buat-Ménard P, editor. *The Role of Air-Sea Exchange in Geochemical Cycling*. Hingham, Massachusetts: D Riedel Publishing Co. p 113–27.
- Maro D, Crabol B, Germain P, Baron Y, Hebert D, Bouisset P. 2002. A study of the near field atmospheric dispersion of emission at height: Comparison of Gaussian plume models (Doury, Pasquill-Briggs, Caire) with krypton-85 measurements taken around La Hague nuclear reprocessing plant. *Radioprotection* 37(C1):1277–82.
- Savoye N, Aminot A, Tréguer P, Fontugne M, Naullet N, Kérouel R. 2003. Dynamics of particulate organic matter δ¹⁵N and δ¹³C during spring phytoplankton blooms in a microtidal ecosystem (Bay of Seine, France). *Marine Ecology Progress Series* 255:27–41.
- Tans PP, Fung IY, Takahashi T. 1990. Observational constraints on the global atmospheric CO₂ budget. *Science* 247:1431–38.
- Wanninkhof R, McGillis WR. 1999. A cubic relationship between air-sea CO₂ exchange and wind speed. *Geophysical Research Letters* 26(13):1889–92.