

resistant to the empiric treatment with piperacillin-tazobactam, but the patient recovered after switching the antibiotic regimen to meropenem. All strains were determined to be identical by amplified-fragment length polymorphism and whole-genome multi-locus sequencing typing (genotype A). New cases occurred, despite the introduction of contact isolation of positive and contact patients. Therefore, weekly point-prevalence screening was introduced, in which more newly colonized patients were identified in the subsequent weeks. Attention to hand hygiene was enforced, and the hypothesis of contamination from “wet” environmental locations was tested by performing cultures of sinks and shower drains. In June and July, 47 of 241 environmental cultures (19.5%) were positive for *E. cloacae* with an identical antibiogram, among which some were typed as genotype A. To diminish the environmental contamination, all siphons of sinks were replaced, and disinfection of sinks and shower drains was intensified using chlorine and soda on a daily basis. Replacement of shower drains was not possible. After this intervention, the incidence of newly colonized patients declined gradually. A change in the regimen of selective gut decontamination in hematology patients was considered as an alternative intervention, but with the decrease in new patient cases, this was not implemented. A final round of environmental cultures at the end of August revealed 8 positive cultures, of which 5 were positive for genotype A. In retrospect, this finding could be explained by the fact that the cleaning team did not follow the intensified instructions for disinfection. From week 29, genotype A *E. cloacae* was no longer cultured in weekly patient screenings. Based on this observation, it is important that in (re)building plans for hospitals, a master plan for the prevention of nosocomial transmission from environment to patients is incorporated.

Funding: None

Disclosures: None

Doi:10.1017/ice.2020.1012

Presentation Type:

Poster Presentation

Rotavirus Vaccination in the NICU: Where Are We? A Rapid Review of Recent Evidence

Mélanie Sicard, CHU Sainte-Justine, University of Montreal; Kristina Bryant, University of Louisville; Martha Muller, University of New Mexico; Caroline Quach, CHU Sainte Justine

Background: Rotavirus is a leading cause of viral acute gastroenteritis (AGE) in infants. Neonates hospitalized in neonatal intensive care units (NICUs) are at risk of rotavirus infections with severe outcomes. The administration of rotavirus vaccines is only recommended, in the United States and Canada, upon discharge from the NICU despite rotavirus vaccine being proven safe and effective in these populations, due to risks of live-attenuated vaccine administration in immunocompromised patients and theoretical risks of rotavirus vaccines strains shedding and transmission. We summarized recent evidence regarding rotavirus vaccines administration in the NICU setting and safety of rotavirus vaccines in preterm infants. **Methods:** We conducted a rapid review of the literature from the past 10 years, searching Medline and Embase, including all study types except reviews, reporting on rotavirus vaccine 1 and rotavirus vaccine 5; NICU setting; shedding or transmission; and/or safety in preterm. One reviewer performed data extraction and quality assessment. **Results:** In total, 31 articles were analyzed. Vaccine-derived virus shedding following rotavirus vaccination existed for nearly all

infants, mostly during the first week after dose 1, with rare transmission described only in the household setting. No case of transmission in the NICU was reported. Adverse events were mild to moderate, occurring in 10%–60% of vaccinated infants. Extreme premature infants or with underlying gastrointestinal failure requiring surgery presented more severe adverse events. **Conclusions:** Recommendations regarding rotavirus vaccine administration in the NICU should be reassessed in light of the relative safety and absence of transmission of rotavirus vaccine strains in the NICU.

Funding: None

Disclosures: Sicard Mélanie: I reference the use of rotavirus vaccines in the NICU setting, which is not recommended; I discuss possible reassessment of these recommendations.

Doi:10.1017/ice.2020.1013

Presentation Type:

Poster Presentation

Rubella Outbreak in Heballi Agasi Ward, Dharwad District, Karnataka, India, 2014–2015

Sree Kalpana Mohankumar, Chief Ministers Comprehensive Health Insurance; Vishweshwarayya Hiremath, District Surveillance unit, Dharwad; Rajashree Koppad, State Surveillance Unit, Bangalore

Background: Countries that have good rubella surveillance, report ~10,000–20,000 rubella cases annually. In India, not many cases of rubella are reported. The Heballi Agasi ward of Dharwad district in Karnataka state, India, reported rubella cases on the last week of January 2015. **Objective:** We investigated the outbreak by time, place, person, and clinical symptoms. **Methods:** We performed a cross-sectional study. We defined a case as any resident of Heballi Agasi who had fever and rash, with or without lymphadenopathy, arthralgia, conjunctivitis, coryza, and cough, after December 15, 2014. We collected sociodemographic details and clinical symptoms of patients. We collected 5 serum samples and sent them to the National Measles Laboratory, Bangalore. We tested for measles and rubella antibodies. We drew an epidemic curve and a spot map. We computed mean age of cases, and we calculated attack rates by mean age and gender. We calculated proportions to describe clinical symptoms, and we interviewed stakeholders regarding rubella vaccination. We continued surveillance until March 2015. **Results:** The population of Heballi Agasi was 1,458. We identified 15 rubella cases (9 girls and 6 boys). The outbreak lasted between December 10, 2014, and February 21, 2015, with a peak on January 16, 2015. The overall attack rate was 1% (15 of 1,458). The mean age of the cases was 6 years (range, 1–23). The attack rate was high (7.7%) among those aged 1–6 years (11 of 143). The attack rate among those aged >6 years was 0.3% (4 of 1,315). In addition to fever and rash, 93% of cases (14 of 15) had coryza, 47% had cough (7 of 15), and 40% had conjunctivitis (6 of 15). Lymphadenopathy was present in only 1 case (1 of 15), and arthralgia was absent among all 15 cases. There was no death among the cases. All 5 sera were positive for rubella and negative for measles. Rubella vaccination was not given for any of the cases because no rubella vaccination is provided in the routine immunization program. **Conclusions:** There was a rubella outbreak in Heballi Agasi ward. Children aged 1–6 years were most affected. We recommend rubella vaccination in the routine immunization.

Funding: None

Disclosures: None

Doi:10.1017/ice.2020.1014