## **General Notes**

		Offered by ARTMENT OF OTO Aichigan Medical So	-	ICHIGAN
	Cours	se Director: John L	Kemink, M.D.	
1989:	2428 April:	22-26 May:	16-20 October:	13–17 November
Mornings	d in a manner relevant for and afternoons are spen	or the otologic surgeo t in the dissection lab	n, utilizing lectures, vid oratory, giving extensiv	leotapes, and dissection ve drilling experience in
Mornings temporal bo provided.		or the otologic surgeo t in the dissection lab	n, utilizing lectures, vid oratory, giving extensiv	ve drilling experience ir

#### CONFERENCE ON THE EUSTACHIAN TUBE AND MIDDLE EAR DISEASES

#### GENEVA, SWITZERLAND

#### 26-29 OCTOBER, 1989

For futher information please contact:

Professor Jacob Sade, c/o The Secretariat, Conference on The Eustachian Tube and Middle Ear Diseases, Intercontinental Hotel Geneva, 7-9 CH. du Petit Saconnex, CH-1211 Geneva 19, Switzerland. **GENERAL NOTES** 

#### VIII ISIAN INTERNATIONAL SYMPOSIUM ON

#### INFECTION AND ALLERGY OF THE NOSE

#### 11-15 JUNE, 1989

#### Sponsored by

#### THE JOHNS HOPKINS UNIVERSITY SCHOOL OF MEDICINE AND

#### THE TRIOLOGIC SOCIETY

#### Stouffer Harborplace Hotel

#### Baltimore, Maryland U.S.A.

For additional information, Abstract submission, and registration, contact:

Congress Secretariat, Office of Continuing Education, The Johns Hopkins Medical Institutions, Turner 22, 720 Rutland Avenue, Baltimore, Maryland 21205 U.S.A.

(301) 955-2959

#### 7<sup>TH</sup> INTERNATIONAL COURSE IN FUNCTIONAL CORRECTIVE NASAL SURGERY 28 May - 2 June, 1989 Utrecht, The Netherlands From 28 May – 2 June, 1989 a course in functional corrective nasal surgery will be given at the University of Utrecht, The Netherlands, under the direction of Prof. E. H. Huizing. The course will consist of lectures, TV-demonstrations, cadaver dissection and live TV-surgery. Teacher Board: Prof W. Pirsig (Ulm), Prof R. Th. R. Wentges (Nijmegen), Prof S. Hellmich (Dortmund), Prof E. B. Kern (Rochester), Prof G. Rettinger (Erlangen), Dr A. F. van Olphen (Utrecht), Dr J. A. M. de Groot (Utrecht) and the Staff of the Department of O.R.L. of the Utrecht University Hospital. Language: English. Information: Prof Dr E. H. Huizing, Department of O.R.L., University Hospital Utrecht, P.O. Box 16250, 3500 CG Utrecht, The Netherlands, Tel: 31 30 373024

129

#### COCHLEAR IMPLANTS

A course for professionals run by UCH/RNID programme

#### 21-23 February, 1989

at

The Institute of Laryngology and Otology, London

Course fee: £150 including lunches, tea and evening reception (reductions available for students etc.)

Write for details and application form to:

The Course Secretary, UCH/RNID Cochlear Implant Programme, Royal Ear Hospital, Huntley Street, London WC1

or phone: 01-387 9300 extension 8761 or 01-387 8033 extension 215

#### ANNOUNCEMENT

3rd Course on Endonasal Surgery of the Paranasal Sinuses, February 10–11, 1989 in Fulda, West Germany. Information:

Prof. Dr W. Draf, Head of the Department for ENT-Diseases and Facial Plastic Surgery, Pacelliallee 4, D-6400 Fulda , Telephone 0661-84481, Telex 49808

#### 4<sup>TH</sup> ASEAN OTORHINOLARYNGOLOGICAL HEAD & NECK CONGRESS SINGAPORE

The 4th ASEAN Otorhinolaryngological Head & Neck Congress will be held from 6-9 April 1989, in Singapore.

Programme includes Plenary Lectures, Panel Discussions, Symposia, "How I do it" sessions, and Free Papers.

For further information, contact

Dr R. E. Stanley, Organizing Secretary, c/o ENT Department, Singapore General Hospital, Outram Road, Singapore 0316, Republic of Singapore.

#### VI ALL INDIA TEMPORAL BONE DISSECTION

#### AND

#### ADVANCED MICRO EAR SURGERY DEMONSTRATION COURSE

#### 22 - 25 FEBRUARY, 1989

Faculty: Dr K. K. Ramalingam, M.B., F.R.C.S., D.L.O.,
Dr A Mahadevaiah, M.B.B.S., D.A.B.O.(U.S.A.)., F.A.A.O.(U.S.A.),
Dr P. G. Viswanathan, M.S., F.R.A.C.S., D.L.O.,
Dr V. S. Siva Sundar, M.S., D.O.R.L.

Under the auspices of the above Foundation, this course is being arranged and will deal mainly with practical and technical aspects of ear operations like various types of Mastoidectomies, Tympanoplasties, Ossicular reconstructions, Stapedectomy, Facial Nerve surgery and inner ear surgery. The course will consist mainly of live demonstration of various operations using closed circuit television and also demonstration by recorded and edited videos. The participants will do temporal bone dissection themselves in the Temporal Bone Laboratory attached to ENT Nursing Home. There will be round table discussion on the various operations and related topics between the participants and Faculty members. Hostel type accommodation, if required, will be arranged. Microscopes, instruments and temporal bones (both dry and wet) will be supplied.

Course Fee: Participant fee \$40; Observer fee \$25.

This includes course material, lunch, tea/coffee and one day Banquet.

#### GRAY'S INN ROAD ENT FELLOWSHIP COURSE

27 February - 10 March, 1989

(For students taking the Final F.R.C.S. in Otolaryngology)

This two week, full-time course is limited to 20 senior students who are about to take the final F.R.C.S. In addition to a series of lectures/demonstrations at the appropriate level, both in the specialty and in related branches of general surgery, the course will include a number of clinical and viva voce sessions, and the setting and correction of test papers.

Course Tutors: Mr C. M. Bailey & Mr C. B. Croft

COURSE FEE: £350

(Meals and refreshments not included apart from the Course Dinner)

All applications must be accompanied by the course fee. Cheques should be made payable to University College London and sent together with the completed application form to:

Cheryl Overington, Secretary to the Dean, The Institute of Laryngology and Otology, 330-332 Gray's Inn Road, London WC1X 8EE.

Telephone: 01-837 8855 Ext. 4218

·

# Instructions to Authors

**Historical Articles**. Articles of this type are generally encouraged, but it is obvious that they have to provide some new information or interpretation, whether it be about a well-known person or for example an instrument associated with him, but those from within a Hospital's own department who have made a hitherto less well-known contribution would be welcome.

Letters to the Editor. This feature is to be re-introduced to give those who wish to comment about a paper previously published within the Journal, an opportunity to express their views. Wherever possible, we shall ask the orginal author to add his further comment, thereby adding to the value of the contribution. Such letters should be sent as soon as possible after publication of each month's issue of the Journal in the hope of including them as soon as possible thereafter.

'Mini-papers, such as those which appear in the British Medical Journal, Lancet, or New England Journal of Medicine, will not be acceptable except on the rare occasion that they bring information of immediate interest to the reader.

**Pathology.** It is proposed to present on a monthly or bi-monthly basis, articles which are pathological interest with particular emphasis on the way the diagnosis was achieved, the techniques involved, etc. It is hoped that clinicians will involve their pathologists in these reports, and all illustrations must be of first-class quality. Where a department, particularly those overseas, may not have such specialized facilities, it may be possible, if a block or several unstained sections are provided for our Adviser to produce or supplement the illustration submitted. Only contributions of particular pathological interest will be accepted. **Radiology.** It is proposed to introduce on a monthly or bi-monthly basis, short reports on cases in which the radiology has been crucial in the making of the diagnosis or the management of a particular case. This spot is to encourage clinicians and radiologists to produce material of particular interest in the specialty and to encourage co-operation in this field. Only presentations with first-class illustrations can be accepted and these must emphasise a problem of unusual clinical interest.

### **Check List for Authors/Secretaries**

1. Title page—Titles should be short with names of the authors, higher degrees only and the city/country. Details of the departments in which the authors work should be put lower down.

An address for correspondence should be supplied together with the author who should receive this and this will ultimately appear beneath the list of references. If the paper was presented at a meeting, the details of this must be given and will be inserted at the bottom of the first page of the printed script.

- 2. Abstract—No paper will be accepted without this and it adds considerably to the Editor's time to have to write and request this if the paper is accepted.
- 3. To follow the Instructions to Authors with the way in which the paper is set out. It is preferred that each section should start on a fresh page with double spacing and wide margins.
- 4. References must be in the Harvard system; to submit a paper using the Vancouver system is automatically to have it returned or rejected.
- 5. Two sets of illustrations must be included, one of half-plate size and the other with the width of 80 mm. Illustrations must be clearly labelled with the author's name on the reverse side and where appropriate with an arrow to give orientation.
- 6. Authors to check manuscript and references to see that these match up particularly for dates and spelling.
- 7. Titles of Journals must be given in full with the date, volume number and first and *last* pages.
- 8. Consent to be obtained from a patient if a photograph of their face is to be reproduced.
- 9. If the author to whom correspondence is to be directed changes his address, he should let the Editorial Office know as soon as possible.
- 10. Page proofs will normally be sent out three months in advance and must be returned as soon as possible.

### The Journal of Laryngology and Otology

Vol 103

January 1989

Instructions to Authors	Cover
Main Articles	4
Tuning forks: the problem of striking: <i>J. Samuel and E. Eitelberg</i>	-
Clinical role of informed tests of hearing: G. G. Browning, I. R. C. Swan and K. K. Chew.	7
Genetic counselling for isolated hearing loss: V. E. Newton	12
Organogenesis of the middle ear structures: <i>B. Ars</i>	16
Does severe water deprivation affect the inner ear?: <i>M. Barbara, D. Bagger-Sjöback, H. Rask-Anderson and R. Filipo</i>	22
Myotic infection of the ear (otomycosis): K. O. Paulose, S. Al Khalifa, P. Shenoy and R. K. Sharma	30
Validity of tympanometry in cases of confirmed otosclerosis: C. Muchnik, M. Hildesheimer, M. Rubinstein and Y. Gleitman	36
Repair of tympanic membrane perforations using hyaluronic acid: An alternative to myringoplasty: <i>Lars-Eric Stenfors</i>	39
Bone pâté repair of the eroded incus: <i>N. B. Solomons and J. M. Robinson .</i>	41
Obliteration of the middle ear cleft: N. S. Violaris and R. Donovan	43
Ototoxicity of ear drops in patients suffering from chronic otitis media: <i>L. Podosthin, M. Fradis and J. Ben David</i>	46
Strategy and benefits of acoustic neuroma searching: D. A. Moffat, D. G. Hardy and D. M. Baguley	51
Role of post-operative electroneuronography in predicting facial nerve recovery after acoustic neuroma removal: <i>G. R. Croxson, D. A. Moffatt, D. G. Hardy and D. M. Baguley</i> .	60
Evaluation of recent experience in hypoglossal-facial nerve anastomosis in the treatment of facial palsy: <i>M. I. Clayton, R. P. Rinon, D. R. Hanson and J. D. Fenwick</i>	63
Treatment of glue ear in relation to radiographic palatal airway size: a predictor for outcome following adenoidectomy?: <i>A. J. Parker and A. R. Maw</i>	66
Effect of respiration on nasopharyngeal radiographs when assessing adenoidal enlargement: <i>P. D. Britton</i>	71
Clinical Records Cholesteatoma of the external auditory canal in hemifacial hypertrophy (hyperplasia): <i>A. K. Gadre and K. C. Gadre</i> Morphology of a columellaform stapes: <i>C. Vande Maele, P. Van Cauwenberge, P.</i> <i>Kluyskeris and L. Vakaet</i> Localized middle ear effusions: Three unusual acute cases in adults: <i>D. A. Parker, A. P. Reid</i>	74 79
Cholesteatoma of the external auditory canal in hemifacial hypertrophy (hyperplasia): <i>A. K. Gadre and K. C. Gadre</i> Morphology of a columellaform stapes: <i>C. Vande Maele, P. Van Cauwenberge, P.</i>	
<ul> <li>Cholesteatoma of the external auditory canal in hemifacial hypertrophy (hyperplasia): A. K. Gadre and K. C. Gadre</li> <li>Morphology of a columellaform stapes: C. Vande Maele, P. Van Cauwenberge, P. Kluyskeris and L. Vakaet</li> <li>Localized middle ear effusions: Three unusual acute cases in adults: D. A. Parker, A. P. Reid and P. J. Patel</li> <li>Carcinoid tumour of the middle ear: H. Kodama, H. Takezawa, T. Suzuki, K. Kobayashi, A. Kataura, J. Wakabayashi and M. Satoh</li> </ul>	79
<ul> <li>Cholesteatoma of the external auditory canal in hemifacial hypertrophy (hyperplasia): A. K. Gadre and K. C. Gadre</li> <li>Morphology of a columellaform stapes: C. Vande Maele, P. Van Cauwenberge, P. Kluyskeris and L. Vakaet</li> <li>Localized middle ear effusions: Three unusual acute cases in adults: D. A. Parker, A. P. Reid and P. J. Patel</li> <li>Carcinoid tumour of the middle ear: H. Kodama, H. Takezawa, T. Suzuki, K. Kobayashi, A. Kataura, J. Wakabayashi and M. Satoh</li> <li>Primary cholesteatoma of the mastoid presenting as cervical abscess: A. A. El-Kholy and J. R. Cherry</li> </ul>	79 83 86 92
<ul> <li>Cholesteatoma of the external auditory canal in hemifacial hypertrophy (hyperplasia): A. K. Gadre and K. C. Gadre</li> <li>Morphology of a columellaform stapes: C. Vande Maele, P. Van Cauwenberge, P. Kluyskeris and L. Vakaet</li> <li>Localized middle ear effusions: Three unusual acute cases in adults: D. A. Parker, A. P. Reid and P. J. Patel</li> <li>Carcinoid tumour of the middle ear: H. Kodama, H. Takezawa, T. Suzuki, K. Kobayashi, A. Kataura, J. Wakabayashi and M. Satoh</li> <li>Primary cholesteatoma of the mastoid presenting as cervical abscess: A. A. El-Kholy and J. R. Cherry</li> <li>Transpalatal approach to the petrous apex: AE. Davis and P. E. Robin</li> </ul>	79 83 86 92 94
<ul> <li>Cholesteatoma of the external auditory canal in hemifacial hypertrophy (hyperplasia): A. K. Gadre and K. C. Gadre</li> <li>Morphology of a columellaform stapes: C. Vande Maele, P. Van Cauwenberge, P. Kluyskeris and L. Vakaet</li> <li>Localized middle ear effusions: Three unusual acute cases in adults: D. A. Parker, A. P. Reid and P. J. Patel</li> <li>Carcinoid tumour of the middle ear: H. Kodama, H. Takezawa, T. Suzuki, K. Kobayashi, A. Kataura, J. Wakabayashi and M. Satoh</li> <li>Primary cholesteatoma of the mastoid presenting as cervical abscess: A. A. El-Kholy and J. R. Cherry</li> <li>Transpalatal approach to the petrous apex: AE. Davis and P. E. Robin</li> <li>Traumatic CSF fistula presenting late as a middle ear effusion: J. T. Brandrick</li> </ul>	79 83 86 92
<ul> <li>Cholesteatoma of the external auditory canal in hemifacial hypertrophy (hyperplasia): A. K. Gadre and K. C. Gadre</li> <li>Morphology of a columellaform stapes: C. Vande Maele, P. Van Cauwenberge, P. Kluyskeris and L. Vakaet</li> <li>Localized middle ear effusions: Three unusual acute cases in adults: D. A. Parker, A. P. Reid and P. J. Patel</li> <li>Carcinoid tumour of the middle ear: H. Kodama, H. Takezawa, T. Suzuki, K. Kobayashi, A. Kataura, J. Wakabayashi and M. Satoh</li> <li>Primary cholesteatoma of the mastoid presenting as cervical abscess: A. A. El-Kholy and J. R. Cherry</li> <li>Transpalatal approach to the petrous apex: AE. Davis and P. E. Robin</li> </ul>	79 83 86 92 94
<ul> <li>Cholesteatoma of the external auditory canal in hemifacial hypertrophy (hyperplasia): A. K. Gadre and K. C. Gadre</li> <li>Morphology of a columellaform stapes: C. Vande Maele, P. Van Cauwenberge, P. Kluyskeris and L. Vakaet</li> <li>Localized middle ear effusions: Three unusual acute cases in adults: D. A. Parker, A. P. Reid and P. J. Patel</li> <li>Carcinoid tumour of the middle ear: H. Kodama, H. Takezawa, T. Suzuki, K. Kobayashi, A. Kataura, J. Wakabayashi and M. Satoh</li> <li>Primary cholesteatoma of the mastoid presenting as cervical abscess: A. A. El-Kholy and J. R. Cherry</li> <li>Transpalatal approach to the petrous apex: AE. Davis and P. E. Robin</li> <li>Traumatic CSF fistula presenting late as a middle ear effusion: J. T. Brandrick</li> <li>Recurrent meningitis associated with meningioma of the mastoid cavity: R. J. Kelleher, J.</li> </ul>	79 83 86 92 94 97
<ul> <li>Cholesteatoma of the external auditory canal in hemifacial hypertrophy (hyperplasia): <i>A. K. Gadre and K. C. Gadre</i></li> <li>Morphology of a columellaform stapes: <i>C. Vande Maele, P. Van Cauwenberge, P. Kluyskeris and L. Vakaet</i></li> <li>Localized middle ear effusions: Three unusual acute cases in adults: <i>D. A. Parker, A. P. Reid and P. J. Patel</i></li> <li>Carcinoid tumour of the middle ear: <i>H. Kodama, H. Takezawa, T. Suzuki, K. Kobayashi, A. Kataura, J. Wakabayashi and M. Satoh</i></li> <li>Primary cholesteatoma of the mastoid presenting as cervical abscess: <i>A. A. El-Kholy and J. R. Cherry</i></li> <li>Transpalatal approach to the petrous apex: <i>AE. Davis and P. E. Robin</i></li> <li>Traumatic CSF fistula presenting late as a middle ear effusion: <i>J. T. Brandrick</i></li> <li>Recurrent meningitis associated with meningioma of the mastoid cavity: <i>R. J. Kelleher, J. A. M. Murray and P. D. Welsby</i></li> <li>Temporal bone venous anomaly of surgical significance: <i>J. C. Shotton, H. Ludman and T.</i></li> </ul>	79 83 86 92 94 97 99
<ul> <li>Cholesteatoma of the external auditory canal in hemifacial hypertrophy (hyperplasia): <i>A. K. Gadre and K. C. Gadre</i></li> <li>Morphology of a columellaform stapes: <i>C. Vande Maele, P. Van Cauwenberge, P. Kluyskeris and L. Vakaet</i></li> <li>Localized middle ear effusions: Three unusual acute cases in adults: <i>D. A. Parker, A. P. Reid and P. J. Patel</i></li> <li>Carcinoid tumour of the middle ear: <i>H. Kodama, H. Takezawa, T. Suzuki, K. Kobayashi, A. Kataura, J. Wakabayashi and M. Satoh</i></li> <li>Primary cholesteatoma of the mastoid presenting as cervical abscess: <i>A. A. El-Kholy and J. R. Cherry</i></li> <li>Transpalatal approach to the petrous apex: <i>AE. Davis and P. E. Robin</i></li> <li>Traumatic CSF fistula presenting late as a middle ear effusion: <i>J. T. Brandrick</i></li> <li>Recurrent meningitis associated with meningioma of the mastoid cavity: <i>R. J. Kelleher, J. A. M. Murray and P. D. Welsby</i></li> <li>Temporal bone venous anomaly of surgical significance: <i>J. C. Shotton, H. Ludman and T. C. S. Cox</i></li> </ul>	79 83 86 92 94 97 99 101
<ul> <li>Cholesteatoma of the external auditory canal in hemifacial hypertrophy (hyperplasia): <i>A. K. Gadre and K. C. Gadre</i></li> <li>Morphology of a columellaform stapes: <i>C. Vande Maele, P. Van Cauwenberge, P. Kluyskeris and L. Vakaet</i></li> <li>Localized middle ear effusions: Three unusual acute cases in adults: <i>D. A. Parker, A. P. Reid and P. J. Patel</i></li> <li>Carcinoid tumour of the middle ear: <i>H. Kodama, H. Takezawa, T. Suzuki, K. Kobayashi, A. Kataura, J. Wakabayashi and M. Satoh</i></li> <li>Primary cholesteatoma of the mastoid presenting as cervical abscess: <i>A. A. El-Kholy and J. R. Cherry</i></li> <li>Transpalatal approach to the petrous apex: <i>AE. Davis and P. E. Robin</i></li> <li>Traumatic CSF fistula presenting late as a middle ear effusion: <i>J. T. Brandrick</i></li> <li>Recurrent meningitis associated with meningioma of the mastoid cavity: <i>R. J. Kelleher, J. A. M. Murray and P. D. Welsby</i></li> <li>Temporal bone venous anomaly of surgical significance: <i>J. C. Shotton, H. Ludman and T. C. S. Cox</i></li> <li>Aneurysm of the anterior inferior cerebellar artery mimicking an intracanalicular acoustic neuroma: <i>M. J. Gleeson, T. C. S. Cox and A. J. Strong</i></li> <li>Mycotic aneurysm of the petrous portion of the internal carotid artery: <i>J. Samuel and C. M.</i></li> </ul>	79 83 86 92 94 97 99 101 107
<ul> <li>Cholesteatoma of the external auditory canal in hemifacial hypertrophy (hyperplasia): <i>A. K. Gadre and K. C. Gadre</i></li> <li>Morphology of a columellaform stapes: <i>C. Vande Maele, P. Van Cauwenberge, P. Kluyskeris and L. Vakaet</i></li> <li>Localized middle ear effusions: Three unusual acute cases in adults: <i>D. A. Parker, A. P. Reid and P. J. Patel</i></li> <li>Carcinoid tumour of the middle ear: <i>H. Kodama, H. Takezawa, T. Suzuki, K. Kobayashi, A. Kataura, J. Wakabayashi and M. Satoh</i></li> <li>Primary cholesteatoma of the mastoid presenting as cervical abscess: <i>A. A. El-Kholy and J. R. Cherry</i></li> <li>Transpalatal approach to the petrous apex: <i>AE. Davis and P. E. Robin</i></li> <li>Traumatic CSF fistula presenting late as a middle ear effusion: <i>J. T. Brandrick</i></li> <li>Recurrent meningitis associated with meningioma of the mastoid cavity: <i>R. J. Kelleher, J. A. M. Murray and P. D. Welsby</i></li> <li>Temporal bone venous anomaly of surgical significance: <i>J. C. Shotton, H. Ludman and T. C. S. Cox</i></li> <li>Aneurysm of the anterior inferior cerebellar artery mimicking an intracanalicular acoustic neuroma: <i>M. J. Gleeson, T. C. S. Cox and A. J. Strong</i></li> <li>Mycotic aneurysm of the petrous portion of the internal carotid artery: <i>M. J. R. R.</i></li> </ul>	<ul> <li>79</li> <li>83</li> <li>86</li> <li>92</li> <li>94</li> <li>97</li> <li>99</li> <li>101</li> <li>107</li> <li>111</li> </ul>
<ul> <li>Cholesteatoma of the external auditory canal in hemifacial hypertrophy (hyperplasia): <i>A. K. Gadre and K. C. Gadre</i></li> <li>Morphology of a columellaform stapes: <i>C. Vande Maele, P. Van Cauwenberge, P. Kluyskeris and L. Vakaet</i></li> <li>Localized middle ear effusions: Three unusual acute cases in adults: <i>D. A. Parker, A. P. Reid and P. J. Patel</i></li> <li>Carcinoid tumour of the middle ear: <i>H. Kodama, H. Takezawa, T. Suzuki, K. Kobayashi, A. Kataura, J. Wakabayashi and M. Satoh</i></li> <li>Primary cholesteatoma of the mastoid presenting as cervical abscess: <i>A. A. El-Kholy and J. R. Cherry</i></li> <li>Transpalatal approach to the petrous apex: <i>AE. Davis and P. E. Robin</i></li> <li>Traumatic CSF fistula presenting late as a middle ear effusion: <i>J. T. Brandrick</i></li> <li>Recurrent meningitis associated with meningioma of the mastoid cavity: <i>R. J. Kelleher, J. A. M. Murray and P. D. Welsby</i></li> <li>Temporal bone venous anomaly of surgical significance: <i>J. C. Shotton, H. Ludman and T. C. S. Cox</i></li> <li>Aneurysm of the anterior inferior cerebellar artery mimicking an intracanalicular acoustic neuroma: <i>M. J. Gleeson, T. C. S. Cox and A. J. Strong</i></li> <li>Mycotic aneurysm of the petrous portion of the internal carotid artery: <i>J. Samuel and C. M. C. Fernandes</i></li> <li>Post-traumatic asymptomatic saccular aneurysm of the internal carotid artery: <i>M. J. R. R. Vanlierde</i></li> </ul>	79 83 86 92 94 97 99 101 107 111 115
<ul> <li>Cholesteatoma of the external auditory canal in hemifacial hypertrophy (hyperplasia): <i>A. K. Gadre and K. C. Gadre</i></li> <li>Morphology of a columellaform stapes: <i>C. Vande Maele, P. Van Cauwenberge, P. Kluyskeris and L. Vakaet</i></li> <li>Localized middle ear effusions: Three unusual acute cases in adults: <i>D. A. Parker, A. P. Reid and P. J. Patel</i></li> <li>Carcinoid tumour of the middle ear: <i>H. Kodama, H. Takezawa, T. Suzuki, K. Kobayashi, A. Kataura, J. Wakabayashi and M. Satoh</i></li> <li>Primary cholesteatoma of the mastoid presenting as cervical abscess: <i>A. A. El-Kholy and J. R. Cherry</i></li> <li>Transpalatal approach to the petrous apex: <i>AE. Davis and P. E. Robin</i></li> <li>Traumatic CSF fistula presenting late as a middle ear effusion: <i>J. T. Brandrick</i></li> <li>Recurrent meningitis associated with meningioma of the mastoid cavity: <i>R. J. Kelleher, J. A. M. Murray and P. D. Welsby</i></li> <li>Temporal bone venous anomaly of surgical significance: <i>J. C. Shotton, H. Ludman and T. C. S. Cox</i></li> <li>Aneurysm of the enterior inferior cerebellar artery mimicking an intracanalicular acoustic neuroma: <i>M. J. Gleeson, T. C. S. Cox and A. J. Strong</i></li> <li>Mycotic aneurysm of the petrous portion of the internal carotid artery: <i>J. Samuel and C. M. C. Fernandes</i></li> <li>Post-traumatic asymptomatic saccular aneurysm of the internal carotid artery: <i>M. J. R. R. Vanlierde</i></li> <li>Recurrent bilateral peripheral facial palsy: <i>N. Stahl and T. Ferit</i></li> </ul>	79 83 86 92 94 97 99 101 107 111 115 117
<ul> <li>Cholesteatoma of the external auditory canal in hemifacial hypertrophy (hyperplasia): <i>A. K. Gadre and K. C. Gadre</i></li> <li>Morphology of a columellaform stapes: <i>C. Vande Maele, P. Van Cauwenberge, P. Kluyskeris and L. Vakaet</i></li> <li>Localized middle ear effusions: Three unusual acute cases in adults: <i>D. A. Parker, A. P. Reid and P. J. Patel</i></li> <li>Carcinoid tumour of the middle ear: <i>H. Kodama, H. Takezawa, T. Suzuki, K. Kobayashi, A. Kataura, J. Wakabayashi and M. Satoh</i></li> <li>Primary cholesteatoma of the mastoid presenting as cervical abscess: <i>A. A. El-Kholy and J. R. Cherry</i></li> <li>Transpalatal approach to the petrous apex: <i>AE. Davis and P. E. Robin</i></li> <li>Traumatic CSF fistula presenting late as a middle ear effusion: <i>J. T. Brandrick</i></li> <li>Recurrent meningitis associated with meningioma of the mastoid cavity: <i>R. J. Kelleher, J. A. M. Murray and P. D. Welsby</i></li> <li>Temporal bone venous anomaly of surgical significance: <i>J. C. Shotton, H. Ludman and T. C. S. Cox</i></li> <li>Aneurysm of the anterior inferior cerebellar artery mimicking an intracanalicular acoustic neuroma: <i>M. J. Gleeson, T. C. S. Cox and A. J. Strong</i></li> <li>Mycotic aneurysm of the petrous portion of the internal carotid artery: <i>J. Samuel and C. M. C. Fernandes</i></li> <li>Post-traumatic asymptomatic saccular aneurysm of the internal carotid artery: <i>M. J. R. R. Vanlierde</i></li> <li>Recurrent bilateral peripheral facial palsy: <i>N. Stahl and T. Ferit</i></li> </ul>	79 83 86 92 94 97 99 101 107 111 115 117 121

Printed in Great Britain and published for the proprietors by Headley Brothers Ltd. The Invicta Press Ashford Kent and London