

Research Paper

Cite this article: Soares GB, Martins MF, Vianna RT, Domingues MV, Adriano EA (2023). Morphology and molecular phylogeny of *Chauhanellus* Bychowsky & Nagibina, 1969 (Monogenoidea) parasitizing marine catfish (Ariidae) from the Atlantic coast of South America: a new species, supplementary taxonomic information and new insights. *Journal of Helminthology* **97**, e32, 1–14. <https://doi.org/10.1017/S0022149X23000135>

Received: 26 September 2022

Revised: 2 February 2023

Accepted: 25 February 2023

Keywords:

Dactylogyridae; Fish parasite; Integrative taxonomy; Neotropical region

Author for correspondence:

G.B. Soares,

E-mail: geusivansoares@gmail.com

Morphology and molecular phylogeny of *Chauhanellus* Bychowsky & Nagibina, 1969 (Monogenoidea) parasitizing marine catfish (Ariidae) from the Atlantic coast of South America: a new species, supplementary taxonomic information and new insights

G.B. Soares¹ , M.F. Martins², R.T. Vianna² , M.V. Domingues³ 
and E.A. Adriano^{1,4} 

¹Programa de Pós-graduação em Biologia Animal, Departamento de Biologia Animal, Instituto de Biologia, Universidade Estadual de Campinas (UNICAMP), Rua Monteiro Lobato, 255, CEP 13083-862 Campinas, São Paulo, Brazil; ²Programa de Pós-graduação em Biologia de Ambientes Aquáticos Continentais, Instituto de Ciências Biológicas, Universidade Federal do Rio Grande (FURG), Avenida Itália km 8, CEP 96203-900, Rio Grande, Rio Grande do Sul, Brazil; ³Programa de Pós-graduação em Biologia Ambiental, Instituto de Estudos Costeiros, Universidade Federal do Pará (UFPA), Travessa Leandro Ribeiro, s/n, Aldeia, CEP 68600-000 Bragança, Pará, Brazil and ⁴Departamento de Ecologia e Biologia Evolutiva, Universidade Federal de São Paulo (UNIFESP), Rua Professor Arthur Riedel, 275, Jardim Eldorado, CEP 09972-270 Diadema, São Paulo, Brazil

Abstract

In this study, a new dactylogyrid species is described from the gills of two ariid fish species of the Brazilian coast, *Genidens barbatus* and *Genidens genidens*, by combining morphological characters with partial 18S rDNA sequences. The new species can be distinguished from its congeners by the following morphological characteristics: a male copulatory organ (MCO), a coiled tubular shaft of around one counterclockwise ring, base with a sclerotized cap; MCO with tapered distal region; a T-shaped accessory piece; a vagina, a cup-shaped vaginal vestibule, slightly sclerotized; and a pharynx comprising a muscular, glandular bulb. Supplementary taxonomic data for *Chauhanellus velum* from *Sciades couma* (type-host) are also presented and new 18S rDNA sequences of *Chauhanellus* spp. from other ariid fish from South America are provided. Phylogenetic analyses based on partial 18S rDNA gene sequences placed *Chauhanellus riograndinensis* n. sp. and *Chauhanellus velum* as two early divergent lineages within *Chauhanellus* from South America. Finally, a way to test the monophyly of *Chauhanellus* and *Hamatopeduncularia* is also discussed, which may be useful for future studies.

Introduction

Ariidae (Siluriformes) species, also known as marine catfish, are widespread in the tropical and subtropical continental shelves of the Atlantic, Indian and Pacific Oceans (Marceniuk *et al.*, 2012). Currently, a total of 156 Ariidae species belonging to 34 genera are known and ~20 species have been found to occur on the Atlantic coast of South America (Marceniuk *et al.*, 2012; Fricke *et al.*, 2022).

There are 77 species of Monogenoidea currently parasitizing Ariidae around the world; 68 Dactylogyridae (*Chauhanellus* Bychowsky & Nagibina, 1969, *Hamatopeduncularia* Yamaguti, 1953; *Neotetraonchus* Bravo-Hollis, 1968 and *Susanlimocotyle* Soares, Domingues & Adriano, 2021), seven Neocalceostomatidae (*Fridericianella* Brandes, 1894, *Neocalceostoma* Tripathi, 1959, *Neocalceostomoides* Kritsky, Mizelle & Bilqees, 1978 and *Thysanotophaptor* Kritsky, Shameem, Kumari, & Krishnaveni, 2012), and two Udonellidae (*Udonella* Johnson, 1835) (Lim *et al.*, 2001; Domingues *et al.*, 2016; Illa *et al.*, 2019; Soares *et al.*, 2021a, b; Soo & Tan, 2021).

Chauhanellus, as amended by Lim (1994), is the second most species-rich genus, with 30 valid species, behind only *Hamatopeduncularia*, which comprises 32 species (Lim *et al.*, 2001; Soares *et al.*, 2021a; Soo & Tan, 2021). *Chauhanellus* species possess a haptor with or usually without a digit-like extension; a dorsal anchor with or without spines at proximal base surface and slightly expanded outer roots; a ventral anchors with outer roots usually expanded and base of inner roots thickened; and a vaginal dextral opening with a sclerotized vaginal tube. Of the *Chauhanellus* species known to date, 28 have been described as infecting ariid marine

Table 1. *Chauhanellus* species from ariid hosts.

<i>Chauhanellus</i> species	Type host	Type locality	References
<i>C. alatus</i> (Chauhan, 1945)	<i>Arius arius</i> ^a	India	Lim et al. (2001)
<i>C. aspinosus</i> Lim, 1994	<i>Arius venosus</i>	Malaysia	Lim et al. (2001)
<i>C. auriculatum</i> Lim, 1994	<i>Arius maculatus</i>	Malaysia	Lim et al. (2001)
<i>C. australis</i> (Young, 1967)	<i>Neoarius australis</i>	Australia	Lim et al. (2001)
<i>C. boegeri</i> Domingues & Fehlaue, 2006	<i>Genidens barbatus</i>	Brazil	Domingues & Fehlaue (2006)
<i>C. caelatus</i> Lim, 1994	<i>Nemapteryx caelata</i> ^b	Malaysia	Lim et al. (2001)
<i>C. chauhani</i> Venkatanarasaiah & Kulkarni, 1990	<i>Plicofollis dussumieri</i> ^c	India	Lim et al. (2001)
<i>C. digitalis</i> Lim, 1994	<i>Hexanematchichthys sagor</i> ^d	Malaysia	Lim et al. (2001)
<i>C. duriensis</i> Lim, 1994	<i>Netuma thalassina</i> ^e	Malaysia	Lim et al. (2001)
<i>C. flexiosus</i> Bychowsky & Nagibina, 1969	<i>A. arius</i> ^a	China	Lim et al. (2001)
<i>C. forcipi</i> Lim, 1994	<i>Hexanematchichthys sagor</i> ^d	Malaysia	Lim et al. (2001)
<i>C. hamatopeduncularoideum</i> Domingues, Soares & Watanabe, 2016	<i>Amphiarus rugispinis</i>	Brazil	Domingues et al. (2016)
<i>C. hypenocleithrum</i> Domingues, Soares & Watanabe, 2016	<i>Sciades proops</i>	Brazil	Domingues et al. (2016)
<i>C. indicus</i> Rastogi, Kumar & Singh, 2004	<i>Wallago attu</i> ^f	India	Rastogi et al. (2004)
<i>C. intermedius</i> Lim, 1994	<i>Hexanematchichthys sagor</i> ^d	Malaysia	Lim et al. (2001)
<i>C. malayanus</i> Lim, 1994	<i>H. sagor</i> ^d	Malaysia	Lim et al. (2001)
<i>C. nagibinae</i> Paperna, 1977	<i>Plicofollis argyropleuron</i> ^g	Kenya	Lim et al. (2001)
<i>C. nengi</i> (Tripathi, 1959)	<i>Nemapteryx nenga</i> ^h	India	Lim et al. (2001)
<i>C. neotropalis</i> Domingues & Fehlaue, 2006	<i>Aspistor luniscutis</i>	Brazil	Domingues & Fehlaue (2006)
<i>C. oculatus</i> Bychowsky & Nagibina, 1969	<i>A. maculatus</i>	China	Lim et al. (2001)
<i>C. osteogeneiosus</i> Lim, 1994	<i>Osteogeneiosus militaris</i>	Malaysia	Lim et al. (2001)
<i>C. pedunculatus</i> Paperna, 1977	<i>Arius</i> sp.	India	Lim et al. (2001)
<i>C. poculus</i> Lim, 1994	<i>A. maculatus</i>	Malaysia	Lim et al. (2001)
<i>C. pulutanus</i> Lim, 1994	<i>A. maculatus</i>	Malaysia	Lim et al. (2001)
<i>C. riograndinensis</i> n. sp.	<i>Genidens barbatus</i>	Brazil	Present study
<i>C. seenghali</i> (Kumar, 2013)	<i>Sperata seenghali</i> ⁱ	India	Domingues et al. (2016)
<i>C. susamlimae</i> Domingues, Soares & Watanabe, 2016	<i>Sciades herzbergii</i>	Brazil	Domingues et al. (2016)
<i>C. trifidus</i> Lim, 1994	<i>Hexanematchichthys sagor</i> ^d	Malaysia	Lim et al. (2001)
<i>C. tuberhamatus</i> Zhang & Ding, 1997	<i>Tachysurus sinensis</i> ^j	China	Lim et al. (2001)
<i>C. velum</i> Domingues, Soares & Watanabe, 2016	<i>Sciades couma</i>	Brazil	Domingues et al. (2016)
<i>C. youngi</i> Kearns & Whittington, 1994	<i>Neoarius graeffei</i> ^k	Australia	Lim et al. (2001)

The new species described in this study is in boldface type.

^aFormerly *Arius falcarius*

^bFormerly *Arius caelatus*

^cFormerly *Tachysurus dussumieri*

^dFormerly *Arius sagor*

^eFormerly *Arius thalassinus*

^fSiluridae

^gFormerly *Arius macrocephalus*

^hFormerly *Arius nenga*

ⁱBagridae

^jFormerly *Arius sinensis*

^kFormerly *Arius graeffei*

catfish worldwide, while only two species have also been reported in non-ariid hosts (see table 1) (Lim et al., 2001; Domingues & Fehlaue, 2006; Domingues et al., 2016).

In South America, only six species of *Chauhanellus* have been described in Atlantic coast ariids (table 1) (Domingues & Fehlaue, 2006; Domingues et al., 2016; Soares et al., 2021a).

However, recent studies indicate that the actual number of species in the region is probably much higher (Soares et al., 2022).

During a study of monogenoids of ariid species captured from the Brazilian coast, a new species of *Chauhanellus* was found in the gills of *Genidens barbatus* (Lacepède) and *Genidens genidens* (Cuvier) and is described based on morphological characters

and partial 18S *rDNA* sequences. Supplementary taxonomic data from *Chauhanellus velum* Domingues, Soares & Watanabe, 2016, a parasite of *Sciades couma* (Valenciennes) (type-host) and new partial 18S *rDNA* sequences of *Chauhanellus* spp. from South American ariids, are also presented. Finally, new insights into *Chauhanellus* are provided based on morphology and molecular phylogenetic evidence.

Material and methods

Sample collection, morphological study and deposit of the helminths

Specimens of distinct ariid species were collected by local fishermen with trammel nets and hooks from four locations on the Brazilian coast between December 2017 to December 2019 under a License for the Collection of Zoological Material (SISBio No. 60666-2 and Sisgen No. AD28DC2) (table 2). The gill arches were removed and placed in vials containing heated water (~65°C), following which each vial was vigorously shaken. The contents of each vial were examined in the laboratory under a dissecting microscope and helminths were removed from the gills or sediment using small probes. Helminths were fixed in either 4% formalin for morphological study or 96% ethanol for molecular characterization. Some specimens were stained with Gomori's trichrome and mounted in Damar gum to examine their internal soft structures, while others were mounted in Hoyer's medium (Humason, 1979; Boeger & Vianna, 2006) for the study of the sclerotized structures. Measurements, all in micrometres, were taken following the procedures of Mizelle & Klucka (1953). Dimensions of organs and other structures represent the highest measurements in the dorso-ventral view; lengths of curved or bent structures (bars and accessory piece) represent the straight-line distances between the extreme ends; anchor length measurements followed Soares *et al.* (2019); and the total length of the male copulatory organ (MCO) was measured using ImageJ (Rasband, 2022). Measurements are presented in micrometres as the mean followed by the range and the number (*n*) of specimens measured is shown in parentheses. Illustrations were prepared with a drawing tube attached to a Leica DM 2500 microscope with differential interference contrast

and phase contrast optics. The soft structures were illustrated using pen and ink, while the hard structures were scanned and redrawn on a digitizing tablet using CorelDraw (2014). Plates were also prepared in CorelDraw (2014). Definitions of prevalence, mean intensity and mean abundance followed Bush *et al.* (1997). The Bray–Curtis similarity test (Bray & Curtis, 1957) was used to evaluate possible morphometric similarities between the specimens of *Chauhanellus* from *S. couma* (present study) and *C. velum* from other previously described hosts. Type specimens, vouchers and hologenophores (Pleijel *et al.*, 2008) were deposited in the Invertebrate Collection of the Museu Paraense Emílio Goeldi (MPEG. PLA), Belém, Pará, Brazil, and the collection of Platyhelminthes of the Adão José Cardoso Museum of Zoology of the State University of Campinas (ZUEC PLA), São Paulo, Brazil, respectively, under No. (MPEG.PLA 000359–000384; ZUEC PLA 186–187). The vouchers of *C. velum* CHIOC 38262 a–b, 38263 were examined for comparative purposes. Nomenclature for hosts followed Marceniuk *et al.* (2012). Details of the new taxa have been submitted to ZooBank following the International Code of Zoological Nomenclature (article 8.5 of the amended version) (International Commission on Zoological Nomenclature, 2012).

Molecular characterization of parasites

Each monogenoid specimen subjected to molecular analysis was divided using fine needles under a dissecting microscope. The anterior half of the body (without the MCO) was placed in a 1.5 ml microtube with 96% ethanol for genomic DNA extraction. The posterior part containing the haptor complex and the MCO were flattened under coverslip pressure and mounted in Hoyer's solution for species identification. These fragments also served as vouchers (hologenophores). Genomic DNA was extracted using a Qiagen Dneasy® Blood and Tissue Kit, according to the manufacturer's protocol, with a final volume of 30 µl. DNA concentration was verified using a NanoDrop 2000 spectrophotometer (Thermo Fisher Scientific, Massachusetts, USA) at 260 nm.

The partial 18S *rDNA* was amplified using a two-round polymerase chain reaction (PCR). In the first round, DNA was amplified with the primer pair WormA and WormB (Littlewood & Olson, 2001). In the second round, for the semi-nested PCRs,

Table 2. Host species, locality (geographical coordinates), and monogenoid species detected in the present effort on each fish species.

Host	N	Locality	Parasite
<i>Amphiarus rugispinis</i>	10	Ajuruteua (0°49'31" N;46°36'29" W), Bragança, PA, Br	<i>Chauhanellus hamatopeduncularoideum</i> ^a <i>Chauhanellus neotropicalis</i>
<i>Aspistor quadriscutis</i>	6	Ajuruteua (0°49'31" N; 46°36'29" W), Bragança, PA, Br	<i>C. neotropicalis</i> ^a
	8	Caratateua (1° 59' 41.91" S; 46° 43' 21.385" W), Bragança, PA, Br	<i>C. neotropicalis</i>
<i>Genidens barbatus</i>	18	Cananéia (25°02'09.2" S; 47°54'57.8" W), SP, Br	<i>Chauhanellus boegeri</i>
	31	Estuary of Patos Lagoon (32° 08' 05.7" S; 52° 06' 11.2" W), RS, Br	<i>C. boegeri</i> <i>Chauhanellus riograndinensis</i> n. sp. ^a
<i>Genidens genidens</i>	17	Estuary of Patos Lagoon (32° 08' 05.7" S; 52° 06' 11.2" W), RS, Br	<i>C. boegeri</i> <i>Chauhanellus riograndinensis</i> n. sp
<i>Sciades couma</i>	6	Caratateua (1° 59' 41.91" S; 46° 43' 21.385" W), Bragança, PA, Br	<i>Chauhanellus hamatopeduncularoideum</i> <i>Chauhanellus velum</i> ^a <i>C. boegeri</i>
<i>Sciades proops</i>	9	Ajuruteua (0°49'31" N;46°36'29" W), Bragança, PA, Br	<i>Chauhanellus hypenocleithrum</i> ^a

N, number of host; PA, Pará; SP, São Paulo; RS, Rio Grande do Sul; BR, Brazil.

^aSpecimen subjected to molecular analysis, for which sequences of 18S *rDNA* were used for the phylogenetic analysis.

the primer combinations were WormA and 1270R (Littlewood & Olson, 2001) and 930F (Littlewood *et al.*, 2008) with WormB, which amplified two overlapping fragments of approximately 1179 base pairs (bp) and 1054 bp, respectively. PCRs were performed in a Mathercycler[®] nexus (Eppendorff, Hamburg, Germany) with a final volume of 25 μ l: 12.5 μ l of DreamTaq Green PCR Master Mix (2 \times) (Thermo Scientific Wilmington, USA), following the manufacturer's recommendations, 0.5 mM of each primer, and 3 μ l of the extracted DNA.

The PCR profile was performed using the cycling described in Soares *et al.* (2021a). The semi-nested PCRs were conducted with 1 μ l of the product of the PCRs, diluted 1:1 in ultrapure water, applying the same cycling conditions. Amplicons were electrophoresed in 2% agarose gel in a TAE buffer (Tris 40 mM, acetic acid 20 mM, EDTA 1 mM) stained with SYBRsafe[®] (Invitrogen, Thermo Fisher Scientific, Massachusetts, USA) alongside a 1 kb Plus DNA Ladder (Invitrogen, Thermo Fisher Scientific, Massachusetts, USA) at 100 V for 30 min. The PCR products were purified using a QIAquick PCR Purification Kit (Qiagen, USA) and sequencing was carried out with the BigDye[®] Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems[™]) in a 3500 DNA sequencing analyser (Applied Biosystems, California, USA) at Helixxa Company (Paulínia, São Paulo, Brazil) or at the Human Genome Research Center, of the University of São Paulo (São Paulo, Brazil), using the same primers as used for rDNA amplification.

Alignment and phylogenetic inference

Contigs were edited using Sequencher 4.1.4 (Gene Codes, Ann Arbor, MI) and deposited in GenBank under the accession numbers listed in table 3. Standard nucleotide Basic Local Alignment Search Tool searches were then conducted (Altschul *et al.*, 1997) to verify the similarity of the newly obtained sequences from the present study with other sequences of monogenoids in the United States National Center for Biotechnology Information (NCBI) BioSystems database (Geer *et al.*, 2009). Alignment of the 18S rDNA was generated using MUSCLE implemented in Geneious version 7.1.3 (Kearse *et al.*, 2012). A total of 34 partial sequences of the 18S rDNA of species belonging to the Dactylogyridea order published in the NCBI BioSystems database (Geer *et al.*, 2009), along with two of the Monocotylidae order (used as the outgroup) were retrieved from GenBank (see table 3) and aligned with five newly generated sequences of *Chauhanellus* spp. from ariids from the Brazilian coast (*Chauhanellus riograndinensis* n. sp., *Chauhanellus neotropicalis* Domingues & Fehlaue, 2006, *Chauhanellus hamatopeduncularoideum* Domingues, Soares & Watanabe, 2016, *Chauhanellus hypenocleithrum* Domingues, Soares & Watanabe, 2016, and *C. velum*). Despite the availability of several partial 18S rDNA sequences, we restricted our analysis to sequences of monogenoids >1600 bp to attain the highest number of variable and phylogenetically informative sites. Forty-one sequences (1647–2200 bp long) were aligned, and the extremes were trimmed, leaving an alignment of 1777 bp in length. The model of evolution was selected by JModelTest 2.1.1 (Darrriba *et al.*, 2012) using the Akaike information criterion. Phylogenetic analyses were performed using the maximum likelihood (ML) and Bayesian inference (BI) methods. ML was performed in PhyML 3.0, implemented via a web server (<http://www.atgc-montpellier.fr/phyml/>) (Guindon *et al.*, 2010), with topology assessed by bootstrapping with 1000 replicates, applying the GTR + I + G model. BI was performed using MrBayes v.3.2.6

(Ronquist *et al.*, 2012), under the same model, with posterior probabilities estimated from 5×10^5 generations with two independent runs of four simultaneous Markov chain Monte Carlo (MCMC) algorithms, sufficient to keep the average standard deviation below 0.001 and the effective sample size (>200) on Tracer v1.7 (Rambaut *et al.*, 2018). The MCMC algorithms with the 1000th tree saved and diagnostics for every 1000th generation with burn-in periods, were set to the first 25,000 generations. Trees were visualized using FigTree 1.3.1 (Rambaut, 2022) and figures prepared using CorelDraw (2014). Genetic divergence was determined using the *P*-distance model matrix in MEGA version 7 (Kumar *et al.*, 2016). Gaps and missing data were deleted.

Results

A total of 100% of the 10 host specimens of *Amphiarus rugispinis* (Valenciennes), 100% of the fourteen *Aspistor quadriscutis* (Valenciennes), five (16%) of the 31 host specimens of *G. barbatus*, seven (42%) of the seventeen *G. genidens*, three (50%) of the six *Sciades couma* and 100% of the nine *S. proops* (Valenciennes) examined were infected with monogenoids (table 2).

The morphological, morphometric and partial 18S rDNA data endorsing the proposition of the new taxon, the supplementary taxonomic of *C. velum* and insights into *Chauhanellus* are presented below.

Taxonomic acts

Taxonomic summary

Class Monogenoidea Bychowsky, 1937

Subclass Polyonchoinea Bychowsky, 1937

Order Dactylogyridea Bychowsky, 1937

Dactylogyridae Bychowsky, 1933

Chauhanellus Bychowsky & Nagibina, 1969

Chauhanellus riograndinensis n. sp. (fig. 1)

Synonym. *Chauhanellus* sp. (Soares *et al.*, 2022)

Type-host. *Genidens barbatus* (Lacepède), (Siluriformes, Ariidae).

Site of infection. Gills.

Type locality. Estuary of Patos Lagoon, Municipality of Rio Grande, Rio Grande do Sul State, Brazil (32° 08' 05.7" S; 52° 06' 11.2" W).

Other records. *Genidens genidens* (Cuvier), Ariidae (prevalence: 42% of 17 hosts; mean intensity: 47; mean abundance: 1.1), Estuary of Patos Lagoon, Municipality of Rio Grande, Rio Grande do Sul State, Brazil (32° 08' 05.7" S; 52° 06' 11.2" W).

Prevalence. 16% of 31 hosts examined.

Mean intensity. 32 parasites per infected host.

Mean abundance. two parasites per host.

Specimens deposited. Holotype, MPEG.PLA 000359; paratypes, MPEG.PLA 000360-000372; vouchers, MPEG.PLA 000373-000379; hologenophore, ZUEC PLA 187.

Representative DNA sequence. 1689 bp long partial sequence of the 18S rDNA gene of one parasite isolates (GenBank accession number, OQ517175).

Etymology: The specific name refers to the municipality of Rio Grande, Rio Grande do Sul State, Brazil, where the type host was collected.

Number of ZooBank. 7E70DF71-791A-4333-B6B7-7989BB6F4EA1

Comparative measurements. table 4

Description. (Based on fourteen specimens, eight mounted in Hoyer's medium and six stained with Gomori's trichrome). Body fusiform, total length excluding haptor 863 (743–972;

Table 3. List of monogenoids included in the phylogenetic analyses, providing host species data, locality, GenBank ID and references. Data for the present study are highlighted with boldface type.

Parasites species	Host	Host family	Locality	GenBank ID	Reference
Dactylogyridae					
Ancyrocephalinae					
<i>Anacanthorus penilabiatus</i>	<i>Piaractus mesopotamicus</i>	Serrasalminidae	Brazil	KU941837	Muller <i>et al.</i> (2016)
<i>Bravohollisia tecta</i>	<i>Pomadasys maculatus</i>	Haemulidae	China	KJ571020	Sun <i>et al.</i> (2014)
<i>Bravohollisia maculatus</i>	<i>P. maculatus</i>	Haemulidae	China	KJ571018	Sun <i>et al.</i> (2014)
<i>Euryhaliotrema johnii</i>	<i>Lutjanus johnii</i>	Lutjanidae	China	EU836214	Sun <i>et al.</i> (2014)
<i>Euryhaliotrematoides annulocirrus</i> ^a	<i>Chaetodon vagabundus</i>	Chaetodontidae	Australia	AY820602	Plaisance <i>et al.</i> (2005)
<i>Euryhaliotrematoides berenguelae</i> ^a	<i>Chaetodon citrinellus</i>	Chaetodontidae	French Polynesia	AY820604	Plaisance <i>et al.</i> (2005)
<i>Euryhaliotrematoides triangulovagina</i> ^a	<i>Chaetodon kleinii</i>	Chaetodontidae	Palau	AY820608	Plaisance <i>et al.</i> (2005)
<i>Euryhaliotrematoides pirulum</i> ^a	<i>Chaetodon lunula</i>	Chaetodontidae	French Polynesia	AY820607	Plaisance <i>et al.</i> (2005)
<i>Haliotrema aurigae</i>	<i>Chaetodon auriga</i>	Chaetodontidae	Australia	AY820610	Plaisance <i>et al.</i> (2005)
<i>Haliotrema scyphovagina</i>	<i>Forcipiger flavissimus</i>	Chaetodontidae	French Polynesia	AY820611	Plaisance <i>et al.</i> (2005)
<i>Lethrinitrema zhanjiangense</i>	<i>Lethrinus nebulosus</i>	Lethrinidae	China	KJ571021	Sun <i>et al.</i> (2014)
<i>Lethrinitrema grossecurvittubu</i>	<i>L. nebulosus</i>	Lethrinidae	China	EU836225	Sun <i>et al.</i> (2014)
<i>Mymarothecium viatorum</i>	<i>Piaractus mesopotamicus</i>	Serrasalminidae	Brazil	KU941838	Muller <i>et al.</i> (2016)
<i>Pseudohaliotrema sphincterporus</i>	<i>Siganus doliatus</i>	Siganidae	Australia	AJ287568	Littlewood & Olson (2001)
Ancylo-discoidinae					
<i>Bychowskyella fossilisi</i>	<i>Heteropneustes fossilisi</i>	Heteropneustidae	India	KT852454	Verma <i>et al.</i> (2017a)
<i>Bychowskyella tchangii</i>	<i>Clarias batrachus</i>	Clariidae	India	KT852455	Verma <i>et al.</i> (2017a)
<i>Chauhanellus boegeri</i>	<i>Sciades herzbergii</i>	Ariidae	Brazil	MW132134	Soares <i>et al.</i> (2021a)
<i>Chauhanellus hamatopeduncularoideum</i>	<i>Amphiarus rugispinis</i>	Ariidae	Brazil	OQ511493	Present study
<i>Chauhanellus hypenocleithrum</i>	<i>Sciades proops</i>	Ariidae	Brazil	OQ511507	Present study
<i>Chauhanellus neotropicalis</i>	<i>Aspistor quadriscutis</i>	Ariidae	Brazil	OQ511509	Present study
<i>Chauhanellus riograndinensis</i> n. sp	<i>Genidens barbatus</i>	Ariidae	Brazil	OQ517175	Present study
<i>Chauhanellus susamlimae</i>	<i>Sciades herzbergii</i>	Ariidae	Brazil	MW144439	Soares <i>et al.</i> (2021a)
<i>Chauhanellus velum</i>	<i>S. herzbergii</i>	Ariidae	Brazil	MW144823	Soares <i>et al.</i> (2021a)
<i>C. velum</i>	<i>Sciades couma</i>	Ariidae	Brazil	OQ511558	Present study
<i>Hamatopeduncularia arii</i>	<i>Arius jella</i>	Ariidae	India	KT252895	Illa <i>et al.</i> (2019)
<i>Hamatopeduncularia bifida</i>	<i>A. jella</i>	Ariidae	India	MK084781	Illa <i>et al.</i> (2019)
<i>Hamatopeduncularia elongata</i>	<i>A. jella</i>	Ariidae	India	MK084780	Illa <i>et al.</i> (2019)
<i>Hamatopeduncularia madhaviae</i>	<i>Plicofollis dussumieri</i>	Ariidae	India	KT252898	Illa <i>et al.</i> (2019)
<i>Hamatopeduncularia thalassini</i>	<i>A. jella</i>	Ariidae	India	KT252900	Illa <i>et al.</i> (2019)
<i>Mizelleus longicirrus</i>	<i>Wallago attu</i>	Siluridae	India	KR296801	Verma <i>et al.</i> (2017b)
<i>Susanlimocotyle narina</i>	<i>S. herzbergii</i>	Ariidae	Brazil	MW144824	Soares <i>et al.</i> (2021a)
<i>Thaparocleidus gangus</i>	<i>W. attu</i>	Siluridae	India	KX364088	Verma <i>et al.</i> (2017a)

(Continued)

Table 3. (Continued.)

Parasites species	Host	Host family	Locality	GenBank ID	Reference
<i>Thaparocleidus wallagonius</i>	<i>W. attu</i>	Siluridae	India	KX364085	Verma et al. (2017a)
Diplectanidae					
<i>Lamellogadus japonicus</i>	<i>Acanthopagrus s. schlegelii</i>	Sparidae	China	EU836236	Sun et al. (2014)
<i>Lamellogadus pagrosomi</i>	<i>Pagrus major</i>	Sparidae	China	EU836235	Sun et al. (2014)
<i>Pseudorhabdosynchus grouperi</i>	<i>Epinephelus coioides</i>	Serranidae	Indonesia	FJ655782	Wu et al. (2005)
Pseudomurraytrematidae					
<i>Pseudomurraytrema ardens</i>	<i>Catostomus ardens</i>	Catostomidae	United States	AJ228793	Littlewood et al. (1998)
Anoplodiscidae					
<i>Anoplodiscus cirruspiralis</i>	<i>Sparus auratus</i>	Sparidae	Australia	AJ287475	Littlewood & Olson (2001)
Sundanonchidae					
<i>Sundanonchus micropeltis</i>	<i>Channa micropeltis</i>	Channidae	Malaysia	AJ287579	Littlewood & Olson (2001)
Monocotylidae					
<i>Calicotyle affinis</i> ^b	<i>Chimaera monstrosa</i>	Chimaeridae	Norway	AJ228777	Littlewood et al. (1999)
<i>Dictyocotyle coeliaca</i> ^b	<i>Amblyraja radiata</i>	Rajidae	United Kingdom	AJ228778	Littlewood et al. (1998)

^a*Euryhalotrematoides* was placed in subjective synonymy with *Euryhalotrema* (Kritsky, 2012).

^bSpecies used as outgroups.

$n = 6$), total width at the level of germarium 141 (126–168; $n = 6$) (fig. 1a). Cephalic margin tapered; poorly developed terminal lobes; four bilateral pairs of head organs with rod-shaped secretion; cephalic glands unicellular, lateral to pharynx. Eyes four, equidistant; accessory chromatic granules absent. Mouth subterminal, midventral, prepharyngeal; pharynx comprising muscular, glandular bulb, spherical, 72 (63–85; $n = 6$) long, 71 (56–88; $n = 6$) wide. Oesophagus elongate; two intestinal caeca, non-confluent posteriorly, lacking diverticula. Common genital pore midventral near level of intestinal bifurcation; genital atrium muscular, unarmed. Gonads intercaecal, testis post-germarial, dorsal to germarium. Testis bacilliform, 120 (109–134; $n = 6$) long, 52 (39–65; $n = 6$) wide. Vas deferens looping left of intestinal caecum; seminal vesicle a dilatation of vas deferens, sigmoid. Single saculiform prostatic reservoir, lying posterior to the base of MCO. Copulatory complex comprising MCO, accessory piece; MCO, sclerotized, coiled tubular shaft of around one counterclockwise ring, base with a sclerotized cap, 85 (73–102; $n = 8$) long, tapered distal region. Accessory piece sclerotized, T-shaped, proximally spatulate, guarding termination of MCO; non-articulated with MCO (fig. 1c). Germarium piriform, 108 (79–164; $n = 6$) long, 69 (48–92; $n = 6$) wide. Uterus delicate. Vagina comprises a vaginal aperture with dextroventral opening, submarginal; vaginal vestibule cup-shaped, slightly sclerotized; narrow vaginal canal, a loop at the distal portion before entering the seminal receptacle (fig. 1d). Seminal receptacle ovoid. Mehlis' glands, bilateral to the uterus. Eggs, ootype not observed. Vitellaria coextensive with intestinal caeca; transverse vitelline duct anterior to seminal receptacle, dorsal to intestinal caeca. Haptor digitiform, 242 (213–273; $n = 4$) long, 222 (180–242; $n = 4$) wide, with four haptor glands (two ventral and two mid-dorsal). Anchors dissimilar.

Ventral anchor, outer 48 (44–55; $n = 8$) long, inner 31 (27–35; $n = 8$) long; base 34 (29–38; $n = 8$) long; with divergent roots; truncate superficial root; expanded deep root, evenly curved elongate shaft, slightly straight short point extending at the level of tip of the superficial root (fig. 1g). Dorsal anchor, outer 46 (41–57; $n = 8$) long, inner 42 (38–51; $n = 8$) long; base 25 (23–27; $n = 8$) long; robust, with inconspicuous roots, slightly curved short shaft, elongate point extending well past the level of the tip of the inner base; anchor spine blunt (fig. 1h). Ventral bar, 114 (93–144; $n = 3$) long, 27 (25–29; $n = 3$) wide, broadly open U-shaped rod with bifid ends for articulation with ventral anchor (fig. 1e). Dorsal bar, 105 (86–141; $n = 5$) long, 21 (18–24; $n = 6$) wide, V-shaped, with bifurcation on both ends, acute anterior protuberance, rounded posterior protuberance; elongated posteromedial process (fig. 1f). Hooks similar in shape, 17 (15–19; $n = 16$) long, shank elongated, without inflation, erect thumb, lightly curved long shaft and delicate point (fig. 1b). Filamentous hook loop comprising 80% of the length of the shank.

Remarks

The new species seems to be closely related to *Chauhanellus susamlimae* Domingues, Soares & Watanabe, 2016, by having a digitate haptor; a ventral anchor with divergent roots, a truncate superficial root and an expanded deep root; evenly curved shaft; a dorsal anchor, robust, with inconspicuous roots; slightly curved shaft; elongated point; and anchor spine blunt. However, *C. rio-grandinensis* n. sp. can be easily distinguished from this species because the new species possesses MCO, a coiled tubular shaft of around one counterclockwise ring, base with a sclerotized cap, MCO with tapered distal region and accessory piece,




Fig. 1. *Chauhanellus riograndinensis* n. sp. (a) Holotype whole-mount, ventral; (b) hook; (c) copulatory complex; (d) vagina; (e) ventral bar; (f) dorsal bar; (g) ventral anchor; (h) dorsal anchor. Scale bars: (a) 100 µm; (b) 10 µm; (c–h) 25 µm.

T-shaped, proximally spatulate, guarding termination of MCO (MCO, a sclerotized tube, sigmoid, and accessory piece comprising an elongated sheath in *C. susamlimae*) and vagina sclerotized (a muscular vagina [unsclerotized] in *C. susamlimae*).

Chauhanellus velum Domingues, Soares & Watanabe, 2016 (fig. 2)

Type-host. *Sciades couma* (Valenciennes) (Siluriformes, Ariidae).

Site of infection. Internal borders of the primary lamellae of the gills.

Type-locality. Fish market, Municipality of Bragança, State of Pará, Brazil.

Other records. *Sciades couma*, Caratateua, Municipality of Bragança, State of Pará, Brazil (1° 59' 41.91" S, 46° 43' 21.385"

W) (present study); *Sciades herzbergii*, Caratateua, Municipality of Bragança, State of Pará, Brazil (Soares *et al.*, 2021a); *Sciades herzbergii*, Furo da Ostra, municipality of Curuçá, State of Pará, Brazil; *Sciades passany*, Fish market, Municipality of Bragança, State of Pará, Brazil (Domingues *et al.*, 2016).

Prevalence. 50% of six hosts examined.

Specimens deposited. Vouchers five (MPEG.PLA 000380-000384).

Specimens studied. Vouchers of *C. velum* CHIOC 38262 a–b, 38263.

Representative DNA sequence. 1654 bp long partial sequence of the 18S rDNA gene of one parasite isolates (GenBank accession number, OQ511558).

Comparative measurements. table 5

Table 4. Comparative measurements (in μm) of specimens of *Chauhanellus riograndinensis* n. sp. from *Genidens barbatus* and *Genidens genidens* from Rio Grande, Rio Grande do Sul, Brazil.

	<i>Genidens barbatus</i>	<i>N</i>	<i>Genidens genidens</i>	<i>N</i>
Male copulatory organ				
Length	85(73–102)	8	79(75–92)	6
Ring	17(16–18)	5	18(17–19)	3
Ventral bar				
Length	114(93–144)	3	109(105–114)	2
Width	27(25–29)	3	15(13–18)	3
Dorsal bar				
Length	105(86–141)	5	77	1
Width	21(18–24)	6	19(16–22)	3
Ventral anchor				
Outer	48(44–55)	8	45(38–52)	7
Inner	31(27–35)	8	27(22–33)	7
Base	34(29–38)	8	31(26–36)	7
Dorsal anchor				
Outer	46(41–57)	8	43(33–53)	8
Inner	42(38–51)	8	38(30–47)	8
Base	25(23–27)	8	20(15–25)	8
Hook length	17(15–19)	16	17(16–19)	16

N, number of parasite.

Redescription. (Based on five specimens, two mounted in Hoyer's medium and three stained with Gomori's trichrome). Body fusiform, total length excluding haptor 898 (766–1143; $n = 3$), total width at the level of germarium 178 (148–232; $n = 3$) (fig. 2a). Cephalic margin tapered; poorly developed terminal lobes; four bilateral pairs of head organs with rod-shaped secretion; cephalic glands not observed. Eyes four, equidistant; accessory chromatic granules absent. Mouth subterminal, midventral; pharynx subspherical, 82 (70–100; $n = 3$) long, 69 (57–79; $n = 3$) wide. Oesophagus short; two intestinal caeca, non-confluent posteriorly, lacking diverticula. Common genital pore midventral, anterior to the copulatory complex near the pharynx; genital atrium muscular, unarmed. Gonads intercaecal, testis post-germarial, dorsal to germarium. Testis subspherical, 83 (72–93; $n = 2$) long, 72 (70–75; $n = 2$) wide. Vas deferens looping left of intestinal caecum; seminal vesicle a dilatation of vas deferens, sigmoid. Single spherical prostatic reservoir, lying posterior to the base of MCO. Copulatory complex comprising MCO, accessory piece; MCO sclerotized, coiled tubular shaft of around one counterclockwise ring, base with an expanded sclerotized cap, 103 (85–113; $n = 4$) long, MCO acute distal region. Accessory piece sclerotized, comprising two regions, proximal region with three small projections (one projection serrated on the inner margin) and duct throughout the MCO pass, distal region an elongated sheath; non-articulated with MCO (fig. 2b). Germarium piriform, 121 (115–130; $n = 3$) long, 64 (50–81; $n = 3$) wide. Uterus delicate. Vagina comprises vaginal aperture with dextroventral opening, submarginal; vaginal vestibule with soft tissue; vaginal canal, short, sclerotized, knob shape, before entering the seminal receptacle. Seminal receptacle ovoid. Mehlis' glands,

bilateral to the uterus. Eggs, ootype not observed. Vitellaria coextensive with intestinal caeca. Haptor subcircular, velum-like, 314 (232–383; $n = 3$) long, 273 (199–331; $n = 3$) wide, with four haptor glands. Anchors dissimilar. Ventral anchor, outer 47 (39–54; $n = 4$) long, inner 44 (38–39; $n = 4$) long; base 39 (37–43; $n = 4$) long; with divergent roots; truncate superficial root; expanded deep root, mildly curved elongate shaft, slightly curved short point extending at the level of the tip of the superficial root (fig. 2f). Dorsal anchor, outer 44 (36–49; $n = 3$) long, inner 32 (31–33; $n = 3$) long; base 33 (28–37; $n = 3$) long; robust, with inconspicuous roots, expanded margin, slightly curved short shaft, elongate point extending well past the level of the tip of the inner base; anchor spine blunt (fig. 2g). Ventral bar 53 (42–67; $n = 5$) long, 12 (10–13; $n = 4$) wide, curved in the posterior direction, with bifid ends for articulation with ventral anchor (fig. 2d). Dorsal bar 53 (38–76; $n = 5$) long, 10 (7–14; $n = 5$) wide, slightly straight shaped, with rounded ends; mid-posteromedial process (fig. 2e). Hooks similar in shape, 18 (17–19; $n = 5$) long, shank elongated, without inflation, erect thumb, evenly curved shaft point (fig. 2c). Filamentous hook loop comprising 80% of the length of the shank.

Remarks

A comparative analysis of the vouchers of *C. velum* (CHIOC 38262 a–b, 38263), provided by Domingues et al. (2016), and specimens of *Chauhanellus* from *S. couma*, Caratateua, Municipality of Bragança, in Pará, Brazil, indicated that they are conspecific, mainly because they both share the same morphology of the copulatory complex, haptor velum-like, bars and anchors (fig. 2) (see also molecular data results). Moreover, the Bray–Curtis morphometric analysis of the morphological structures of *C. velum* from *S. couma* (present study) and *S. herzbergii*, *S. couma* and *S. passany* (table 5) identified a similarity of 88–95% across the specimens from each host (fig. 3; Online supplementary table S1). In addition, we provide supplementary morphological data and new illustrations of *C. velum* (fig. 2).

The ventral anchor observed in the specimens in the present study possesses a truncate superficial root and an expanded deep root (fig. 2f). This characteristic was also observed in the vouchers examined (CHIOC 38262 a–b, 38263); however, not as clearly as in the specimens of the present study, as it appears to be compressed by the coverslip sheet, making its definition difficult. It is true that the species has a truncate superficial root, although it was not clear in the drawing from the original description (Domingues et al., 2016, p. 312, fig. 27). The compression of this structure by the coverslip sheet may have caused a misinterpretation by Domingues et al. (2016). In addition, with respect to the intestinal caeca, vas deferens, genital pore, genital atrium, seminal vesicle, seminal receptacle, uterus and Mehlis' gland, none of which were observed in the original description of the species, the present study provides a better definition of these structures (see above).

Molecular data

The sequencing of the partial 18S rDNA of *C. riograndinensis* n. sp. was 1689 bp in length. Beyond this, four new partial 18S rDNA sequences were obtained for four other species of *Chauhanellus* (*C. neotropicalis* from *A. quadriscutis* – 1699 bp long, *C. hamatopeduncularoideum* from *A. rugispinis* [type-host] – 1611 bp long, *C. hypenocleithrum* from *S. proops* [type-host] – 1653 bp long and *C. velum* from *S. couma* [type-host] – 1654 bp long) (table 2).




Fig. 2. *Chauhanellus velum* Domingues, Soares & Watanabe, 2016. (a) Voucher specimen whole-mount, ventral; (b) copulatory complex; (c) hook; (d) ventral bar; (e) dorsal bar; (f) ventral anchor; (g) dorsal anchor. Scale bars: (a) 100 µm; (c) 10 µm; (b) 25 µm; (d–g) 50 µm.

The genetic divergence between *Chauhanellus* species and monogenoid species from Siluriformes was compared, varying from 3.2 to 11.8% (table 6). Interspecific divergence within *Chauhanellus* ranged from 0.3 to 4.6% (7–189 bp). The genetic divergence among *C. riograndinensis* n. sp. and other *Chauhanellus* species was between 2.3 and 4.3% (39–77 pb). The divergence between *C. riograndinensis* n. sp. and the most similar morphological species, *C. susamlimae*, was 2.2% (39 pb). The smallest interspecific distance was observed between *C. boegeri* Domingues & Fehlaue, 2006 and *C. neotropicalis* at only 0.3% (7 bp), while *C. velum* was the most genetically distant species of *Chauhanellus* (4.6%). There was no intraspecific divergence between sequences of *C. velum* from the distinct hosts *S. herzbergii* and *S. couma*.

Phylogenetic evidence

The ML and BI phylogenetic analyses based on the 18S rDNA gene converged with similar topologies, and only the BI tree

was presented, with the statistical support of both methods (fig. 4). Monogenoid species from Siluriformes fish arose in clade S, with highly supportive nodes in both ML and BI analyses, and were further divided into three subclades, S1, S2 and S3 (fig. 4). S1 exclusively comprises parasites of freshwater catfish from the Oriental region: *Mizellus longicirrus* (Tripathi, 1959) from Siluridae, *Bychowskyella* spp. from Heteropneustidae and Clariidae and *Thaparocleidus* spp. from Siluridae. S2 clustered exclusively parasites of marine catfish (Ariidae) from South America and the Oriental region: *Susanlimocotyle narina* from *S. herzbergii* arises forming a strongly supported lineage closely related to *Hamatopeduncularia* spp. (*Hamatopeduncularia arii* Yamaguti, 1953, *Hamatopeduncularia bifida* Illa, Shameem, Serra, Melai, Mangam, Basuri, Petroni & Modeo, 2019, *Hamatopeduncularia elongata* Lim, 1996, *Hamatopeduncularia thalassini* Bychowsky & Nagibina, 1968 [all from *Arius jella* Day] and *Hamatopeduncularia madhaviae* Illa, Shameem, Serra, Melai, Mangam, Basuri, Petroni & Modeo, 2019, from *Plicofollis dussumieri* [Valenciennes]) from Oriental ariids (fig. 4).

Table 5. Comparative measurements (in μm) of specimens of *Chauhanellus velum* Domingues, Soares & Watanabe 2016 from *Sciades couma*, *Sciades herzbergii* and *Sciades passany* from different ariids' hosts.

	<i>S. couma</i> ^a	<i>N</i>	<i>S. couma</i> ^b	<i>N</i>	<i>S. herzbergii</i> ^b	<i>N</i>	<i>S. passany</i> ^b	<i>N</i>
Male copulatory organ length	103(85–113)	4	110	1	104(97–112)	2	85	1
Ventral bar								
Length	53(42–67)	5	46	1	49(46–52)	2	55(52–57)	2
Width	12(10–13)	4	11	1	10(9–11)	2	14(12–15)	2
Dorsal bar								
Length	53(38–76)	5	52	1	49 (41–57)	2	67 (64–69)	2
Width	10(7–14)	5	12	1	14 (12–15)	2	15	2
Ventral anchor								
Outer	47(39–54)	4	62	1	43 (41–44)	2	37	1
Inner	44(38–39)	4	–	–	28 (25–31)	2	37	1
Base	39(37–43)	4	40	1	29 (27–31)	2	45	1
Dorsal anchor								
Outer	44(36–49)	3	–	–	123	1	165	1
Inner	32(31–33)	3	–	–	115	1	150	1
Base	33(28–37)	3	–	–	75	1	63	1
Hooks length	18(17–19)	5	16	2	16 (15–17)	2	16	2

N, number of parasite.

^aPresent study.

^bDomingues et al. (2016).

Subclade S3 arises in a sister position to clade S2 and comprises *Chauhanellus* species that are exclusive parasites of marine catfish (Ariidae) from South America: *C. hamatopeduncularoideum* from *A. rugispinis*, and *C. susamlimae* from *S. herzbergii*, appearing as the derived species in a sister position to *C. hypenocleithrum* from *S. proops*. In turn, with strong support, this lineage appears as sister to the one composed of *C. boegeri* from *S. herzbergii* and *C. neotropicalis* from *A. quadriscutis*. *Chauhanellus riograndinensis* n. sp., a parasite of *G. barbuis* and *G. genidens*, and *C. velum* from *S. couma* and *S. herzbergii*, arose as early divergent *Chauhanellus* species.

Discussion

Several genera have been proposed for the monogenoids of ariid fishes worldwide (Soares et al., 2021a, b). *Fridericianella*,

Neocalceostoma, *Neocalceostomoides* and *Thysanotohaptor* are harboured in the Neocalceostomatidae and *Udonella* in Udonellidae (Soares et al., 2021a, b). The four others, *Chauhanellus*, *Hamatopeduncularia*, *Neotetraonchus* and *Susanlimocotyle* belong to the Dactylogyridae (Soares et al., 2021a).

Of the six valid *Chauhanellus* species so far reported to be infesting South American ariid catfishes (table 1), only three have available sequences (Soares et al., 2021a). In the present study, we sequenced all known South American *Chauhanellus* species, as well as the new species described here.

Only two prior reports exist of *Chauhanellus* species parasitizing *G. barbuis* and *G. genidens* from the Brazilian coast: *C. boegeri* on *G. barbuis* and *G. genidens* from Guaratuba, Paraná, Brazil (Domingues & Fehlaue, 2006); and an undetermined species of




Fig. 3. Morphometric similarity dendrogram by Bray-Curtis method for *Chauhanellus velum* from different hosts and the new species described.

Table 6. Pairwise genetic identities of 18S rDNA sequences selected from Dactylogyridae species from Siluriformes adjusted for missing data.

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19
1. <i>Mizelleus longicirrus</i> KR296801	–	189	162	214	225	454	465	208	192	430	194	399	403	184	346	237	246	222	223
2. <i>Bychowskyella fossilisi</i> KT852454	7.6	–	56	174	188	195	206	182	164	186	168	186	183	155	323	225	223	205	206
3. <i>Bychowskyella tchangii</i> KT852455	7.2	1.6	–	150	159	197	199	176	158	176	163	176	175	151	296	195	198	179	180
4. <i>Thaparocleidus gangus</i> KX364088	9.8	7.2	6.7	–	114	167	184	164	131	148	138	153	151	135	291	177	177	152	155
5. <i>Thaparocleidus wallagonius</i> KX364085	10.2	8.1	7.4	5.3	–	178	198	174	152	167	147	169	167	144	299	198	199	174	177
6. <i>Susanlimocotyle narina</i> MW144824	11.2	9.9	9.5	8.6	8.9	–	242	115	78	196	76	196	193	74	259	110	105	76	75
7. <i>Chauhanellus velum</i> MW144823 ^a	11.8	10.3	9.8	9.4	9.9	6.6	–	0	77	189	84	179	184	80	279	119	131	104	103
8. <i>C. velum</i>^b	11.8	10.3	9.8	9.4	9.9	6.6	0	–	76	78	83	78	73	80	214	107	118	95	96
9. <i>Chauhanellus riograndinensis</i> n. sp	10.5	9.1	8.5	7.6	8.6	4.4	4.3	4.3	–	45	51	43	39	39	195	77	105	55	57
10. <i>Chauhanellus boegeri</i> MW132134	10.4	9.1	8.6	7.8	8.4	4.3	4.4	4.4	2.7	–	7	104	99	32	251	89	96	62	60
11. <i>Chauhanellus neotropicalis</i>	10.6	9.3	8.8	8	8.4	4.4	4.6	4.6	3	0.3	–	36	32	36	202	82	88	59	59
12. <i>Chauhanellus hamatopeduncularoideum</i>	10.7	9.3	8.6	7.9	8.4	4.6	4.3	4.3	2.3	1.7	1.8	–	44	21	255	93	100	67	66
13. <i>Chauhanellus susamlimae</i> MW144439	10.7	9.3	8.6	8.1	8.5	4.3	4.1	4.1	2.2	1.5	1.7	0.6	–	15	251	86	93	60	61
14. <i>Chauhanellus hypenocleithrum</i>	10.6	8.9	8.3	7.9	8.3	4.3	4.4	4.4	2.2	1.8	2	1	0.8	–	194	80	86	59	60
15. <i>Hamatopeduncularia elongate</i> KT252896	15.2	13.4	12.9	13	13	11.2	12.4	12.4	10.7	10.9	11.1	10.9	10.9	11.3	–	232	220	232	232
16. <i>Hamatopeduncularia madhaviae</i> KT252898	11.0	9.7	9.2	8.6	9.1	5.1	6	6	4.3	4.3	4.4	4.4	4.1	4.6	9.5	–	75	68	68
17. <i>Hamatopeduncularia bifida</i> KT252899	11.2	9.8	9.3	8.9	9.6	4.9	6.7	6.7	4.6	4.3	5	4.9	4.6	5	9.1	3.2	–	62	65
18. <i>Hamatopeduncularia thalassini</i> KT252900	10.5	9.2	8.8	7.7	8.4	3.7	5.4	5.4	3.2	3.5	3.5	3.6	3.4	3.6	9.7	3.2	2.8	–	15
19. <i>Hamatopeduncularia arii</i> KT252895	10.5	9.3	8.9	8	8.7	3.7	5.5	5.5	3.4	3.5	3.5	3.6	3.5	3.6	9.7	3.2	3.1	0.7	–

The upper triangular matrix shows the number of differences in nucleotides and the lower triangular matrix shows the differences in terms of percentage of nucleotides.

Sequences obtained in the present work are shown in in boldface type.

^aIn *Sciades herzbergii*.

^bIn *Sciades couma*.

Despite the fact that *G. barbuis* from Cananéia, São Paulo, Brazil (present study) and *G. barbuis* and *G. genidens* from Guaratuba, Paraná state, Brazil (Domingues & Fehlaue, 2006) had been examined, the current new species was not found in these locations. On the other hand, *C. boegeri* from Guaratuba, Paraná, Brazil, was found in *G. barbuis* and *G. genidens* obtained from the current study area (i.e. Cananéia, São Paulo, south-eastern Brazil and Rio Grande, Rio Grande do Sul state, southern Brazil), as well as on *S. couma* from Caratateua, Pará, northern Brazil (table 2). According to Lim *et al.* (2001), the presence of different species of *Chauhanellus* on the same host species in different biogeographical regions (e.g. *Arius maculatus* (Thunberg) from Hainan, China, and from the western coast of Peninsular Malaysia) suggests that habitat differences may affect the presence or absence of certain parasite species. However, the limitations of insufficient numbers of host samples and potentially misidentified host species should also be considered.

Conclusion

The present study provides morphological and molecular characterization (partial 18S rDNA) of a new species of *Chauhanellus* and new 18S rDNA sequences of *Chauhanellus* spp. from South American ariids. Our results showed that *C. riograndinensis* n. sp. and *C. velum* represent two early divergent lineages within *Chauhanellus* from South America. Moreover, the confirmation of the conspecificity of *Chauhanellus* specimens from *S. couma* (present study) and *C. velum* from *S. couma*, *S. herzbergii*, and *S. passany* reported by other authors emphasizes the need for an integrative taxonomic approach to ensure accurate delimitation of monogenoid species. Finally, we suggest one way for future studies that seek to test the monophyly of *Chauhanellus* and *Hamatopeduncularia*.

Supplementary material. To view supplementary material for this article, please visit <https://doi.org/10.1017/S0022149X23000135>.

Acknowledgement. We thank Marcelo Knoff, Coleção Helmintológica do Instituto Oswaldo Cruz (Brazil), who allowed us access to specimens under his care.

Financial support. The present study was partly supported by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (the Coordination for the Improvement of Higher Education Personnel) – Brazil (CAPES) – Finance Code 001. G.B. Soares was supported by a doctoral scholarship #2017/17531-0, São Paulo Research Foundation (FAPESP). M. V. Domingues and E. A. Adriano received research productivity grants from the Brazilian Fostering Agency CNPq (respectively, grant #309896/2019-3 and #301886/2016-4).

Conflicts of interest. None.

Ethical standards. Permission to collect the hosts was given by the Brazilian Institute of Environment and Renewable Natural Resources (IBAMA) under a License for the Collection of Zoological Material (SISBio No. 60666-2) and for access to genetic heritage (Sisgen No. AD28DC2).

References

Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W and Lipman DJ (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Research* **25**(17), 3389–3402.

Barriel V (1994) Molecular phylogenies and how to code insertion/deletion events. *Comptes Rendus de l'Académie des Sciences Serie III, Sciences de la Vie* **317**(7), 693–701.

Boeger WA and Vianna RT (2006) Monogenoidea. pp. 42–116. In Thatcher V (Ed.) *Aquatic biodiversity in Latin America*. Sofia–Moscow, Amazon fish parasites eds Pensoft Publishers.

Bray JR and Curtis JT (1957) An ordination of the upland forest communities of southern Wisconsin. *Ecological Monographs* **27**(4), 325–349.

Bush AO, Lafferty KD, Lotz JM and Shostak W (1997) Parasitology meets ecology on its own terms: Margolis *et al.* revisited. *Journal of Parasitology* **83**(4), 575–583.

Bychowsky BE and Nagibina LF (1969) Ancyrocephalinae (Dactylogyridae, Monogenoidea) from fishes of the Family Ariidae. *Parazitologiya* **3**(4), 357–368.

CorelDraw Com (2014) CorelDRAW Graphics Suite. Available at https://www.coreldraw.com/en/product/coreldraw/?x-vehicle=ppc_brkws&glid=Cj0KCQIA0oagBhDHARIsAI-BbgelYakN3YSfEHFBPCDr2uXHVfRcFy_75S6BvobnPpFBpzzCyGz2g_UaAj2-EALw_wcB (accessed April 2022).

Darriba D, Taboada GL, Doallo R and Posada D (2012) jModelTest 2: more models, new heuristics and parallel computing. *Nature Methods* **9**(8), 772.

Domingues MV and Fehlaue KH (2006) New species of *Chauhanellus* (Monogenoidea, Platyhelminthes) from the gills of Southern Atlantic marine catfishes (Siluriformes, Ariidae) on the Neotropical Region. *Zootaxa* **1365**(1), 61–68.

Domingues MV, Soares GB and Watanabe A (2016) Monogenoidea (Polyonchoinea: Dactylogyridae) parasitizing the gills of marine catfish (Siluriformes: Ariidae) inhabiting the Atlantic Amazon Coast of Brazil. *Zootaxa* **4127**(2), 301–326.

Fricke R, Eschmeyer WN and Van der Laan R (2022) *Eschmeyer's catalog of fishes*. Available at <http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp> (accessed 7 July 2022).

Geer LY, Marchler-Bauer A, Geer RC, Han L, He J, He S, Liu C, Shi W and Bryant SH (2009) The NCBI BioSystems database. *Nucleic Acids Research* **38**(Database issue), D492–D496.

Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W and Gascuel O (2010) New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Systematic Biology* **59**(3), 307–321.

Humason GL (1979) *Animal tissue techniques*. 661 pp. San Francisco, WH Freeman.

Illa K, Shameem U, Serra V, Melai M, Mangam S, Basuri CK, Petroni G and Modeo L (2019) Multidisciplinary investigation on the catfish parasite *Hamatopeduncularia* Yamaguti, 1953 (Monogenoidea: Dactylogyridae): description of two new species from India, and phylogenetic considerations. *The European Zoological Journal* **86**(1), 132–155.

International Commission on Zoological Nomenclature (2012) Amendment of Articles 8, 9, 10, 21 and 78 of the International Code of Zoological Nomenclature to expand and refine methods of publication. *Bulletin of Zoological Nomenclature* **69**(1), 161–169.

Kearn GC and Whittington ID (1994) Ancyrocephaline monogeneans of the genera *Chauhanellus* and *Hamatopeduncularia* from the gills of the blue catfish, *Arius graeffei*, in the Brisbane River and Moreton Bay, Queensland, Australia, with descriptions of four new species. *International Journal for Parasitology* **24**(4), 569–588.

Kearse M, Moir R, Wilson A, *et al.* (2012) Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* **28**(12), 1647–1649.

Kritsky DC (2012) Dactylogyrids (Monogenoidea: Polyonchoinea) parasitizing the gills of snappers (Perciformes: Lutjanidae): revision of *Euryhaliotrema* with new and previously described species from the Red Sea, Persian Gulf, the eastern and Indo-West Pacific Ocean, and the Gulf of Mexico. *Zoologia* **29**(3), 227–276.

Kumar S, Stecher G and Tamura K (2016) MEGA7: molecular evolutionary genetics analysis version 7.0. *Molecular Biology and Evolution* **33**(7), 1870–1874.

Lim LHS (1994) *Chauhanellus* Bychowsky & Nagibina, 1969 (Monogenea) from ariid fishes (Siluriformes) of Peninsular Malaysia. *Systematic Parasitology* **28**(1), 99–124.

Lim LHS (1996) Eight new species of *Hamatopeduncularia* Yamaguti, 1953 (Monogenea: Ancyrocephalidae) from Ariidae of Peninsular Malaysia. *Systematic Parasitology* **33**(1), 53–71.

Lim LHS, Timofeeva TA and Gibson DI (2001) Dactylogyridae monogeneans of the siluriform fishes of the old world. *Systematic Parasitology* **50**(3), 159–197.

- Littlewood DTJ and Olson PD** (2001) SSU rDNA and the Platyhelminthes: signal, noise, conflict and compromise. pp. 262–278. In Littlewood DTJ and Bray RA (Eds) *Interrelationships of the Platyhelminthes*. London, Taylor & Francis.
- Littlewood DTJ, Rohde K and Clough KA** (1998) The phylogenetic position of *Udonella* (Platyhelminthes). *International Journal for Parasitology* **28**(8), 1241–1250.
- Littlewood DTJ, Rohde K and Clough KA** (1999) The interrelationships of all major groups of Platyhelminthes: phylogenetic evidence from morphology and molecules. *Biological Journal of the Linnean Society* **66**(75), 114.
- Littlewood DTJ, Waeschenbach A and Nikolov PN** (2008) In search of mitochondrial markers resolving the phylogeny of cyclophyllidean tapeworms (Platyhelminthes, Cestoda) - a test study with Davaineidae. *Acta Parasitologica* **53**(2), 133–144.
- Marceniuk AP, Menezes NA and Brito MR** (2012) Phylogenetic analysis of the family Ariidae (Ostariophysi: Siluriformes), with a hypothesis on the monophyly and relationships of the genera. *Zoological Journal of the Linnean Society* **165**(3), 534–669.
- Mizelle JD and Klucka AR** (1953) Studies on monogenetic trematodes. XVI. Dactylogyridae from Wisconsin fishes. *American Midland Naturalist* **49**(3), 720–733.
- Muller MI, Ceccarelli PS and Ueta MT** (2016) Supplementary studies on *Anacanthorus* and *Mymarothecium viatorum* (Monogenea: Dactylogyridae) from *Piaractus mesopotamicus* (Characiformes: Serrasalminidae) in Brazil. *Acta Parasitologica* **61**(3), 508–515.
- Plaisance L, Littlewood DTJ, Olson PD and Morand S** (2005) Molecular phylogeny of gill monogeneans (Platyhelminthes, Monogenea, Dactylogyridae) and colonization of Indo-West Pacific butterflyfish hosts (Perciformes, Chaetodontidae). *Zoologica Scripta* **34**(4), 425–436.
- Pleijel F, Jondelius U, Norlinder E, Nygren A, Oxelman B, Schander C, Sundberg P and Thollesson M** (2008) Phylogenies without roots? A plea for the use of vouchers in molecular phylogenetic studies. *Molecular Phylogenetics and Evolution* **48**(1), 369–371.
- Rambaut A** (2022) *FigTree v1.4.4: Tree Figure Drawing Tool*. Available at <http://tree.bio.ed.ac.uk/software/figtree/> (accessed 26 June 2022).
- Rambaut A, Drummond AJ, Xie D, Baele G and Suchard MA** (2018) Posterior summarisation in Bayesian phylogenetics using Tracer 1.7. *Systematic Biology* **67**(5), 901–904.
- Rasband WS** (2022) *ImageJ*. Bethesda, Maryland, USA, US National Institutes of Health. Available at <http://imagej.nih.gov/ij/> (accessed 22 July 2022).
- Rastogi P, Kumar K and Singh HS** (2004) Review of the genus *Chauhanellus* (Young, 1967) Bychowsky and Nagibina 1969 with a report on a new species from freshwater fishes of Meerut (U.P.), India. *Uttar Pradesh Journal of Zoology* **24**(2), 121–128.
- Ronquist F, Teslenko M, Mark PV, et al.** (2012) MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* **61**(3), 539–542.
- Soares GB, Magalhães KX, Silva AC, Carneiro JS, Barbosa LL, Costa NGS and Domingues MV** (2019) Monogeneoids (Polyonchoinea, Dactylogyridae) from *Hydrolycus armatus* (Characiformes, Cynodontidae) with the description of a new species of *Rhinoxenus* and the proposal of a new genus from the Xingu River, Pará, Brazil. *Zootaxa* **4700**(2), 229–245.
- Soares GB, Domingues MV and Adriano EA** (2021a) An integrative taxonomic study of *Susanlimocotyle narina* n. gen. n. sp. (Monogeneoidea, Dactylogyridae) from the nasal cavities of a marine catfish (Siluriformes, Ariidae) from the Atlantic Amazon Coast of Brazil and new molecular data of *Chauhanellus* spp. *Parasitology International* **81**(1), 102271.
- Soares GB, Domingues MV and Adriano EA** (2021b) Morphological and molecular characterization of *Udonella brasiliensis* n. sp. (Monogeneoidea), an epibiont on *Caligus* sp. parasite of Ariidae from the southeastern coast of Brazil. *Parasitology International* **83**(1), 102371.
- Soares GB, Adriano EA, Domingues MV and Balbuena JA** (2022) Diversification processes between monogeneoids (Dactylogyridae) and their marine catfish (Siluriformes: Ariidae) from the Atlantic coast of South America. *Parasitology* **150**(2), 184–194.
- Soo OYM and Tan WB** (2021) *Hamatopeduncularia* Yamaguti, 1953 (Monogenea: Ancyrodiscoididae) from catfish off peninsular Malaysia: description of two new species and insights on the genus. *Parasitology International* **81**(1), 102282.
- Sun Y, Li M and Yang T** (2014) Studies on *Lethrinitrema* Lim & Justine, 2011 (Monogenea: Dactylogyridae), with the description of two new species, a key to the genus and a phylogenetic analysis based on rDNA sequences. *Systematic Parasitology* **88**(2), 119–139.
- Verma C, Chaudhary A and Singh HS** (2017a) Redescription of two species of *Thaparocleidus* (Monogenea: Dactylogyridae), with the description of *T. armillatus* sp. n. from *Wallago attu* and a phylogenetic analysis based on 18S rDNA sequences. *Acta Parasitologica* **62**(3), 652–665.
- Verma C, Chaudhary A and Singh HS** (2017b) Morphology, molecular and systematic analyses of *Bychowskyella* (Monogenea: Dactylogyridae) in siluriform fish from India. *Journal of Helminthology* **91**(2), 197–205.
- Wilkinson M** (1995) Coping with abundant missing entries in phylogenetic inference using parsimony. *Systematic Biology* **44**(4), 501–514.
- Winnepenninckx B and Backeljau T** (1996) 18S rRNA alignments derived from different secondary structure models can produce alternative phylogenies. *Journal of Zoological Systematics and Evolutionary Research* **34**(3), 135–143.
- Wu XY, Li AX, Zhu XQ and Xie MQ** (2005) Description of *Pseudorhabdosynochus seabassi* sp. n. (Monogenea: Diplectanidae) from *Lates calcarifer* and revision of the phylogenetic position of *Diplectanum grouperi* (Monogenea: Diplectanidae) based on rDNA sequence data. *Folia Parasitologica* **52**(3), 231–240.