

## Meat intake in relation to semen quality and reproductive hormone levels among young men in Spain

Ana B. Maldonado-Cárceles<sup>1,2\*</sup>, Lidia Mínguez-Alarcón<sup>2,3</sup>, Jaime Mendiola<sup>2</sup>, Jesús Vioque<sup>4,5</sup>, Niels Jørgensen<sup>6</sup>, Julián J. Árense-Gonzalo<sup>2</sup>, Alberto M. Torres-Cantero<sup>2,7</sup> and Jorge E. Chavarro<sup>1,8,9</sup>

<sup>1</sup>Department of Nutrition, Harvard School of Public Health, Boston, MA 02115, USA

<sup>2</sup>Department of Health and Social Sciences, Division of Preventive Medicine and Public Health, University of Murcia School of Medicine, IMIB-Arrixaca, 30100 Espinardo, Murcia, Spain

<sup>3</sup>Department of Environmental Health, Harvard School of Public Health, Boston, MA 02115, USA

<sup>4</sup>CIBER Epidemiology and Public Health (CIBERESP), Institute of Health-ISCIII, 28029 Madrid, Spain

<sup>5</sup>Department of Public Health, Alicante Institute for Health and Biomedical Research (ISABIAL-FISABIO Foundation), Miguel Hernández University, 03550 Alicante, Spain

<sup>6</sup>Department of Growth and Reproduction, International Center for Research and Research Training in Endocrine Disruption of Male Reproduction and Child Health (EDMaRC), Rigshospitalet, University of Copenhagen, DK-2100 Copenhagen, Denmark

<sup>7</sup>Regional Campus of International Excellence “Campus Mare Nostrum”, University of Murcia, 30003 Murcia, Spain

<sup>8</sup>Department of Epidemiology, Harvard School of Public Health, Boston, MA 02115, USA

<sup>9</sup>Channing Division of Network Medicine, Department of Medicine, Brigham and Women’s Hospital and Harvard Medical School, Boston, MA 02115, USA

(Submitted 24 July 2018 – Final revision received 13 October 2018 – Accepted 31 October 2018 – First published online 18 December 2018)

### Abstract

The aim of this study was to assess the associations of intake of different types of meat with semen parameters and reproductive hormones in healthy young men. This cross-sectional study included 206 men, 18–23 years, from Southern Spain. All men completed a validated FFQ, underwent a physical examination, and provided blood and semen samples. Multivariable linear regression was used to evaluate the associations between meat intake with semen quality parameters and reproductive hormones. Total meat intake was unrelated to semen quality or reproductive hormone levels. When subgroups of meat were separately considered, however, shellfish intake was positively related to progressive motility. The adjusted percentages of progressively motile spermatozoa for men in increasing quartiles of shellfish intake were 45.2, 42.0, 49.4 and 53.2% with a significant linear trend across quartiles ( $P_{\text{trend}} \leq 0.001$ ). In contrast, men who consumed organ meats had significantly lower progressive sperm motility (51.5 v. 42.8%;  $P=0.001$ ) and higher luteinising hormone levels (4.0 v. 4.6 IU/l;  $P=0.03$ ) compared with men who did not consume organ meats. Intake of shellfish and organ meats was low in this population, however. Given the scarcity of data on the relation between specific types of meat with semen quality and reproductive hormone levels, additional research is needed to confirm or refute these findings.

**Key words:** Meat: Fish: Shellfish: Semen motility: Semen quality parameters

During the last decades, decreasing trends in sperm counts have been reported in multiple studies<sup>(1–4)</sup>. The reasons for this downward trend have not been fully elucidated, but exposures adversely affecting prenatal testicular development as well as exposures and lifestyle factors during childhood and young adulthood have been suggested<sup>(4,5)</sup>. However, relatively little work has been devoted to understanding the modifiable determinants of semen quality, as a marker of men’s reproductive potential<sup>(6)</sup>, which could lead to the design of clinical and public health interventions<sup>(7)</sup>.

Diet is a modifiable risk factor that can be assessed in clinical practice and inform actions for disease prevention<sup>(8)</sup>. Previous epidemiological studies suggest that diet may be associated with semen parameters<sup>(9–14)</sup>. Animal food intake, and in particular intake of red meats and fish, has received particular attention since their nutritional profile and their contamination with persistent organic pollutants and hormonal residues may affect testicular function<sup>(11,15–19)</sup>. Although literature is scarce, red meats have consistently been associated with worse semen quality, while fish intake appears to have the opposite

**Abbreviations:** E2, oestradiol; LH, luteinising hormone.

\* **Corresponding author:** A. B. Maldonado-Cárceles, fax +34 868 88 3947, email [anab.maldonadocarceles@gmail.com](mailto:anab.maldonadocarceles@gmail.com)

relation<sup>(19–22)</sup>. However, these studies have been conducted in different populations and settings, with diverse age and health status, and only one previous work was performed in healthy young men<sup>(19)</sup>. Therefore, the aim of this study is to investigate the association of intakes of meats, including red, white, processed and fish meats, with semen quality parameters and reproductive hormones among healthy young men from Spain.

## Methods

### Study population

The Murcia Young Men's Study (MYMS) was a cross-sectional study conducted in 2010–2011 that enrolled healthy men between 18 and 23 years of age at the University of Murcia (Region of Murcia, Spain). MYMS is part of a multicentre international study (Finland, Denmark, Spain and USA) aimed at evaluating the role of environmental contaminants on semen quality. The same study population has been used in previously published works<sup>(10,12,23,24)</sup>. Participants were recruited through flyers posted at university campuses. Men were eligible to participate if they were born in Spain after 31 December 1987, and were able to contact their mother to complete a questionnaire.

A total of 240 men contacted the study staff. Of these, 223 (92.9%) met the eligibility criteria and 215 completed a study visit and agreed to participate in the study. During the visit, men underwent a physical examination, completed questionnaires concerning diet, lifestyle, medical and reproductive history and provided semen and blood samples. Six men who reported implausible total energy intake (>20920 kJ/d) and three who had a hydrocele were excluded from the analysis. The final analytical sample comprised 206 men (85.8%). Participants received a €50 gift card for their participation in the study. Written informed consent was obtained from all subjects. The Research Ethics Committee of the University of Murcia approved this study (no. 495/2010, approved 14 May 2010).

### Physical examination

Men's height and weight were measured using a wall stadiometer and digital scale (Tanita SC 330-S) to determine their BMI. The presence of varicocele, hydrocele or other scrotal abnormalities was recorded. Testicular volume was assessed with a Prader orchidometre (Andrology Australia). All physical examinations were performed by the same investigator (J. M.) to minimise variability in study measures.

### Dietary assessment

Participants completed a 101-item semi-quantitative FFQ previously validated in several adult populations in Spain<sup>(25,26)</sup>. Men reported their usual intake of foods and supplements over the past year. The FFQ had nine categories for frequency of consumption, ranging from never to  $\geq 6$  times/d. Total meat intake was defined as the sum of red meat, white meat and fish. We grouped specific meat types in seven groups (online Supplementary Table S1): three types of red meat, one type white meat and three types of fish.

### Semen analysis

Men were instructed to abstain from ejaculation for 48 h before sample collection but were not excluded if this was not the case. At the time of sample collection, they also provided information on time of the previous ejaculation. Semen samples were obtained by masturbation at the clinic, and lubricants were not used. Ejaculate volume was estimated by sample weight, assuming a semen density of 1.0 g/ml. Sperm concentration was evaluated by haemocytometer (Improved Neubauer; Hauser Scientific Inc.). Sperm motility was evaluated as either motile or immotile according to the WHO criteria<sup>(27)</sup>, and the percentage of total motile sperm and progressive motility were calculated. Smears for morphology were made, air-dried, fixed, Papanicolaou stained and evaluated using strict criteria<sup>(28)</sup>. Total sperm count (volume  $\times$  sperm concentration) was also calculated. The same specialist biologist carried out all the semen analyses. An external quality control on semen samples throughout the study period was carried out in collaboration with the University of Copenhagen's Department of Growth and Reproduction. To assess inter-laboratory variation in sperm concentration analysis, five sets of duplicate semen samples (600  $\mu$ l each) were sent by mail during the study period from the University of Copenhagen's Department of Growth and Reproduction to the Murcia Andrology Laboratory. The samples were blinded, non-diluted fresh sperm specimens from regular semen donors that were preserved by adding 10  $\mu$ l of a 3 M sodium azide solution per 1 ml of the ejaculate after liquefaction. Results showed no systematic differences. The mean inter-examiner CV was 4.0%, ranging between 1.7 and 7.1%.

### Reproductive hormone measurement

Blood samples were drawn from men's cubital veins and centrifuged; the serum was separated, stored and frozen at  $-80^{\circ}\text{C}$ . Serum samples were shipped to Rigshospitalet (Copenhagen, Denmark) and stored at  $-20^{\circ}\text{C}$  until hormone analysis was performed. The methods have been outlined previously<sup>(29)</sup>. Briefly, hormone assessments were performed simultaneously to reduce intra-laboratory variations. Serum levels of follicle-stimulating hormone (FSH), luteinising hormone (LH) and sex hormone-binding globulin (SHBG) were determined using time-resolved immunofluorometric assays (DELFLIA; PerkinElmer). Intra- and inter-assay variations were <5% in each of the three assays. Serum total testosterone (TT) levels were determined using a time-resolved fluoroimmunoassay (dissociation enhanced lanthanide fluoroimmunoassay; DELFLIA) with intra- and inter-assay variations of <8%. Oestradiol (E2) was measured by RIA (Pantex) with an intra-assay variation of <8% and an inter-assay variation of <13%. Inhibin B levels were determined by a specific two-sided enzyme immunometric assay (Oxford Bio-Innovation Ltd) with intra- and inter-assay variations of 13 and 18%, respectively. When inhibin B was above approximately 100 pg/ml, the intra-assay variation was <7% and the inter-assay variation was <6%. The majority of the men had levels above 100 pg/ml. Calculated free testosterone (FT) levels were determined using the equation of Vermeulen *et al.*<sup>(30)</sup>, assuming a fixed albumin of 43.8 g/l.

### Statistical analysis

Sample size was calculated assuming anticipating an effect size of 0.15, for a prespecified  $\alpha$  and power level of 5 and 80%, respectively. With a non-response rate of 20%, a minimum sample size of 157 subjects was needed<sup>(31)</sup>. Participants' characteristics were summarised using the median and interquartile range (IQR) and absolute frequency and percentage across quartiles of total meat intake. Statistical differences across quartiles of meat intake, red meat intake and fish meat intake were assessed using Kruskal–Wallis tests for continuous variables and  $\chi^2$  tests for categorical variables. Semen volume, sperm concentration, total sperm count, morphologically normal sperm, FSH levels and E2 levels were  $\log(\ln)$  transformed before analysis to improve normality. After analysis, we back-transformed them to show values in original scale. Multivariate linear regression models were used to investigate the association of each group of meat intake (in categories) with semen quality parameters and reproductive hormones with raw data and after adjusting for confounders. We used ANCOVA to calculate adjusted reproductive outcomes and 95% CI as dependent continuous variables for each category of meat intake and covariates as independent variables. An association was considered when we found a statistically significant linear trend among categories of meat intake as ordinal variable. Explicit substitution analyses were performed for the foods for which the unspecified substitution was statistically significant.

Confounding was assessed based on participants' characteristics associated with meat intake and semen parameters or reproductive hormones. Based on this criteria, models were adjusted for age (years), BMI ( $\text{kg}/\text{m}^2$ ), smoking (current smoker *v.* not current smoker), physical activity (h/week), television (TV) watching (h/week) and total energy intake (kJ/d). The remaining types of meat (continuous) and dietary patterns (continuous) were also included into the models to address the possibility that observed associations would be explained by the overall food choices rather than intake of specific meats. Specifically, as described elsewhere in this population<sup>(23)</sup>, we used principal components analysis with orthogonal transformation based on predefined food groups<sup>(32)</sup> to obtain uncorrelated factors (dietary patterns) with simpler structures. Terms for intake of meats were excluded from this analysis. We retained the first two factors (online Supplementary Table S2) based on the amount of variance explained by the factor, the scree plot and the interpretability of the factors. Every subject was given a score for the two identified patterns according to their food consumption, and each participant appears in results for both dietary patterns. All analyses of semen parameters were also adjusted for abstinence time. Analyses of sperm motility were additionally adjusted for time to start semen analysis, and analyses of reproductive hormones were adjusted for time of blood draw. Analyses were performed with the statistical software IBM SPSS 21.0 (IBM Corporation).

### Results

Participants were predominantly Caucasian (97.6%) and non-smokers (68.1%). They had a median age of 20.5 (IQR 19.6–21.5)

years and BMI of 23.7 (IQR 21.8–25.4)  $\text{kg}/\text{m}^2$ . The median physical activity and TV watching was 9 and 20 h/week, respectively (Table 1, online Supplementary Table S3). A total of seventy-one subjects had a surgical scar in the genital area or lower abdomen mostly due to frenectomies or circumcision (53.5%), appendectomy (16.9%) and inguinal hernia repair (9.9%) (data not shown). The most consumed meat product was processed red meat (29%), followed by dark meat fish (22%), white meat (18%), unprocessed red meat (12%), white meat fish (11%), shellfish (5%) and organ meats (3%). Men in highest quartile of total meat intake had a higher total energy intake. Meat intake was positively related to physical activity and to the Mediterranean and Western dietary patterns, and inversely related to TV watching. Red meat intake was positively related to energy intake and the Western dietary pattern, and negatively related to TV watching. Fish intake was positively associated with energy intake, the Mediterranean dietary pattern and physical activity. No other subject characteristics were significantly different across quartiles of total meat intake, red meat or fish intake (Table 1, online Supplementary Table S3). Semen parameters and reproductive hormone levels were within normal limits for healthy adult men (Table 1, online Supplementary Table S3).

Total meat intake was unrelated to semen quality parameters (Table 2, online Supplementary Table S4) or reproductive hormone levels (Table 3, online Supplementary Table S5). Similarly, when the major categories of meat were examined, we found no association of intakes of red meats, white meats or fish with semen quality parameters (Table 2) or reproductive hormone levels (Table 3).

We then evaluated intake of sub-categories of meat in relation to semen quality and reproductive hormones (online Supplementary Tables S6–S9). In these analyses, shellfish intake was positively related to progressive sperm motility (online Supplementary Tables S6 and S8). The multivariate adjusted progressive sperm motility for men in increasing quartiles of shellfish intake was 45.2 (95% CI 42.1, 48.3), 42.0 (95% CI 38.2, 45.7), 49.4 (95% CI 46.9, 51.8) and 53.2 (95% CI 50.0, 56.4)% ( $P_{\text{trend}} < 0.001$ ) (Fig. 1, online Supplementary Table S8). This association was similar when we used men who reported no intake of shellfish as the reference category and divided the remaining men in quartiles of intake. In this analysis, the multivariable adjusted progressive sperm motility for men who did not consume shellfish and men in increasing quartiles of intake was 44.8, 44.7, 42.2, 49.4 and 53.2%, respectively ( $P_{\text{trend}} < 0.001$ ), and similar pattern was found for total sperm motility. Furthermore, there was an inverse relation between shellfish intake and E2 levels (online Supplementary Table S4). The multivariable-adjusted serum E2 concentrations for men in increasing quartiles of shellfish intake were 77.2, 86.6, 76.1 and 67.8 pmol/l ( $P_{\text{trend}} = 0.02$ ) (online Supplementary Table S4). Last, we examined the difference in semen quality parameters associated with the substitution of other types of meat with shellfish. Consuming shellfish instead of organ meats was associated with higher total and progressive sperm motility ( $P_{\text{trend}} < 0.001$ ). A similar association was observed when shellfish was eaten instead of white meat fishes (motile sperm,  $P_{\text{trend}} = 0.03$ ; progressive motility,  $P_{\text{trend}} = 0.004$ ) or dark meat fishes (motile sperm,  $P_{\text{trend}} = 0.03$ ; progressive motility,  $P_{\text{trend}} = 0.006$ ) (data not shown).

**Table 1.** Participant characteristics according to intakes of red meat or fish\* (Murcia Young Men's Study (n 206)) (Medians and interquartile ranges (IQR) for continuous variables; numbers of participants and percentages for categorical variables)

	Total cohort (n 206)		Red meats				P†	Fish				P†
	Median	IQR	Q1 (n 53) (lowest; 0–0.72 servings/d)		Q4 (n 51) (highest; 1.43–4.25 servings/d)			Q1 (n 51) (lowest; 0–0.53 servings/d)		Q4 (n 51) (highest; 1.25–4.66 servings/d)		
			Median	IQR	Median	IQR		Median	IQR	Median	IQR	
Age (years)	20.5	19.6–21.5	20.7	19.6–21.4	20.4	19.6–21.6	0.28	20.7	19.6–21.3	20.4	19.7–21.9	0.58
Caucasian							0.52					0.54
<i>n</i>		201		52		49			51		50	
%		97.6		100		96.1			96.2		100	
BMI (kg/m <sup>2</sup> )	23.7	21.8–25.4	23.9	22.1–25.1	23.1	21.9–25.6	0.63	24.4	22.1–25.6	23.8	22.1–25.7	0.44
Smoking							0.92					0.35
<i>n</i>		65		14		17			22		15	
%		31.9		28.0		33.3			41.5		30.6	
Testicular volume (ml)	21.0	19.5–24.0	21.0	19.0–24.3	20.5	19–23	0.23	21.0	19.8–24.3	22.3	20.0–24.5	0.22
History of cryptorchidism							0.94					0.30
<i>n</i>		4		2		0			1		0	
%		1.9		3.8		0.0			1.9		0.0	
Varicocele							0.35					0.74
<i>n</i>		14		4		4			4		2	
%		6.8		7.7		7.8			7.5		4.0	
Inguinal hernia repair‡							0.93					0.42
<i>n</i>		7		2		1			3		2	
%		3.4		3.8		2.0			5.7		4.0	
Surgical scar‡							0.12					0.77
<i>n</i>		71		14		24			18		15	
%		34.5		26.9		47.1			34.0		30.0	
Use of hormones§							0.37					0.42
<i>n</i>		1		0		0			1		0	
%		0.5		0.0		0.0			1.9		0.0	
Use of dietary supplements							0.32					0.57
<i>n</i>		29		7		7			6		6	
%		14.1		13.5		13.7			11.3		12.0	
Energy intake (kJ/d)	9543.7	7926.2–12253.3	7892.3	6202.8–9515.7	11905.2	10017.3–14798.0	<0.001	9194.3	7231.2–11526.5	11864.6	9533.7–14015.1	<0.001
Physical activity (h/week)	9.0	6.0–13.0	8.0	5.0–12.0	9.0	6.0–15.0	0.36	7.0	5.0–10.0	10.0	6.8–14.3	0.07
Television watching (h/week)	20.0	14.0–41.0	29.0	14.0–41.0	20.0	14.0–35.0	0.07	26.0	14.0–47.0	20.0	14.0–36.5	0.24
Abstinence time (h)	71.0	59.8–92.0	71.0	58.8–98.5	67.0	50.0–93.0	0.68	72.0	64.5–93.0	72.0	61.3–92.5	0.66
Time to semen analysis (min)	35.0	30.0–45.0	32.5	30.0–40.0	40.0	30.0–45.0	0.10	35.0	30.0–45.0	40.0	30.0–45.0	0.33
Time of blood draw (min)	245.0	112.5–270.0	255.0	142.5–288.8	240.0	125–270	0.73	255.0	157.5–277.5	242.5	101.3–281.3	0.23
Mediterranean diet pattern score¶	–0.2	–0.7 to 0.4	–0.3	–0.8 to 0.1	–0.2	–0.7 to 0.7	0.30	–0.7	–1.0 to 0.2	0.3	–0.1 to 1.1	<0.001
Western diet pattern score¶¶	–0.2	–0.7 to 0.5	–0.5	–0.9 to 0.0	0.2	–0.3 to 1.0	<0.001	0.0	–0.5 to 0.6	–0.2	–0.8 to 0.8	0.34
Semen volume (ml)	3.0	2.0–4.0	3.2	2.3–4.4	2.7	1.9–3.7	0.42	3.0	2.1–3.8	3.0	2.0–4.1	0.21
Sperm concentration (millions/ml)	43.4	21.9–72.3	46.2	23.0–74.1	48.7	18.7–79.3	0.96	43.4	25.6–74.7	55.6	19.1–80.5	0.81
Total sperm count (millions)	120.1	63.3–212.5	132.2	90.2–210.1	106.3	46.8–196.2	0.60	129.5	75.1–196.4	115.1	67.6–249.8	0.43
Sperm motility (%)	57.1	50.7–63.8	57.6	46.8–63.9	59.0	54.3–65.7	0.28	59.2	51–66.2	54.9	50.7–63.1	0.40
Progressive motility (%)	48.3	41.3–55.2	47.0	39.8–56.5	50.3	42.4–58.3	0.10	50.3	42.4–56.9	44.7	38.5–51.0	0.10
Morphologically normal sperm (%)	9.0	6.0–14.0	8.0	6.0–12.8	7.5	5.0–13.8	0.58	9.0	5.5–15.0	9.0	5.0–14.0	0.95
Luteinising hormone (IU/l)	4.0	2.8–5.3	3.9	2.7–5.1	4.3	3.2–5.6	0.49	4.1	2.8–5.3	3.9	2.6–5.0	0.52

A. B. Maldonado-Cárceles *et al.*

Table 1. Continued

	Total cohort (n 206)		Red meats				P†	Fish				P†
			Q1 (n 53) (lowest; 0–0.72 servings/d)		Q4 (n 51) (highest; 1.43–4.25 servings/d)			Q1 (n 51) (lowest; 0–0.53 servings/d)		Q4 (n 51) (highest; 1.25–4.66 servings/d)		
	Median	IQR	Median	IQR	Median	IQR	Median	IQR	Median	IQR		
Follicle-stimulating hormone (IU/l)	2.2	1.6–3.3	2.3	1.5–3.0	2.4	1.9–3.5	0.68	2.0	1.5–3.0	2.1	1.3–3.1	0.03
Oestradiol (pmol/l)	76.0	63.0–91.2	75.5	61.0–86.0	80.0	67.0–106.0	0.05	72.0	59.0–93.0	77.0	60.8–95.3	0.59
Calculated free testosterone (nmol/l)	13.4	10.7–17.1	13.7	10.3–16.9	15.5	11.3–18.7	0.05	12.7	10.5–17.0	14.1	10.4–18.7	0.72
Total testosterone (nmol/l)	21.2	17.1–26.6	20.9	16.6–27.2	22.0	18.8–27.2	0.11	20.1	16.6–25.5	21.4	16.6–28.1	0.52
Inhibin B (pg/ml)	193.0	147.0–246.0	181.5	145.3–239.3	184.0	145.0–227.0	0.65	203.0	146.5–249.5	201.0	157.5–266.5	0.21
Sex hormone-binding globulin (nmol/l)	30.0	23.0–39.0	29.5	21.3–40.5	30.0	24.0–36.0	0.93	27.0	20.5–39.5	29.5	24.0–37.3	0.22

Q, quartile.

\* Total red meat includes processed and unprocessed red meats and organ meat and total fish meat includes white fish meat, dark fish meat and shellfish.

† Kruskal–Wallis test for continuous variables and  $\chi^2$  test for categorical variables.

‡ Physical examination in the genital area (including lower abdomen).

§ Self-report of any use of dehydroepiandrosterone, androstenedione, creatinine, steroids or other muscle buildings.

|| Vitamins and minerals.

¶ Dietary pattern scores without total meat intake.

**Table 2.** Multivariable adjusted\* semen parameters in relation to meat intake (Murcia Young Men's Study (n 206))

Meat intake, servings/d (range)	n	Volume†		Sperm concentration†		Total sperm count†		Motile sperm‡		Progressive motility‡		Morphologically normal sperm‡	
		ml	95 % CI	10 <sup>6</sup> /ml	95 % CI	10 <sup>6</sup>	95 % CI	%	95 % CI	%	95 % CI	%	95 % CI
<b>Total meat intake</b>													
Q1 (0–1.78)	53	2.8	2.3, 3.4	33.0	24.1, 45.2	92.7	65.6, 130.8	54.9	51.5, 58.2	45.2	41.8, 48.7	7.6	6.3, 9.3
Q2 (>1.78–2.38)	51	2.6	2.1, 3.1	40.3	30.3, 53.6	103.1	75.0, 141.7	57.7	54.7, 60.8	48.6	45.5, 51.8	8.6	7.2, 10.4
Q3 (>2.38–3.23)	51	3.1	2.6, 3.7	39.0	29.5, 51.7	121.0	88.8, 165.0	56.6	53.6, 59.6	49.1	45.9, 52.2	8.7	7.3, 10.4
Q4 (>3.23–9.12)	51	2.6	2.1, 3.3	39.7	28.6, 55.0	105.1	73.3, 150.5	58.7	55.3, 62.2	49.8	46.1, 53.4	10.2	8.3, 12.5
<i>P</i> <sub>trend</sub>		0.93		0.52		0.52		0.24		0.12		0.10	
<b>Total red meat intake§</b>													
Q1 (0–0.72)	52	2.8	2.3, 3.4	38.1	28.1, 51.8	105.3	74.6, 148.6	55.8	52.6, 59.0	46.3	43.0, 49.6	8.4	6.9, 10.2
Q2 (>0.72–1.01)	53	2.8	2.3, 3.3	37.2	28.1, 49.2	103.1	75.6, 140.6	55.5	52.6, 58.4	46.1	43.1, 49.1	8.9	7.5, 10.6
Q3 (>1.01–1.42)	50	2.7	2.3, 3.3	38.2	28.7, 51.0	104.3	75.9, 143.3	58.1	55.1, 61.1	50.3	47.2, 53.3	9.3	7.8, 11.2
Q4 (>1.42–4.25)	51	2.8	2.3, 3.4	37.7	27.6, 51.6	106.4	75.2, 150.7	58.4	55.1, 61.7	49.9	46.5, 53.3	8.4	6.9, 10.2
<i>P</i> <sub>trend</sub>		0.98		0.97		0.96		0.20		0.07		0.89	
<b>Total white meat intake  </b>													
Q1 (0–0.21)	58	2.7	2.3, 3.3	37.9	28.8, 49.8	104.3	76.9, 141.5	55.2	52.4, 58.1	46.9	44.0, 49.9	9.7	8.1, 11.6
Q2 (>0.21–0.35)	43	2.9	2.4, 3.5	28.7	21.2, 38.8	83.5	59.6, 117.0	56.6	53.5, 59.7	48.0	44.8, 51.2	7.7	6.4, 9.4
Q3 (>0.35–0.57)	62	3.1	2.6, 3.6	37.3	28.9, 48.3	113.9	85.4, 152.0	56.8	54.1, 59.4	47.9	45.1, 50.6	9.2	7.8, 10.8
Q4 (>0.57–2.57)	43	2.3	1.9, 2.9	50.8	36.2, 71.1	118.5	81.4, 172.6	59.9	56.4, 63.5	50.2	46.5, 53.8	8.0	6.4, 10.0
<i>P</i> <sub>trend</sub>		0.59		0.23		0.44		0.09		0.26		0.43	
<b>Total fish intake</b>													
Q1 (0–0.53)	53	2.7	2.2, 3.2	38.0	28.6, 50.6	101.4	74.1, 138.9	56.5	53.5, 59.5	47.6	44.5, 50.7	8.0	6.7, 9.7
Q2 (>0.53–0.77)	53	2.6	2.1, 3.1	33.0	24.8, 44.0	84.4	61.6, 115.8	56.2	53.2, 59.2	46.8	43.7, 49.9	8.4	7.0, 10.1
Q3 (>0.77–1.24)	50	3.3	2.8, 4.0	34.8	26.1, 46.4	114.7	83.3, 157.7	57.9	54.8, 60.9	50.3	47.1, 53.5	8.0	6.7, 9.6
Q4 (>1.24–4.66)	50	2.6	2.1, 3.3	46.9	33.7, 65.4	124.1	86.2, 178.8	57.3	53.8, 60.8	47.9	44.3, 51.5	10.8	8.8, 13.3
<i>P</i> <sub>trend</sub>		0.52		0.51		0.33		0.58		0.53		0.13	

Q, quartile.

\* Adjusted for energy content intake, intakes of the remaining meats, dietary patterns, age, BMI, smoking, physical activity, television watching and abstinence time.

† Back-transformed to original scale.

‡ Additionally adjusted for time to start semen analysis (min).

§ Includes processed and unprocessed red meat and organ meat.

|| Includes chicken with and without skin, rabbit, quail and duck.



**Table 3.** Multivariable adjusted\* reproductive hormone levels in relation to meat intake (Murcia Young Men's Study (n 206))

Meat intake, servings/d (range)	n	LH		FSH†		Oestradiol‡		Free testosterone		Total testosterone		Inhibin B		SHBG	
		IU/l	95% CI	IU/l	95% CI	pmol/l	95% CI	nmol/l	95% CI	nmol/l	95% CI	pg/ml	95% CI	nmol/l	95% CI
<b>Total meat intake</b>															
Q1 (0–1.78)	53	4.4	3.9, 5.0	2.3	1.9, 2.7	75.2	68.6, 82.4	14.7	13.2, 16.2	22.3	20.2, 24.4	212.9	188.4, 237.4	30.6	27.4, 33.9
Q2 (>1.78–2.38)	51	3.9	3.4, 4.3	2.3	1.9, 2.7	78.5	72.4, 85.2	13.7	12.3, 15.1	21.8	19.9, 23.7	197.8	175.7, 219.8	33.8	30.9, 36.7
Q3 (>2.38–3.23)	51	4.5	4.1, 5.0	2.2	1.9, 2.6	77.9	71.7, 84.5	14.8	13.5, 16.2	22.4	20.5, 24.3	203.1	181.0, 225.2	31.1	28.2, 34.0
Q4 (>3.23–9.12)	51	4.1	3.5, 4.7	2.4	2.0, 2.9	71.7	65.1, 78.9	13.7	12.1, 15.4	20.8	18.6, 23.0	196.9	171.0, 222.8	30.6	27.2, 34.0
<i>P</i> <sub>trend</sub>		0.87		0.91		0.59		0.73		0.52		0.75		0.89	
<b>Total red meat intake‡</b>															
Q1 (0–0.72)	52	4.1	3.6, 4.6	2.2	1.8, 2.5	74.1	68, 80.7	14.4	12.9, 15.8	22	20.0, 23.9	209.9	186.8, 232.9	31.4	28.3, 34.5
Q2 (>0.72–1.01)	53	4.1	3.7, 4.6	2.2	1.9, 2.6	76.7	70.9, 83.1	14.5	13.2, 15.8	21.9	20.1, 23.8	213.8	192.4, 235.2	30.6	27.8, 33.5
Q3 (>1.01–1.42)	50	4.4	3.9, 4.9	2.2	1.9, 2.6	71.2	65.7, 77.3	12.7	11.3, 14.1	20.1	18.2, 22.0	207.5	185.7, 229.4	32.7	29.8, 35.7
Q4 (>1.42–4.25)	51	4.3	3.8, 4.9	2.6	2.2, 3.1	80.9	74, 88.4	15.4	13.8, 16.9	23.2	21.2, 25.3	179.8	155.7, 203.9	31.3	28.1, 34.6
<i>P</i> <sub>trend</sub>		0.51		0.21		0.44		0.86		0.80		0.12		0.79	
<b>Total white meat intake§</b>															
Q1 (0–0.21)	58	4.1	3.7, 4.6	2.4	2.0, 2.7	72.0	66.6, 77.9	13.7	12.4, 15.0	20.9	19.1, 22.7	215.2	194.2, 236.2	30.7	27.9, 33.5
Q2 (>0.21–0.35)	43	4.5	4.0, 5.0	2.5	2.1, 2.9	75.7	69.4, 82.7	14.1	12.7, 15.6	21.7	19.6, 23.7	190.2	166.6, 213.7	31.2	28.0, 34.3
Q3 (>0.35–0.57)	62	4.3	3.8, 4.7	2.1	1.9, 2.5	76.3	70.9, 82.2	14.8	13.5, 16.0	22.5	20.8, 24.2	195.8	176.0, 215.5	31.8	29.1, 34.4
Q4 (>0.57–2.57)	43	4.1	3.5, 4.7	2.2	1.9, 2.7	80.1	72.7, 88.2	14.3	12.7, 15.9	22.3	20.0, 24.5	208.7	182.9, 234.5	32.5	29.1, 36.0
<i>P</i> <sub>trend</sub>		0.92		0.42		0.11		0.41		0.26		0.56		0.42	
<b>Total fish meat intake</b>															
Q1 (0–0.53)	53	4.3	3.8, 4.8	2.1	1.8, 2.5	75	69.1, 81.3	14.4	13.0, 15.8	22.1	20.2, 24.0	208.5	187.2, 229.8	32.2	29.3, 35.1
Q2 (>0.53–0.77)	53	4.5	4.0, 5.0	2.7	2.3, 3.2	79.6	73.4, 86.2	14.4	13.0, 15.8	22.8	20.9, 24.7	181.2	160.0, 202.4	33.8	30.9, 36.7
Q3 (>0.77–1.24)	50	4.3	3.8, 4.8	2.4	2.1, 2.9	75	69.1, 81.5	13.7	12.3, 15.1	20.6	18.7, 22.5	202.4	180.7, 224.2	29.6	26.6, 32.5
Q4 (>1.24–4.66)	50	3.8	3.3, 4.3	1.9	1.6, 2.3	73.3	67, 80.2	14.4	12.9, 16.0	21.7	19.6, 23.8	219.8	196.0, 243.6	30.3	27.0, 33.5
<i>P</i> <sub>trend</sub>		0.27		0.54		0.63		0.82		0.49		0.46		0.21	

Meat intake and testicular function


LH, luteinising hormone; FSH, follicle-stimulating hormone; SHBG, sex hormone-binding globulin; Q, quartile.

\* Adjusted for energy content intake, intakes of the remaining meats, dietary patterns, age, BMI, smoking, physical activity, television watching and time of blood draw.

† Back-transformed to original scale.

‡ Includes processed and unprocessed red meat and organ meat.

§ Includes chicken with and without skin, rabbit, quail and duck.



**Fig. 1.** Shellfish intake in relation to progressive sperm motility among healthy young men. Estimates are adjusted for total energy intake, intakes of other meats, dietary patterns, age, BMI, smoking, physical activity, television watching, abstinence time and time to start semen analysis. Q, quartile.

We also found an inverse association between organ meat intake and sperm motility (online Supplementary Tables S6 and S8). The multivariate percentage of progressively motile sperm for men who did not consume organ meats was 50.2 (48.3–52.1)% compared with 44.8 (95% CI 42.4, 47.2)% for men who consumed organ meats ( $P=0.001$ ) (online Supplementary Table S8). LH levels for organ meat consumers were higher than those of non-consumers (4.6 *v.* 4.0 IU/l) after multivariable adjustment ( $P=0.03$ ). Levels of TT, FT and E2, as well as the TT:LH and FT:LH ratios were unrelated to organ meat intake (data not shown). In addition, unprocessed red meat intake was positively associated with LH ( $P_{\text{trend}}=0.02$ ) and SHBG ( $P_{\text{trend}}=0.001$ ) in adjusted models. No other associations were identified with semen quality parameters (online Supplementary Table S3) or reproductive hormone levels (online Supplementary Table S4).

As shellfish intake was related to higher progressive motility and lower E2 levels, and organ meat to lower progressive motility and higher LH levels, we examined whether LH or E2 were related to progressive motility in this population, and whether further adjustment for these hormones affected the association of shellfish and organ meat intake with semen quality. LH and E2 were unrelated to semen quality in this population, and additional adjustment did not change the previously observed associations (data not shown).

## Discussion

In this cross-sectional study of healthy young men, we found that total meat intake was unrelated to semen quality or reproductive hormone levels. When subgroups of meat were separately considered, however, shellfish intake was associated with higher progressive motility and organ meat intake was inversely related to this outcome. We also found association with reproductive hormones that did not explain the association with semen quality. The remaining types of meat were unrelated to semen quality parameters or reproductive hormone levels. The relevance of the observed association of shellfish

and organ meat intakes with semen quality and reproductive hormones should be further evaluated, given that these specific relations have only been evaluated in a small number of previous studies<sup>(19)</sup>.

Our findings are consistent with those of previous studies that explored the association between fish and seafood intake with semen quality. Two case-control studies among men in fertility clinics have reported significantly higher shellfish intake among normospermic men than among oligoasthenoteratospermic patients<sup>(19)</sup> and greater seafood intakes as a protective factor for asthenozoospermia<sup>(21)</sup>. Greater sperm motility has also been previously related to higher consumption of fish and other seafood<sup>(33)</sup>, while in other study no associations were found with any semen indicators<sup>(34)</sup>. Afeiche *et al.*<sup>(22)</sup> reported a positive association between intake of dark meat fish intake and total sperm count as well as a positive relation between white meat fish intake and morphologically normal sperm among men presenting to a fertility clinic but not among healthy young men<sup>(19)</sup>.

Shellfish is an important source of PUFA<sup>(35)</sup>, and previous work has related higher consumption of *n*-3 PUFA to better sperm motility<sup>(36)</sup>. However, because we did not observe an association between dark meat fish intake and semen quality in this study, and previous work in this cohort has not found associations between intakes of *n*-3 and semen quality parameters or reproductive hormones<sup>(12,24)</sup>, intake of *n*-3 PUFA is unlikely to explain our findings. Instead, the relation between shellfish intake with sperm motility may be due to the intake of micronutrients highly concentrated in shellfish such as Zn<sup>(35)</sup>. Zn is involved in male reproductive function including spermatogenesis, sperm maturation and sperm activation<sup>(37)</sup>. A recent meta-analysis showed that Zn supplementation in infertile men was associated with better semen quality parameters including sperm motility<sup>(37)</sup>.

To our knowledge, only three previous studies have addressed the association between intake of organ meat and semen parameters. Organ meat intake has been previously hypothesised to lower semen quality<sup>(19)</sup>. Contrary to our findings, a similar study among young men in the USA found organ consumption to be positively related to sperm motility<sup>(19)</sup>. Previous studies among infertile men have not found a relationship between organ consumption and sperm motility<sup>(19,22)</sup>. As literature is scarce and inconsistent, this association may represent a chance finding and more studies are needed to clarify this discrepancy.

Several limitations should be discussed. The cross-sectional design is not strong to distinguish causal from non-causal relationships and the small sample size could result in missing true associations. However, results of the semen analysis were unknown to participants, essentially blinding them to the study outcome, and thus decreasing the likelihood of reverse causation. Diet was assessed using an FFQ and like all other diet assessment tools, FFQ are subject to measurement error. However, this questionnaire has been validated previously in Spanish populations<sup>(27)</sup>. In addition, FFQ are known to be better at ranking than at estimating exact intakes. Our analytic strategy of making comparisons between quantiles of intake rather than relying on continuous measures of intake is not only better aligned with this characteristic of FFQ but also protects



against the influence of individuals with very high intakes, thus resulting in conservative estimates of the associations examined. Moreover, as the consumption of shellfish and organ meats was very low (3–5%) and the multiple comparisons developed in the statistical analysis, we cannot rule out the possibility of residual confounding or chance findings that could explain the associations reported. However, we adjusted for a large number of potential confounders, including known predictors of semen quality as well as lifestyle factors associated with meat intake, including other dietary behaviours as captured by data-derived dietary patterns. Finally, although the homogeneity of study participants may have increased the internal validity of the study, it may limit the generalisability of our results to men facing difficulties with fertility.

In conclusion, we found no relation of total meat intake nor intake of major types of meat (red meats, white meats and fish) with markers of testicular function. Nevertheless, analyses of subgroups of meat suggest that intake of shellfish may be beneficial for sperm motility, whereas the intake of organ meats may have the opposite relation. Due to the scarcity of literature in this population and in specific types of meats, additional research is needed to confirm or refute these findings.

### Acknowledgements

This work was supported by the Seneca Foundation, Regional Agency of Science and Technology, grant no. 08808/PI/08; Ministerio de Ciencia e Innovación, Instituto Carlos III (FIS), grant no. PI10/00985; National Institutes of Health, grant no. P30DK046200.

A. M. T.-C., J. M. and N. J. were involved in study conception and study design. J. M. was involved in study execution and acquisition of data. A. B. M.-C. performed the statistical analysis. A. B. M.-C., L. M.-A., J. E. C. and J. J. A.-G. contributed to data analysis and interpretation. A. B. M.-C., L. M.-A. and J. E. C. drafted the manuscript. J. V. and N. J. were involved in critical revision for important intellectual content. All authors provided substantial intellectual contributions and approved the final version of the manuscript.

The authors declare that there are no conflicts of interest.

### Supplementary material

For supplementary material/s referred to in this article, please visit <https://doi.org/10.1017/S0007114518003458>

### References

- Swan SH, Elkin EP & Fenster L (2000) The question of declining sperm density revisited: an analysis of 101 studies published 1934–1996. *Environ Health Perspect* **108**, 961–966.
- Kumar N & Singh AK (2015) Trends of male factor infertility, an important cause of infertility: a review of literature. *J Hum Reprod Sci* **8**, 191–196.
- Levine H, Jørgensen N, Martino-Andrade A, *et al.* (2017) Temporal trends in sperm count: a systematic review and meta-regression analysis. *Hum Reprod Update* **23**, 646–659.
- Virtanen HE, Jørgensen N & Toppari J (2017) Semen quality in the 21(st) century. *Nat Rev Urol* **14**, 120–130.
- Nordkap L, Joensen UN, Blomberg Jensen M, *et al.* (2012) Regional differences and temporal trends in male reproductive health disorders: semen quality may be a sensitive marker of environmental exposures. *Mol Cell Endocrinol* **355**, 221–230.
- Povey AC, Clyma JA, McNamee R, *et al.* (2012) Modifiable and non-modifiable risk factors for poor semen quality: a case-referent study. *Hum Reprod* **27**, 2799–2806.
- Centers for Disease Control and Prevention (2014) *National Public Health Action Plan for the Detection, Prevention, and Management of Infertility*. Atlanta: CDC.
- Mahe G, Carsin M, Zeeny M, *et al.* (2011) Dietary pattern, a modifiable risk factor that can be easily assessed for atherosclerosis vascular disease prevention in clinical practice. *Public Health Nutr* **14**, 319–326.
- Jensen TK, Swan SH, Skakkebaek NE, *et al.* (2010) Caffeine intake and semen quality in a population of 2,554 young Danish men. *Am J Epidemiol* **171**, 883–891.
- Mínguez-Alarcón L, Mendiola J, López-Espín JJ, *et al.* (2012) Dietary intake of antioxidant nutrients is associated with semen quality in young university students. *Hum Reprod* **27**, 2807–2814.
- Jensen TK, Heitmann BL, Blomberg Jensen M, *et al.* (2013) High dietary intake of saturated fat is associated with reduced semen quality among 701 young Danish men from the general population. *Am J Clin Nutr* **97**, 411–418.
- Chavarro JE, Mínguez-Alarcón L, Mendiola J, *et al.* (2014) *Trans* fatty acid intake is inversely related to total sperm count in young healthy men. *Hum Reprod* **29**, 429–440.
- Giahi L, Mohammadmoradi S, Javidan A, *et al.* (2016) Nutritional modifications in male infertility: a systematic review covering 2 decades. *Nutr Rev* **74**, 118–130.
- Salas-Huetos A, Bulló M & Salas-Salvadó J (2017) Dietary patterns, foods and nutrients in male fertility parameters and fecundability: a systematic review of observational studies. *Hum Reprod Update* **23**, 371–389.
- Swan SH, Liu F, Overstreet JW, *et al.* (2007) Semen quality of fertile US males in relation to their mothers' beef consumption during pregnancy. *Hum Reprod* **22**, 1497–1502.
- Gasull M, Bosch de Basea M, Puigdomènech E, *et al.* (2011) Empirical analyses of the influence of diet on human concentrations of persistent organic pollutants: a systematic review of all studies conducted in Spain. *Environ Int* **37**, 1226–1235.
- Attaman JA, Toth TL, Furtado J, *et al.* (2012) Dietary fat and semen quality among men attending a fertility clinic. *Hum Reprod* **27**, 1466–1474.
- Vested A, Giwercman A, Bonde JP, *et al.* (2014) Persistent organic pollutants and male reproductive health. *Asian J Androl* **16**, 71–80.
- Afeiche MC, Williams PL, Gaskins AJ, *et al.* (2014) Meat intake and reproductive parameters among young men. *Epidemiology* **25**, 323–330.
- Mendiola J, Torres-Cantero AM, Moreno-Grau JM, *et al.* (2009) Food intake and its relationship with semen quality: a case-control study. *Fertil Steril* **91**, 812–818.
- Eslamian G, Amirjannati N, Rashidkhani B, *et al.* (2012) Intake of food groups and idiopathic asthenozoospermia: a case-control study. *Hum Reprod* **27**, 3328–3336.
- Afeiche MC, Gaskins AJ, Williams PL, *et al.* (2014) Processed meat intake is unfavorably and fish intake favorably associated with semen quality indicators among men attending a fertility clinic. *J Nutr* **144**, 1091–1098.
- Cutillas-Tolín A, Mínguez-Alarcón L, Mendiola J, *et al.* (2015) Mediterranean and Western dietary patterns are related to

- markers of testicular function among healthy men. *Hum Reprod* **30**, 2945–2955.
24. Mínguez-Alarcón L, Chavarro JE, Mendiola J, *et al.* (2017) Fatty acid intake in relation to reproductive hormones and testicular volume among young healthy men. *Asian J Androl* **19**, 184–190.
  25. Vioque J & Gonzalez L (1991) Validity of a food frequency questionnaire (preliminary results). *Eur J Cancer Prev* **1**, 19–20.
  26. Vioque J, Navarrete-Muñoz EM, Gimenez-Monzó D, *et al.* (2013) Reproducibility and validity of a food frequency questionnaire among pregnant women in a Mediterranean area. *Nutr J* **12**, 26.
  27. World Health Organization (2010) *WHO Laboratory Manual for the Examination and Processing of Human Semen*, 5th ed. Geneva: WHO.
  28. Menkveld R, Stander FS, Kotze TJ, *et al.* (1990) The evaluation of morphological characteristics of human spermatozoa according to stricter criteria. *Hum Reprod* **5**, 586–592.
  29. Asklund C, Jørgensen N, Skakkebaek NE, *et al.* (2007) Increased frequency of reproductive health problems among fathers of boys with hypospadias. *Hum Reprod* **22**, 2639–2646.
  30. Vermeulen A, Verdonck L & Kaufman JM (1999) A critical evaluation of simple methods for the estimation of free testosterone in serum. *J Clin Endocrinol Metab* **84**, 3666–3672.
  31. Soper DS (2018) Free statistics calculators – *a-priori* sample size calculator for multiple regression. <http://www.danielsoper.com/statcalc> (accessed January 2018).
  32. Hu FB, Rimm EB, Stampfer MJ, *et al.* (2000) Prospective study of major dietary patterns and risk of coronary heart disease in men. *Am J Clin Nutr* **72**, 912–921.
  33. Vujkovic M, de Vries JH, Dohle GR, *et al.* (2009) Associations between dietary patterns and semen quality in men undergoing IVF/ICSI treatment. *Hum Reprod* **24**, 1304–1312.
  34. Braga DP, Halpern G, Figueira Rde C, *et al.* (2012) Food intake and social habits in male patients and its relationship to intracytoplasmic sperm injection outcomes. *Fertil Steril* **97**, 53–59.
  35. Dong FM (2009) *The Nutritional Value of Shellfish*. Washington: University of Washington.
  36. Safarinejad MR (2011) Effect of omega-3 polyunsaturated fatty acid supplementation on semen profile and enzymatic anti-oxidant capacity of seminal plasma in infertile men with idiopathic oligoasthenoteratospermia: a double-blind, placebo-controlled, randomised study. *Andrologia* **43**, 38–47.
  37. Zhao J, Dong X, Hu X, *et al.* (2016) Zinc levels in seminal plasma and their correlation with male infertility: a systematic review and meta-analysis. *Sci Rep* **6**, 22386.